

THE MONOIDAL CENTRE AS A LIMIT

To Aurelio Carboni for his sixtieth birthday

ROSS STREET

ABSTRACT. The centre of a monoidal category is a braided monoidal category. Monoidal categories are monoidal objects (or pseudomonoids) in the monoidal bicategory of categories. This paper provides a universal construction in a braided monoidal bicategory that produces a braided monoidal object from any monoidal object. Some properties and sufficient conditions for existence of the construction are examined.

1. Introduction

During question time after a talk [St2] at the Fields Institute, Peter Schauenburg asked whether the centre construction on a monoidal category (see [JS]) would fit into the general framework of [DMS] that I was describing. At the time I could not see how to do it. Reinforced by Peter's interest, the question stayed with me. During preparation of the paper [St3] on descent theory, intended for a publication arising from the same Fields Institute workshop, the answer began to dawn on me. Another topic at the top of my mind recently (in work with Michael Batanin and Alexei Davydov) has been Hochschild cohomology, and this too turns out to be relevant.

2. The centre of a monoidal object

In any monoidal bicategory \mathcal{M} , with tensor product \otimes and unit I , we use the terms *pseudomonoid* and *monoidal object* for an object A equipped with a binary multiplication $m : A \otimes A \longrightarrow A$ and a unit $j : I \longrightarrow A$ which are associative and unital up to coherent invertible 2-cells. A *monoidal morphism* $f : A \longrightarrow A'$ is a morphism equipped with coherent 2-cells $m \circ (f \otimes f) \Longrightarrow f \circ m$ and $j \Longrightarrow f \circ j$. The monoidal morphism is called *strong* when the coherent 2-cells are both invertible. A *monoidal 2-cell* is one compatible with these last coherent 2-cells. With the obvious compositions, this defines a bicategory $\mathbf{Mon} \mathcal{M}$ of pseudomonoids in \mathcal{M} . For example, if \mathcal{M} is the cartesian-monoidal 2-category \mathbf{Cat} of categories, functors and natural transformations then $\mathbf{Mon} \mathcal{M}$ is the 2-category \mathbf{MonCat} of monoidal categories, monoidal functors and monoidal natural transformations as defined in [EK]. We now suppose \mathcal{M} is braided. In fact, by the coherence result of

Received by the editors 2003-05-06 and, in revised form, 2003-07-31.

Published on 2004-12-05.

2000 Mathematics Subject Classification: 18D10.

Key words and phrases: braiding, centre, pseudofunctor, descent.

© Ross Street, 2004. Permission to copy for private use granted.

[GPS], we suppose \mathcal{M} is a braided Gray monoid in the sense of [DS]. The braiding for \mathcal{M} is denoted by

$$c_{X,Y} : X \otimes Y \xrightarrow{\sim} Y \otimes X$$

for objects X and Y , together with pseudonaturality isomorphisms $c_{f,g}$ for morphisms f and g .

For any monoidal object A of \mathcal{M} , a morphism $u : U \longrightarrow A$ is called a *centre piece* when it is equipped with an invertible 2-cell

$$\begin{array}{ccc} U \otimes A & \xrightarrow{c_{U,A}} & A \otimes U \\ u \otimes 1_A \downarrow & \xRightarrow{\gamma} & \downarrow 1_A \otimes u \\ A \otimes A & & A \otimes A \\ & \xrightarrow{m} & A \end{array}$$

such that the following equality holds.

$$\begin{array}{c} \begin{array}{ccccc} U \otimes A \otimes A & \xrightarrow{c_{U,A \otimes A}} & A \otimes A \otimes U & & \\ u \otimes 1 \otimes 1 \swarrow & 1 \otimes m \searrow & \cong & m \otimes 1 \swarrow & 1 \otimes 1 \otimes u \searrow \\ A \otimes A \otimes A & & U \otimes A & \xrightarrow{c_{U,A}} & A \otimes U & \cong & A \otimes A \otimes A \\ 1 \otimes m \searrow & u \otimes 1 \downarrow & \xRightarrow{\gamma} & 1 \otimes u \downarrow & m \otimes 1 \swarrow & & \\ A \otimes A & & A \otimes A & & A \otimes A & & \\ m \otimes 1 \swarrow & & m \downarrow & & m \downarrow & & \\ A \otimes A & & A & & A & & \\ & \xrightarrow{m} & & \xrightarrow{m} & & & \end{array} \\ \parallel \\ \begin{array}{ccccc} U \otimes A \otimes A & \xrightarrow{c_{U,A \otimes A}} & A \otimes A \otimes U & & \\ u \otimes 1 \otimes 1 \swarrow & & \cong & & 1 \otimes 1 \otimes u \searrow \\ A \otimes A \otimes A & & A \otimes U \otimes A & & A \otimes A \otimes A \\ \gamma \otimes 1 \xRightarrow{\cong} & 1 \otimes u \otimes 1 \downarrow & & 1 \otimes \gamma \xRightarrow{\cong} & \\ A \otimes A \otimes A & & A \otimes A \otimes A & & A \otimes A \otimes A \\ m \otimes 1 \swarrow & & m \otimes 1 \swarrow & & 1 \otimes m \searrow \\ A \otimes A & & A \otimes A & & A \otimes A \\ & \xrightarrow{m} & & \xrightarrow{m} & \end{array} \end{array}$$

A *morphism* $\sigma : u \Longrightarrow v : U \longrightarrow A$ of *centre pieces* is a 2-cell $\sigma : u \Longrightarrow v$ such that

$$\gamma(m \circ (\sigma \otimes 1_A)) = (m \circ (1_A \otimes \sigma) \circ c_{U,A})\gamma .$$

We write $\mathcal{CP}(U, A)$ for the category of centre pieces so obtained. Using the pseudonaturality of the braiding for \mathcal{M} , we see that we have a pseudofunctor

$$\mathcal{CP}(-, A) : \mathcal{M}^{\text{op}} \longrightarrow \mathbf{Cat}$$

defined on morphisms $f : V \longrightarrow U$ by composition; that is, the functor

$$\mathcal{CP}(f, A) : \mathcal{CP}(U, A) \longrightarrow \mathcal{CP}(V, A)$$

takes a centre piece u with γ to $u \circ f$ with the 2-cell obtained by pasting the square containing the pseudonaturality isomorphism $c_{f,1_A}$ onto the top of the pentagon containing γ .


The *centre of A* is a birepresenting object $\mathcal{Z}A$ (in the sense of [St0]) for the pseudo-functor $\mathcal{CP}(-, A)$. This means we have a centre piece $i : \mathcal{Z}A \longrightarrow A$, composition with which induces an equivalence of categories

$$\mathcal{M}(U, \mathcal{Z}A) \simeq \mathcal{CP}(U, A) .$$

It follows that the centre of A is unique up to equivalence if it exists.

2.1. PROPOSITION. *The centre $\mathcal{Z}A$ of a monoidal object A is a braided monoidal object in the sense of [DS], and the morphism $i : \mathcal{Z}A \longrightarrow A$ is strong monoidal.*


PROOF. The composite $\mathcal{Z}A \otimes \mathcal{Z}A \xrightarrow{i \otimes i} A \otimes A \xrightarrow{m} A$ equipped with the 2-cell



is a centre piece. So, up to a unique invertible 2-cell, there is a morphism

$$m : \mathcal{Z}A \otimes \mathcal{Z}A \longrightarrow \mathcal{Z}A$$

and an invertible 2-cell $i \circ m \cong m \circ (i \otimes i)$ compatible with the 2-cells of the two centre pieces. Also $j : I \longrightarrow A$, equipped with the obvious 2-cell, is a centre piece and so induces a morphism $j : I \longrightarrow \mathcal{Z}A$, with $i \circ j \cong j$. Largish pasting diagrams prove that $\mathcal{Z}A$ becomes monoidal with $i : \mathcal{Z}A \longrightarrow A$ strong monoidal. The braiding for $\mathcal{Z}A$ is the invertible 2-cell



whose composite with $i : \mathcal{Z}A \longrightarrow A$ is the pasting composite below.

$$\begin{array}{ccccc}
 \mathcal{Z}A \otimes \mathcal{Z}A & \xrightarrow{c_{\mathcal{Z}A, \mathcal{Z}A}} & \mathcal{Z}A \otimes \mathcal{Z}A & & \\
 \downarrow m & \searrow 1 \otimes i & \downarrow C_{1,i} \cong & \swarrow i \otimes 1 & \downarrow m \\
 \mathcal{Z}A & \xrightarrow{c_{\mathcal{Z}A, A}} & A \otimes \mathcal{Z}A & \xrightarrow{c_{A, \mathcal{Z}A}} & \mathcal{Z}A \\
 \cong \downarrow & \cong \downarrow & \downarrow 1 \otimes i & \downarrow 1 \otimes i & \cong \downarrow \\
 \mathcal{Z}A & \xrightarrow{i \otimes 1} & A \otimes A & \xrightarrow{\gamma} & A \otimes A \\
 \downarrow i & \downarrow m & \downarrow m & \downarrow m & \downarrow i \\
 A & & A & & A
 \end{array}$$

The braiding axioms follow from the defining property of a centre piece. ■

The following two propositions are routinely proved.

2.2. PROPOSITION. *If $F : \mathcal{M} \longrightarrow \mathcal{N}$ is a braided monoidal pseudofunctor and $u : U \longrightarrow A$ is a centre piece for the monoidal object A in \mathcal{M} then $Fu : FU \longrightarrow FA$ is canonically a centre piece for the monoidal object FA in \mathcal{N} . There is a canonical comparison morphism $F\mathcal{Z}A \longrightarrow \mathcal{Z}FA$ provided the centres of A and FA exist.*

2.3. PROPOSITION. *The pseudofunctor $\mathcal{CP}(-, A) : \mathcal{M}^{\text{op}} \longrightarrow \mathbf{Cat}$ preserves all weighted bicategorical limits that exist in \mathcal{M}^{op} and, that as colimits in \mathcal{M} , are preserved by $- \otimes A$ and $- \otimes A \otimes A$.*

3. Existence

In this section we shall look at conditions on the braided monoidal bicategory \mathcal{M} for monoidal centres to exist. Because of Proposition 2.3, we expect $\mathcal{CP}(-, A) : \mathcal{M}^{\text{op}} \longrightarrow \mathbf{Cat}$ to be birepresentable when each $- \otimes A$ preserves colimits and a “special birepresentability theorem” applies to \mathcal{M} .

Recall from [DS] that a monoidal bicategory \mathcal{M} is called *left [right] closed* when, for each object B , the pseudofunctor $- \otimes B : \mathcal{M} \longrightarrow \mathcal{M}$ [$B \otimes - : \mathcal{M} \longrightarrow \mathcal{M}$] has a right biadjoint (and so preserves bicategorical colimits). We call \mathcal{M} *closed* when it is both left and right closed; we denote the right biadjoint of $- \otimes B$ by $[B, -] : \mathcal{M} \longrightarrow \mathcal{M}$ and we have a family of equivalences

$$\mathcal{M}(A \otimes B, C) \cong \mathcal{M}(A, [B, C]) ,$$

pseudonatural in each variable, called the *closedness equivalences*. Taking $A = [B, C]$, we find an evaluation morphism $\text{ev} : [B, C] \otimes B \longrightarrow C$ that, up to isomorphism, is taken to the identity by the closedness equivalence.

Assume \mathcal{M} is braided and left closed. It follows that \mathcal{M} is closed. From any monoidal object A we shall construct a Hochschild-like truncated pseudo-cosimplicial object $\mathcal{C}A$:

$$A \begin{array}{c} \xrightarrow{\partial_0} \\ \xrightarrow{\partial_1} \end{array} [A, A] \begin{array}{c} \xrightarrow{\partial_0} \\ \xrightarrow{\partial_1} \\ \xrightarrow{\partial_2} \end{array} [A \otimes A, A]$$

as follows. The morphism $\partial_0 : A \longrightarrow [A, A]$ corresponds under the closedness equivalence to $m : A \otimes A \longrightarrow A$. The morphism $\partial_1 : A \longrightarrow [A, A]$ corresponds under the closedness equivalence to the composite $A \otimes A \xrightarrow{c_{A,A}} A \otimes A \xrightarrow{m} A$. The morphisms

$$\partial_0, \partial_1, \partial_2 : [A, A] \longrightarrow [A \otimes A, A]$$

correspond under the closedness equivalence to the morphisms

$$[A, A] \otimes A \otimes A \begin{array}{c} \xrightarrow{m \circ (\text{ev} \otimes 1_A)} \\ \xrightarrow{\text{ev} \circ (1 \otimes m)} \\ \xrightarrow{m \circ (1_A \otimes \text{ev}) \circ (c_{[A,A],A} \otimes 1_A)} \end{array} A .$$

One easily finds the coherent invertible 2-cells

$$\partial_0 \circ \partial_0 \cong \partial_1 \circ \partial_0, \quad \partial_0 \circ \partial_1 \cong \partial_2 \circ \partial_0, \quad \partial_2 \circ \partial_1 \cong \partial_1 \circ \partial_1 .$$

3.1. PROPOSITION. *The bicategorical limit of the pseudo-cosimplicial diagram $\mathcal{C}A$ is the centre of A .*

PROOF. The proof is by transport across the closedness equivalences. ■

It is shown in [St0] how to construct this pseudo-descent-like limit in a bicategory \mathcal{M} that admits finite products, iso-inserters, and cotensoring with the arrow category **2**.

3.2. COROLLARY. *In any finitely complete, closed, braided monoidal bicategory, every monoidal object has a centre. Any braided monoidal pseudofunctor that is strong closed and finite-limit preserving preserves centres.*

Examples of such bicategories abound. Let \mathcal{A} be any (small) braided promonoidal 2-category (in particular, \mathcal{A} could be a braided monoidal category). Take \mathcal{M} to be the 2-category $[\mathcal{A}, \mathbf{Cat}]$ of 2-functors from \mathcal{A} to \mathbf{Cat} . This is a complete and cocomplete 2-category and so is also complete and cocomplete as a bicategory. It becomes closed monoidal under the Day convolution tensor product defined by the coends

$$(F \otimes G)A = \int^{B,C} P(B, C; A) \times FB \times GC$$

in \mathbf{Cat} . The braiding is induced by that on \mathcal{A} and the symmetry on \mathbf{Cat} . A good example of an appropriate \mathcal{A} is provided by the category of automorphisms of a groupoid \mathcal{G} as described in Section 7, Example 9 of [DS].

Alternatively we could take \mathcal{M} to be the 2-category $\text{Hom}(\mathcal{A}, \mathbf{Cat})$ of pseudofunctors from \mathcal{A} to \mathbf{Cat} . This is complete and cocomplete as a bicategory. It becomes a closed monoidal bicategory under the convolution tensor product defined by the pseudocoends

$$(F \otimes G)A = \int_{ps}^{B,C} P(B, C; A) \times FB \times GC$$

in \mathbf{Cat} . Again, the braiding is induced by that on \mathcal{A} and the symmetry on \mathbf{Cat} .

3.3. REMARK. After submission of this paper, Steve Lack provided some helpful insights for which I am very grateful and will now explain. We begin with the observation that the centre piece category $\mathcal{CP}(U, A)$ is precisely the descent category of the truncated pseudocosimplicial diagram

$$\mathcal{M}(U, A) \rightrightarrows \mathcal{M}(U \otimes A, A) \rightrightarrows \mathcal{M}(U \otimes A \otimes A, A) .$$

Allowing U to vary, we obtain the pseudofunctor $\mathcal{CP}(-, A) : \mathcal{M}^{\text{op}} \longrightarrow \mathbf{Cat}$ as the descent object of a truncated pseudocosimplicial object in $\text{Hom}(\mathcal{M}^{\text{op}}, \mathbf{Cat})$. If \mathcal{M} is closed, my Hochschild description of the centre is recaptured. Returning to any braided monoidal bicategory \mathcal{M} , we have the convolution braided monoidal structure on $\text{Hom}(\mathcal{M}^{\text{op}}, \mathbf{Cat})$. The diagram for the descent object $\mathcal{CP}(-, A)$ lifts to the bicategory $\text{MonHom}(\mathcal{M}^{\text{op}}, \mathbf{Cat})$ of monoidal objects for the convolution; the objects of $\text{MonHom}(\mathcal{M}^{\text{op}}, \mathbf{Cat})$ are (weakly) monoidal pseudofunctors and the morphisms are monoidal pseudonatural transformations. It follows that $\mathcal{CP}(-, A)$ and $\mathcal{CP}(-, A) \longrightarrow \mathcal{M}(-, A)$ are in $\text{MonHom}(\mathcal{M}^{\text{op}}, \mathbf{Cat})$; moreover, $\mathcal{CP}(-, A) : \mathcal{M}^{\text{op}} \longrightarrow \mathbf{Cat}$ is a braided pseudofunctor between braided monoidal bicategories. The largish diagrams proving the last sentence may even be more readily comprehensible than those referred to in the proof of Proposition 2.1. If $\mathcal{CP}(-, A)$ is representable by some object $\mathcal{Z}A$ then, because the Yoneda embedding $\mathcal{M} \longrightarrow \text{Hom}(\mathcal{M}^{\text{op}}, \mathbf{Cat})$ is strong monoidal, we deduce that $\mathcal{Z}A$ is braided monoidal.

References

- [DMS] B. Day, P. McCrudden and R. Street, Dualizations and antipodes, *Applied Categorical Structures* **11** (2003) 229–260.
- [DS] B. Day and R. Street, Monoidal bicategories and Hopf algebroids, *Advances in Math.* **129** (1997) 99–157.
- [EK] S. Eilenberg and G.M. Kelly, Closed categories, *Proc. Conf. Categorical Algebra at La Jolla 1965* (Springer-Verlag, Berlin 1966) 421–562.
- [GPS] R. Gordon, A.J. Power and R. Street, Coherence for tricategories, *Memoirs Amer. Math. Soc.* **117** (1995) #558.

- [JS] A. Joyal and R. Street, Tortile Yang-Baxter operators in tensor categories, *J. Pure Appl. Algebra* **71** (1991) 43–51.
- [St0] R. Street, Fibrations in bicategories, *Cahiers topologie et géométrie différentielle* **21** (1980) 111–160; **28** (1987) 53–56.
- [St1] R. Street, The quantum double and related constructions, *J. Pure Appl. Algebra* **132** (1998) 195–206.
- [St2] R. Street, Formal representation theory, Talk at the “Workshop on Categorical Structures for Descent and Galois Theory, Hopf Algebras and Semiaabelian Categories” (Fields Institute, 24 September, 2002); slides and audio at http://www.fields.utoronto.ca/audio/02-03/galois_and_hopf/street/.
- [St3] R. Street, Categorical and combinatorial aspects of descent theory, *Applied Categorical Structures* (to appear; also see math.CT/0303175).

Centre of Australian Category Theory,
Macquarie University, N.S.W. 2109, Australia
Email: street@maths.mq.edu.au

This article may be accessed via WWW at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/13/13/13-13.dvi>, [ps](ftp://ftp.tac.mta.ca/pub/tac/html/volumes/13/13/13-13.ps)

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

The method of distribution of the journal is via the Internet tools `WWW/ftp`. The journal is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION. Individual subscribers receive (by e-mail) abstracts of articles as they are published. Full text of published articles is available in .dvi, Postscript and PDF. Details will be e-mailed to new subscribers. To subscribe, send e-mail to `tac@mta.ca` including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, `rrosebrugh@mta.ca`.

INFORMATION FOR AUTHORS. The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{\LaTeX}2_{\text{e}}$ is the preferred flavour. $\text{T}_{\text{E}}\text{X}$ source of articles for publication should be submitted by e-mail directly to an appropriate Editor. They are listed below. Please obtain detailed information on submission format and style files from the journal's WWW server at `http://www.tac.mta.ca/tac/`. You may also write to `tac@mta.ca` to receive details by e-mail.

EDITORIAL BOARD.

Michael Barr, McGill University: `barr@barrs.org`, *Associate Managing Editor*

Lawrence Breen, Université Paris 13: `breen@math.univ-paris13.fr`

Ronald Brown, University of Wales Bangor: `r.brown@bangor.ac.uk`

Jean-Luc Brylinski, Pennsylvania State University: `jlb@math.psu.edu`

Aurelio Carboni, Università dell'Insubria: `aurelio.carboni@uninsubria.it`

Valeria de Paiva, Palo Alto Research Center: `paiva@parc.xerox.com`

Martin Hyland, University of Cambridge: `M.Hyland@dpms.cam.ac.uk`

P. T. Johnstone, University of Cambridge: `ptj@dpms.cam.ac.uk`

G. Max Kelly, University of Sydney: `maxk@maths.usyd.edu.au`

Anders Kock, University of Aarhus: `kock@imf.au.dk`

Stephen Lack, University of Western Sydney: `s.lack@uws.edu.au`

F. William Lawvere, State University of New York at Buffalo: `wlawvere@buffalo.edu`

Jean-Louis Loday, Université de Strasbourg: `loday@math.u-strasbg.fr`

Ieke Moerdijk, University of Utrecht: `moerdijk@math.uu.nl`

Susan Niefield, Union College: `niefiels@union.edu`

Robert Paré, Dalhousie University: `pare@mathstat.dal.ca`

Robert Rosebrugh, Mount Allison University: `rrosebrugh@mta.ca`, *Managing Editor*

Jiri Rosicky, Masaryk University: `rosicky@math.muni.cz`

James Stasheff, University of North Carolina: `jds@math.unc.edu`

Ross Street, Macquarie University: `street@math.mq.edu.au`

Walter Tholen, York University: `tholen@mathstat.yorku.ca`

Myles Tierney, Rutgers University: `tierney@math.rutgers.edu`

Robert F. C. Walters, University of Insubria: `robert.walters@uninsubria.it`

R. J. Wood, Dalhousie University: `rjwood@mathstat.dal.ca`