

NUCLEI OF CATEGORIES WITH TENSOR PRODUCTS

ALEXEI DAVYDOV

ABSTRACT. Following the analogy between algebras (monoids) and monoidal categories the construction of nucleus for non-associative algebras is simulated on the categorical level.

Nuclei of categories of modules are considered as an example.

1. Introduction

The property

$$xJ(y, z, w) - J(xy, z, w) + J(x, yz, w) - J(x, y, zw) + J(x, y, z)w = 0 \quad (1)$$

of the *associator* $J(x, y, z) = x(yz) - (xy)z$ guarantees that the subspaces

$$N_l(A) = \{a \in A : J(a, x, y) = 0 \forall x, y \in A\}, \quad (2)$$

$$N_m(A) = \{a \in A : J(x, a, y) = 0 \forall x, y \in A\}, \quad (3)$$

$$N_r(A) = \{a \in A : J(x, y, a) = 0 \forall x, y \in A\} \quad (4)$$

of an algebra A are associative subalgebras (see for example [4]). We call these subalgebras *left*, *middle* and *right nucleus* respectively.

In section 2.1 we categorify these constructions by defining left, middle and right nuclei $\mathcal{N}_l(\mathcal{G}), \mathcal{N}_m(\mathcal{G}), \mathcal{N}_r(\mathcal{G})$ for a category \mathcal{G} with a tensor product (*magmoidal* category). The tensor product transports to the nuclei. Moreover there is a natural choice of associativity constraint making them *semigroupal* categories (the difference between semigroupal and monoidal is the lack of unit object).

The construction resembles the well-know categorification of the centre of an associative algebra (the so called *monoidal centre* [3]). In both cases the categorifications are categories of pairs. The first component of a pair is an object of the original category when the second should be seen as a specialisation of the appropriate constraint (braiding in the case of the monoidal centre, associativity in the case of nucleus). Coherence then

The work on the paper was supported by Australian Research Council grant DP00663514. The author thanks members of the Australian Category Seminar for stimulating discussions and the referees for numerous corrections and suggestions. Special thanks are due to R. Street for invaluable support during the work on the paper.

Received by the editors 2007-03-12 and, in revised form, 2007-10-08.

Transmitted by RJ Wood. Published on 2007-10-11.

2000 Mathematics Subject Classification: 18D10.

Key words and phrases: monoidal categories, monoidal functors, quasi-bialgebras.

© Alexei Davyдов, 2007. Permission to copy for private use granted.

used to define tensor product of pairs and to check the desired properties of resulting monoidal categories.

Categorical analogues of the inclusions $N_l(A), N_m(A), N_r(A) \subset A$ are tensor product preserving (forgetful) functors $\mathcal{N}_l(\mathcal{G}), \mathcal{N}_m(\mathcal{G}), \mathcal{N}_r(\mathcal{G}) \longrightarrow \mathcal{G}$.

If an algebra A is already associative, its nuclei coincide with it. For a semigroupal category \mathcal{G} we only get semigroupal functors $\mathcal{G} \longrightarrow \mathcal{N}_l(\mathcal{G}), \mathcal{N}_m(\mathcal{G}), \mathcal{N}_r(\mathcal{G})$, which split the corresponding forgetful functor. Semigroupal structures on \mathcal{G} are in 1-to-1 correspondence with splittings of the forgetful functor.

If a (non-associative) algebra A has a unit (an element $e \in A$ such that $ex = xe = x$ for any $x \in A$) then the unit belongs to all nuclei $N_e(A)$ making them associative unital. In section 2.5 we modify the constructions of nuclei for the case of a magmoidal category with unit. The modified constructions give monoidal categories $\mathcal{N}_l^1(\mathcal{G}), \mathcal{N}_m^1(\mathcal{G}), \mathcal{N}_r^1(\mathcal{G})$.

Constructions of self-symmetries of categorical nuclei which we describe in section 2.9 do not have any analogues in algebra. Here we associate, to any natural transformation of the tensor product functor on \mathcal{G} , monoidal autoequivalences of the nuclei of \mathcal{G} .

In section 2.13 we define certain subcategories of nuclei and show how they are acted upon by the self-symmetries of section 2.9.

Note that in the classical algebraic construction the multiplication map of an algebra A makes it a left $N_l(A)$ -module, a right $N_r(A)$ -module, and a left- and right- $N_m(A)$ -module. Moreover, left multiplications by elements of $N_l(A)$ commute with right multiplications by elements of $N_m(A)$ and $N_r(A)$; while right multiplications by elements of $N_r(A)$ commute with left multiplications by elements of $N_m(A)$ and $N_l(A)$. Finally, the multiplication map $A \otimes A \longrightarrow A$ factors through $A \otimes_{N_m(A)} A$ and is a morphism of $N_l(A)$ - $N_r(A)$ -bimodules. In section 2.16 we categorify some of these constructions by defining actions of nuclei $\mathcal{N}(\mathcal{G})$ on the category \mathcal{G} .

This could give an impression that coherence laws of monoidal categories guarantee that categorifications of algebraic constructions are nicely coherent themselves. Unfortunately it is not always true, especially if commutativity enters the picture. For example, if A is a commutative (non-associative) algebra, then different nuclei are related as follows

$$N_l(A) = N_r(A) \subset N_m(A).$$

Indeed, the following chain of equalities proves that an element a of $N_l(A)$ belongs to $N_m(A)$:

$$x(ay) = (ay)x = a(yx) = a(xy) = (ax)y = (xa)y.$$

To see that a belongs to $N_r(A)$ one can use:

$$x(ya) = (ya)x = (ay)x = a(yx) = (yx)a = (xy)a.$$

On the level of categories we have similar relations between nuclei of a category \mathcal{G} with commutative tensor product:

$$\mathcal{N}_l(\mathcal{G}) \simeq \mathcal{N}_r(\mathcal{G}) \longrightarrow \mathcal{N}_m(\mathcal{G}).$$

But the functors between different nuclei are not monoidal.

In the second part of the paper we deal with a similar construction making maps between associative algebras multiplicative. Let $f : A \longrightarrow B$ be a linear map of associative algebras. Define $m_f(x, y) = f(xy) - f(x)f(y)$. The property

$$f(x)m_f(y, z) - m_f(xy, z) + m_f(x, yz) - m_f(x, y)f(z) = 0$$

implies that the subspaces

$$M_l(f) = \{a \in A : m_f(a, x) = 0 \ \forall x \in A\}, \quad M_r(f) = \{a \in A : m_f(x, a) = 0 \ \forall x \in A\}$$

of A are subalgebras (we call them the left and right *multiplicants* of the map f). It is straightforward to see that the restrictions of f to $M_l(f)$ and $M_r(f)$ are homomorphisms of algebras.

Again we face problems when we try to bring in commutativity. For example, the left and right multiplicants $M_l(f), M_r(f)$ of a linear map $f : A \longrightarrow B$ between commutative algebras coincide. The following chain of equalities shows that any $a \in M_l(f)$ belongs to $M_r(f)$:

$$f(xa) = f(ax) = f(a)f(x) = f(x)f(a).$$

But the equivalence $\mathcal{M}_l(F) \longrightarrow \mathcal{M}_r(F)$ between multiplicants of a functor $F : \mathcal{G} \longrightarrow \mathcal{H}$ between symmetric monoidal categories fails to be monoidal.

2. Nuclei of a category with a tensor product


2.1. NUCLEI AS SEMIGROUPAL CATEGORIES. Let \mathcal{G} be a category with a tensor product functor $\otimes : \mathcal{G} \times \mathcal{G} \longrightarrow \mathcal{G}$ without any associativity condition (*magmoidal* category). Following (2) define the *left nucleus* $\mathcal{N}_l(\mathcal{G})$ of \mathcal{G} as the category of pairs (A, a) , where $A \in \mathcal{G}$ and a denotes a natural family of isomorphisms

$$a_{X,Y} : A \otimes (X \otimes Y) \longrightarrow (A \otimes X) \otimes Y$$


indexed by all pairs of objects $X, Y \in \mathcal{G}$. A morphism $(A, a) \longrightarrow (B, b)$ in $\mathcal{N}_l(\mathcal{G})$ is a morphism $f : A \longrightarrow B$ in \mathcal{G} such that the diagram

$$\begin{array}{ccc} A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}} & (A \otimes X) \otimes Y \\ \downarrow f \otimes (X \otimes Y) & & \downarrow (f \otimes X) \otimes Y \\ B \otimes (X \otimes Y) & \xrightarrow{b_{X,Y}} & (B \otimes X) \otimes Y \end{array}$$

is commutative for any $X, Y \in \mathcal{G}$. Define the tensor product on the category $\mathcal{N}_l(\mathcal{G})$ by $(A, a) \otimes (B, b) = (A \otimes B, a|b)$, where $a|b$ is defined by the pentagon diagram:



The following diagram implies that for two morphisms $f : (A, a) \longrightarrow (B, b), g : (A', a') \longrightarrow (B', b')$ the tensor product $f \otimes g$ is a morphism between tensor products of pairs $(A, a) \otimes (A', a') \longrightarrow (B, b) \otimes (B', b')$.




In particular, the forgetful functor $\mathcal{N}_i(\mathcal{G}) \longrightarrow \mathcal{G}$ sending (A, a) to A is magmoidal.

It follows from the diagram below that for $(A, a), (B, b), (C, c) \in \mathcal{N}_i(\mathcal{G})$ the isomorphism $a_{B, C} : A \otimes (B \otimes C) \longrightarrow (A \otimes B) \otimes C$ belongs to the nucleus $\mathcal{N}_i(\mathcal{G})$. Indeed, it is equivalent to the commutativity of the square

$$\begin{array}{ccc}
 (A \otimes (B \otimes C)) \otimes (X \otimes Y) & \xrightarrow{(a|(b|c))_{X, Y}} & ((A \otimes (B \otimes C)) \otimes X) \otimes Y \\
 \downarrow a_{B, C \otimes (X \otimes Y)} & & \downarrow (a_{B, C \otimes X}) \otimes Y \\
 ((A \otimes B) \otimes C) \otimes (X \otimes Y) & \xrightarrow{((a|b)|c)_{X, Y}} & (((A \otimes B) \otimes C) \otimes X) \otimes Y
 \end{array}$$

which is one of the faces of the diagram $(K_5$ in Stasheff's notation [5]):



Two other square faces of this diagram are commutative by naturality while the pentagon faces commute by the definition of the tensor product in $\mathcal{N}_i(\mathcal{G})$. Thus

$$\phi_{(A,a),(B,b),(C,c)} : (A, a) \otimes ((B, b) \otimes (C, c)) \longrightarrow ((A, a) \otimes (B, b)) \otimes (C, c),$$

defined by $\phi_{(A,a),(B,b),(C,c)} = a_{B,C}$, is a morphism in the category $\mathcal{N}_i(\mathcal{G})$.

2.2. THEOREM. *The category $\mathcal{N}_i(\mathcal{G})$, with tensor product and associativity constraint defined above, is semigroupal.*

PROOF. All we need to verify is that the associativity constraint satisfies the pentagon axiom, which says that the diagram

$$\begin{array}{ccc}
 & (A \otimes B) \otimes (C \otimes D) & \\
 \nearrow^{a_{B,C \otimes D}} & & \searrow^{(a|b)_{C,D}} \\
 A \otimes (B \otimes (C \otimes D)) & & ((A \otimes B) \otimes C) \otimes D \\
 \searrow^{A \otimes b_{C,D}} & & \nearrow^{a_{B,C \otimes D}} \\
 A \otimes ((B \otimes C) \otimes D) & \xrightarrow{a_{B \otimes C,D}} & (A \otimes (B \otimes C)) \otimes D
 \end{array}$$

is commutative for any $(A, a), (B, b), (C, c), (D, d) \in \mathcal{N}_l(\mathcal{G})$. This obviously follows from the definition of $a|b$. ■

Semigroupal structures on the magmoidal category \mathcal{G} correspond to functors $\mathcal{G} \rightarrow \mathcal{N}_l(\mathcal{G})$ which split the forgetful functor $\mathcal{N}_l(\mathcal{G}) \rightarrow \mathcal{G}$. Note that this is close parallel with the corresponding result for monoidal centres (Proposition 4 from [2]).

2.3. THEOREM. *If \mathcal{G} is a semigroupal category with the associativity constraint ϕ , the assignment $A \mapsto (A, \phi_{A,-,-})$ defines a semigroupal functor $\mathcal{G} \rightarrow \mathcal{N}_l(\mathcal{G})$ whose composition $\mathcal{G} \rightarrow \mathcal{N}_l(\mathcal{G}) \rightarrow \mathcal{G}$ with the forgetful functor is the identity.*

Conversely, a functor $\mathcal{G} \rightarrow \mathcal{N}_l(\mathcal{G})$, whose composition $\mathcal{G} \rightarrow \mathcal{N}_l(\mathcal{G}) \rightarrow \mathcal{G}$ with the forgetful functor is the identity, defines an associativity constraint on \mathcal{G} thus turning it into a semigroupal category.

Moreover these constructions are mutually inverse.

PROOF. The pentagon axiom for ϕ implies that $\phi_{A,-,-} \circ \phi_{B,-,-} = \phi_{A \otimes B,-,-}$, so that $A \mapsto (A, \phi_{A,-,-})$ is a strict semigroupal functor.

A splitting $\mathcal{G} \rightarrow \mathcal{N}_l(\mathcal{G})$ of the forgetful functor $\mathcal{N}_l(\mathcal{G}) \rightarrow \mathcal{G}$ must have the form: $A \mapsto (A, a)$. It is straightforward to see that $\phi_{A,X,Y} = a_{X,Y}$ is an associativity constraint on \mathcal{G} . ■

Diagrams of this section (as well as their faces, edges and vertices) can be parameterized by certain partially parenthesised words of objects (morphisms) of \mathcal{G} . For example, the tensor product of two objects $X \otimes Y$ corresponds to the word XY , tensor products $A \otimes (X \otimes Y)$ and $(A \otimes X) \otimes Y$ correspond to the words $A(XY)$ and $(AX)Y$ respectively. The word AXY corresponds to the associativity morphism $a_{X,Y}$ and the word $ABXY$ to the pentagon diagram which was used to define the tensor product in $\mathcal{N}_l(\mathcal{G})$. Five edges of this pentagon correspond to the five ways of placing a pair of brackets $()$ in the word $ABXY$:

$$A(BXY), (AB)XY, A(BX)Y, AB(XY), (ABX)Y$$

which represent

$$A \otimes b_{X,Y}, (a|b)_{X,Y}, a_{B \otimes X,Y}, a_{B,X \otimes Y}, a_{B,X} \otimes Y$$

respectively. The diagram (5) for tensor product of morphisms corresponds to the word $fgXY$. Finally, Stasheff's polytope corresponds to the word $ABCXY$. Again its faces are parameterized by placements of a single pair of brackets () in the word $ABCXY$. Six pentagon faces of the Stasheff polytope correspond to the words:

$$A(BCXY), (AB)CXY, A(BC)XY, AB(CX)Y, ABC(XY), (ABCX)Y,$$

while three square faces are labeled by:

$$(ABC)XY, A(BCX)Y, AB(CXY).$$

The fact that the associativity morphism $a_{B,C}$ is a morphism in the nucleus $\mathcal{N}_l(\mathcal{G})$ follows from the commutativity of the face $(ABC)XY$. Finally, the pentagon axiom for its associativity is encoded by $ABCD$.

Using the word presentation for diagrams we can sketch the proof of the analog of theorem 2.2 for $\mathcal{N}_m(\mathcal{G})$. Define $\mathcal{N}_m(\mathcal{G})$ to be the category of pairs (A, a) , where $A \in \mathcal{G}$ and a is a natural collection of isomorphisms

$$a_{X,Y} : X \otimes (A \otimes Y) \longrightarrow (X \otimes A) \otimes Y$$

which we label by the word XAY . As before morphisms in $\mathcal{N}_m(\mathcal{G})$ are morphisms in \mathcal{G} compatible with natural isomorphisms. We can define the tensor product of pairs $(A, a) \otimes (B, a) = (A \otimes B, a|b)$ using the pentagon with label $XABY$. The diagram of shape $XfgY$ implies that the tensor product of morphisms $f \otimes g$ is a morphism of tensor products. It follows from commutativity properties of the shape $XABCY$ that the formula $\phi_{(A,a),(B,b),(C,c)} = b_{A,C}$ defines a natural collection of isomorphisms in $\mathcal{N}_m(\mathcal{G})$:

$$\phi_{(A,a),(B,b),(C,c)} : (A, a) \otimes ((B, b) \otimes (C, c)) \longrightarrow ((A, a) \otimes (B, b)) \otimes (C, c).$$

The word $ABCD$ gives the pentagon axiom for this constraint.

Similarly, define $\mathcal{N}_r(\mathcal{G})$ to be the category of pairs (A, a) , with

$$a_{X,Y} : X \otimes (Y \otimes A) \longrightarrow (X \otimes Y) \otimes A$$

which we label by the word XYA . Again, morphisms in $\mathcal{N}_r(\mathcal{G})$ are morphisms in \mathcal{G} compatible with the natural isomorphisms. The tensor product of pairs $(A, a) \otimes (B, a) = (A \otimes B, a|b)$ is defined by the pentagon with the label $XYAB$. The diagram of shape $XYfg$ implies that the tensor product of morphisms $f \otimes g$ is a morphism of tensor products. The shape $XYABC$ implies that the formula $\phi_{(A,a),(B,b),(C,c)} = c_{A,B}$ defines a natural collection of isomorphisms in $\mathcal{N}_r(\mathcal{G})$:

$$\phi_{(A,a),(B,b),(C,c)} : (A, a) \otimes ((B, b) \otimes (C, c)) \longrightarrow ((A, a) \otimes (B, b)) \otimes (C, c).$$

Finally, the word $ABCD$ gives the pentagon axiom for this constraint.

2.5. UNITS. In this section we slightly modify the construction to incorporate units. First we need to modify the setup.

Now let \mathcal{G} be a magmoidal category with *unit* (an object $1 \in \mathcal{G}$ with natural isomorphisms $l_X : 1 \otimes X \rightarrow X$, $r_X : X \otimes 1 \rightarrow X$ such that $l_1 = r_1$). Define the left *unital nucleus* $\mathcal{N}_l^1(\mathcal{G})$ as the category of pairs (A, a) where as before $a_{X,Y} : A \otimes (X \otimes Y) \rightarrow (A \otimes X) \otimes Y$ is natural collection of isomorphism but now satisfying *normalisation* conditions: the diagrams

$$\begin{array}{ccc}
 A \otimes (1 \otimes X) & \xrightarrow{a_{1,X}} & (A \otimes 1) \otimes X \\
 \searrow A \otimes l_X & & \swarrow r_{A \otimes X} \\
 & & A \otimes X
 \end{array}
 \qquad
 \begin{array}{ccc}
 A \otimes (X \otimes 1) & \xrightarrow{a_{X,1}} & (A \otimes X) \otimes 1 \\
 \searrow A \otimes r_X & & \swarrow A \otimes X \\
 & & A \otimes X
 \end{array}$$

commute for any $X \in \mathcal{G}$.

Define a natural collection of isomorphisms $i_{X,Y} : 1 \otimes (X \otimes Y) \rightarrow (1 \otimes X) \otimes Y$ as the composition $1 \otimes (X \otimes Y) \xrightarrow{l_{X \otimes Y}} X \otimes Y \xrightarrow{(l_{X \otimes Y})^{-1}} (1 \otimes X) \otimes Y$.

2.6. LEMMA. *The pair $(1, i)$ is an object of $\mathcal{N}_l^1(\mathcal{G})$.*

PROOF. The normalisation conditions for the collection i follow from the commutative diagrams:

$$\begin{array}{ccc}
 1 \otimes (1 \otimes Y) & \xrightarrow{i_{1,Y}} & (1 \otimes 1) \otimes Y \\
 \searrow l_{1 \otimes Y} & & \swarrow l_{1 \otimes Y} \\
 & & 1 \otimes Y \\
 \swarrow 1 \otimes l_Y & & \searrow r_{1 \otimes Y}
 \end{array}$$

$$\begin{array}{ccc}
 1 \otimes (X \otimes 1) & \xrightarrow{i_{1,1}} & (1 \otimes X) \otimes 1 \\
 \searrow 1 \otimes r_X & & \swarrow r_{1 \otimes X} \\
 & & 1 \otimes X \\
 \swarrow l_{X \otimes 1} & & \searrow l_{X \otimes 1} \\
 & & X \\
 & & \uparrow r_X \\
 & & X \otimes 1
 \end{array}$$

Here the equation $l_{1 \otimes Y} = 1 \otimes l_Y$ (used in the first diagram) follows from the naturality of l :

$$\begin{array}{ccc}
 1 \otimes (1 \otimes Y) & \xrightarrow{l_{1 \otimes Y}} & 1 \otimes Y \\
 1 \otimes l_Y \downarrow & & \downarrow l_Y \\
 1 \otimes Y & \xrightarrow{l_Y} & Y
 \end{array}$$



In the second diagram we use the naturality of l with respect to r_X and the naturality of r with respect to l_X . ■

2.7. THEOREM. *The category $\mathcal{N}_l^1(\mathcal{G})$ is monoidal with unit object $(1, i)$ and unit natural isomorphisms*

$$l_{(A,a)} : (1, i) \otimes (A, a) \longrightarrow (A, a), \quad r_{(A,a)} : (A, a) \otimes (1, i) \longrightarrow (A, a) \quad (6)$$

defined by $l_{(A,a)} = l_A$, $r_{(A,a)} = r_A$.

PROOF. First we need to check that the tensor product of pairs as it was defined for $\mathcal{N}_l(\mathcal{G})$ preserves the normalisation conditions. This follow from the commutative diagrams:





Thus the subcategory $\mathcal{N}_l^1(\mathcal{G})$ is closed under tensor product in $\mathcal{N}_l(\mathcal{G})$ and is a semigroupal category. By lemma 2.6, the pair $(1, i)$ is an object of $\mathcal{N}_l^1(\mathcal{G})$. We need to check that the unit isomorphisms (6) are morphisms in the category $\mathcal{N}_l^1(\mathcal{G})$. It is equivalent to the

commutativity of the squares:

$$\begin{array}{ccc}
 (1A)(XY) \xrightarrow{(i|a)_{X,Y}} ((1A)X)Y & & (A1)(XY) \xrightarrow{(a|i)_{X,Y}} ((A1)X)Y \\
 \downarrow l_{A,XY} & & \downarrow r_{A,XY} \\
 A(XY) \xrightarrow{a_{X,Y}} (AX)Y & & A(XY) \xrightarrow{a_{X,Y}} (AX)Y \\
 \downarrow (l_{AX})Y & & \downarrow (r_{AX})Y
 \end{array}$$

which follows from the commutative diagrams:



Finally, the coherence for the unit isomorphisms

$$\begin{array}{ccc}
 (A, a) \otimes ((1, i) \otimes (B, b)) & \xrightarrow{\phi_{(A,a),(1,i),(B,b)}} & ((A, a) \otimes (1, i)) \otimes (B, b) \\
 \searrow (A,a) \otimes l_{(B,a)} & & \swarrow r_{(A,a) \otimes (B,b)} \\
 & (A, a) \otimes (B, b) &
 \end{array}$$

boils down to

$$\begin{array}{ccc}
 A \otimes (1 \otimes B) & \xrightarrow{a_{1,B}} & (A \otimes 1) \otimes B \\
 & \searrow l_b & \swarrow r_A \\
 & A \otimes B &
 \end{array}$$

which is commutative by the definition of $\mathcal{N}_l^1(\mathcal{G})$. ■

The constructions of the following sections can be made unital (monoidal instead of just semi-groupal). For the sake of brevity we will not be doing it.

2.9. FUNCTORIALITY. Here we assign, to an invertible natural self-transformation (an automorphism) c of the tensor product functor $\otimes : \mathcal{G} \times \mathcal{G} \longrightarrow \mathcal{G}$, a strict semigroupal autoequivalence $\mathcal{N}_l(c) : \mathcal{N}_l(\mathcal{G}) \longrightarrow \mathcal{N}_l(\mathcal{G})$ of the nucleus. For an object (A, a) of $\mathcal{N}_l(\mathcal{G})$ set $\mathcal{N}_l(c)(A, a)$ to be (A, a^c) where a^c is defined by the diagram:

$$\begin{array}{ccccc}
 A \otimes (X \otimes Y) & \xrightarrow{c_{A, X \otimes Y}} & A \otimes (X \otimes Y) & \xrightarrow{A \otimes c_{X, Y}} & A \otimes (X \otimes Y) \\
 \downarrow \alpha_{X, Y} & & & & \downarrow \alpha_{X, Y}^c \\
 (A \otimes X) \otimes Y & \xrightarrow{c_{A \otimes X, Y}} & (A \otimes X) \otimes Y & \xrightarrow{c_{A, X \otimes Y}} & (A \otimes X) \otimes Y
 \end{array}$$

2.10. PROPOSITION. *The functor $\mathcal{N}_l(c)$ is strict semigroupal.*

PROOF. The following commutative diagram shows that $(a|b)^c = (a^c|b^c)$ for $(A, a), (B, b)$ from $\mathcal{N}_l(\mathcal{G})$:

Now we establish the link between natural transformations of the identity functor on \mathcal{G} and monoidal natural transformations of functors of the form $\mathcal{N}_l(c)$. First we define an action of the group $\text{Aut}(\text{id}_{\mathcal{G}})$ of automorphisms of the identity functor on the group $\text{Aut}(\otimes)$ of automorphisms of the tensor product functor. For $c \in \text{Aut}(\otimes)$ and $f \in \text{Aut}(\text{id}_{\mathcal{G}})$, define $c^f \in \text{Aut}(\otimes)$ by

$$c_{X,Y}^f = (f_X \otimes f_Y)c_{X,Y}f_{X \otimes Y}^{-1}.$$

Note that by naturality of c , $c_{X,Y}$ commutes with $f_X \otimes f_Y$ and by naturality of f , $c_{X,Y}$ commutes with $f_{X \otimes Y}$ so $c_{X,Y}^f$ coincides with

$$(f_X \otimes f_Y)f_{X \otimes Y}^{-1}c_{X,Y} = c_{X,Y}(f_X \otimes f_Y)f_{X \otimes Y}^{-1}.$$

This in particular implies that $\text{Aut}(\text{id}_{\mathcal{G}})$ acts on the group $\text{Aut}(\otimes)$:

$$c_{X,Y}^{fg} = c_{X,Y}(fg_X \otimes fg_Y)fg_{X \otimes Y}^{-1} = c_{X,Y}(f_X \otimes f_Y)f_{X \otimes Y}^{-1}(g_X \otimes g_Y)g_{X \otimes Y}^{-1} = (c^f)_{X,Y}^g.$$

2.12. PROPOSITION. *A natural transformation $f \in \text{Aut}(\text{id}_{\mathcal{G}})$ of the identity functor defines a natural transformation $\mathcal{N}_l(c) \longrightarrow \mathcal{N}_l(c^f)$.*

PROOF. We need to show that for any $(A, a) \in \mathcal{N}_l(\mathcal{G})$ the map $f_A : A \longrightarrow A$ is a morphism $(A, a) \longrightarrow (A, a^{c^f})$ in $\mathcal{N}_l(\mathcal{G})$ or that the square

$$\begin{array}{ccc} A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}^c} & (A \otimes X) \otimes Y \\ f_{A \otimes (X \otimes Y)} \downarrow & & \downarrow (f_A \otimes X) \otimes Y \\ A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}^{c^f}} & (A \otimes X) \otimes Y \end{array}$$

is commutative. The commutativity of this square is equivalent to the commutativity of the bottom square of the following diagram:

$$\begin{array}{ccccc} & & A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}} & (A \otimes X) \otimes Y \\ & f_{A \otimes (X \otimes Y)} \nearrow & \downarrow & f_{A \otimes X} \otimes Y \nearrow & \downarrow \\ A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}} & (A \otimes X) \otimes Y & & \downarrow c_{A \otimes X, Y}^f \\ \downarrow c_{A, X \otimes Y} & & \downarrow c_{A, X \otimes Y}^f & & \downarrow c_{A \otimes X, Y}^f \\ & f_{A \otimes f_{X \otimes Y}} \nearrow & A \otimes (X \otimes Y) & & (A \otimes X) \otimes Y \\ A \otimes (X \otimes Y) & & \downarrow & f_{A, X} \otimes f_Y \nearrow & \downarrow c_{A, X \otimes Y}^f \\ \downarrow A \otimes c_{X, Y} & & \downarrow A \otimes c_{X, Y}^f & & \downarrow c_{A, X \otimes Y}^f \\ & f_{A \otimes (f_X \otimes f_Y)} \nearrow & A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}^{c^f}} & (A \otimes X) \otimes Y \\ & & \downarrow c_{A, X \otimes Y} & & \downarrow c_{A, X \otimes Y}^f \\ A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}^c} & (A \otimes X) \otimes Y & & \downarrow c_{A, X \otimes Y}^f \\ & & \downarrow & f_{(A \otimes f_X) \otimes f_Y} \nearrow & \\ & & A \otimes (X \otimes Y) & & \downarrow c_{A, X \otimes Y}^f \\ & & \downarrow & & \\ & & A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}^c} & (A \otimes X) \otimes Y \\ & & \downarrow & & \downarrow \\ & & A \otimes (X \otimes Y) & & (A \otimes X) \otimes Y \end{array}$$

■

It is clear that the natural transformation $\mathcal{N}_l(f) : \mathcal{N}_l(c) \longrightarrow \mathcal{N}_l(c^f)$ is monoidal iff $f \in \text{Aut}(\text{id}_{\mathcal{G}})$ is monoidal, i.e. iff $f_{X \otimes Y} = f_X \otimes f_Y$ for any $X, Y \in \mathcal{G}$.

2.13. SUBCATEGORIES OF NUCLEI. Here we define semigroupal (monoidal) subcategories in nuclei anchored at quasi-monoidal functors. Let \mathcal{G} be a magmoidal category, let \mathcal{H} be a monoidal (semi-groupal) category and let $F : \mathcal{G} \longrightarrow \mathcal{H}$ be a functor equipped with a natural collection of isomorphisms $F_{X,Y} : F(X \otimes Y) \longrightarrow F(X) \otimes F(Y)$ (a *quasi-monoidal* functor). Define a full subcategory $\mathcal{N}_l(F)$ of $\mathcal{N}_l(\mathcal{G})$ consisting of pairs (A, a) such that the following diagram commutes:

$$\begin{array}{ccc}
 F(A \otimes (X \otimes Y)) & \xrightarrow{F_{X,Y}} & F(A) \otimes F(X \otimes Y) & \xrightarrow{F(A) \otimes F_{X,Y}} & F(A) \otimes (F(X) \otimes F(Y)) & (7) \\
 \downarrow F(a_{X,Y}) & & & & \downarrow \psi_{F(A), F(X), F(Y)} \\
 F((A \otimes X) \otimes Y) & \xrightarrow{F_{A \otimes X, Y}} & F(A \otimes X) \otimes F(Y) & \xrightarrow{F_{X,A} \otimes F(Y)} & (F(A) \otimes F(X)) \otimes F(Y)
 \end{array}$$

here ψ is the associativity constraint of \mathcal{H} .

2.14. PROPOSITION. *The subcategory $\mathcal{N}_l(F)$ is closed under tensor product in $\mathcal{N}_l(\mathcal{G})$. The quasi-monoidal functor F lifts to a monoidal (semi-groupal) functor $\overline{F} : \mathcal{N}_l(F) \longrightarrow \mathcal{H}$.*


PROOF. We need to verify that the tensor product $(A \otimes B, a|b)$ of two pairs $(A, a), (B, b)$ from $\mathcal{N}_l(F)$ is in $\mathcal{N}_l(F)$. It follows from the commutativity of the diagram:

Define a new quasi-monoidal structure F^c on F by

$$F_{X,Y}^c : F(X \otimes Y) \xrightarrow{F(c_{X,Y})} F(X \otimes Y) \xrightarrow{F_{X,Y}} F(X) \otimes F(Y)$$

2.15. PROPOSITION. *The semigroupal functor $\mathcal{N}_i(c)$ defines an equivalence between the semi-groupal subcategories $\mathcal{N}_i(F)$ and $\mathcal{N}_i(F^c)$.*

PROOF. We need to show that the coherence diagram (7) commutes if we replace $F_{X,Y}$ by $F_{X,Y}^c$ and $a_{X,Y}$ by $a_{X,Y}^c$. This follows from the commutative diagram:



■

2.16. ACTIONS OF NUCLEI. Recall that a monoidal (semigroupal) category \mathcal{C} acts on a category \mathcal{M} if there is given a monoidal (semigroupal) functor $\mathcal{C} \rightarrow \text{End}(\mathcal{M})$ into the category of endofunctors on \mathcal{M} . In that case \mathcal{M} is called a (left) *module category* over \mathcal{C} or a (left) \mathcal{C} -category.

Let $L : \mathcal{C} \rightarrow \text{End}(\mathcal{M})$, $L' : \mathcal{C} \rightarrow \text{End}(\mathcal{M}')$ be two actions. A functor $F : \mathcal{M} \rightarrow \mathcal{M}'$ between \mathcal{C} -categories is a \mathcal{C} -functor if it is equipped with a collection of isomorphisms $F_{A,X} : F(L(A)(X)) \rightarrow L'(A)(F(X))$ natural in $A \in \mathcal{C}$, $X \in \mathcal{M}$ such that the diagram

$$\begin{array}{ccccc}
 F(L(AB)(X)) & \xrightarrow{F_{AB,X}} & L'(AB)(F(X)) & \xrightarrow{L'_{A,B}(F(X))} & L'(A)(L'(B)(F(X))) & (8) \\
 \downarrow F_{L(A),B(X)} & & & & \parallel \\
 F(L(A)(L(B)(X))) & \xrightarrow{F_{A,L(B)(X)}} & L'(A)(F(L(B)(X))) & \xrightarrow{L'(A)(F_{B,X})} & L'(A)(L'(B)(F(X)))
 \end{array}$$

commutes.

For a magmoidal category \mathcal{G} define a functor $L : \mathcal{N}_i(\mathcal{G}) \rightarrow \text{End}(\mathcal{G})$ by $L(A, a)(X) = A \otimes X$. Define a monoidal structure $L_{(A,a),(B,b)} : L((A, a) \otimes (B, b)) \rightarrow L(A, a) \circ L(B, b)$

PROOF. We need to check that the natural collection (9) satisfies the coherence condition (8). It follows from the diagram:

$$\begin{array}{ccccc}
 L((A, a)(B, b))(XY) & \xrightarrow{L_{(A,a),(B,b)}(X)Y} & L(A, a)(L(B, b)(X))Y & & \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & ((AB)X)Y & \xrightarrow{a_{B,XY}} & (A(BX))Y & \\
 \downarrow \otimes_{(A,a)(B,b),X,Y} & \uparrow & \downarrow \otimes_{(A,a),L(B,b)(X),Y} & \uparrow & \\
 L((A, a)(B, b))(XY) & & L(A, a)(L(B, b)(X)Y) & & \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & (AB)(XY) & & A((BX)Y) & \\
 \downarrow L_{(A,a),(B,b)}(XY) & \uparrow (a|b)_{X,Y} & \downarrow L_{(A,a)}(\otimes_{(B,b),X,Y}) & \uparrow a_{BX,Y} & \\
 L(A, a)(L(B, b)(XY)) & = & (L(A, a)L(B, b))(XY) & & \\
 \downarrow & \uparrow a_{B,XY} & \downarrow & \uparrow Ab_{X,Y} & \\
 & A(B(XY)) & = & A(B(XY)) &
 \end{array}$$

■

In the next proposition we characterise left nucleus $\mathcal{N}_l(\mathcal{G})$ as a category of certain endofunctors on \mathcal{G} . For a magmoidal category \mathcal{G} a functor $F : \mathcal{G} \rightarrow \mathcal{G}$ will be called (right) \mathcal{G} -invariant if it is equipped with a natural collection of isomorphisms $f_{X,Y} : F(X \otimes Y) \rightarrow F(X) \otimes Y$. A morphism $c : F \rightarrow G$ of \mathcal{G} -invariant functors is a natural transformation such that the diagram

$$\begin{array}{ccc}
 F(X \otimes Y) & \xrightarrow{f_{X,Y}} & F(X) \otimes Y \\
 \downarrow c_{X \otimes Y} & & \downarrow c_{X \otimes Y} \\
 G(X \otimes Y) & \xrightarrow{g_{X,Y}} & G(X) \otimes Y
 \end{array}$$

commutes. Define $\text{End}_{\mathcal{G}}(\mathcal{G})$ to be the category of \mathcal{G} -invariant functors. The composite of two \mathcal{G} -invariant functors $(F, f), (G, g)$ has a structure of \mathcal{G} -invariant functor:

$$\begin{array}{ccc}
 (F \circ G)(X \otimes Y) & \xrightarrow{(f \circ g)_{X,Y}} & (F \circ G)(X) \otimes Y \\
 \parallel & & \parallel \\
 F(G(X \otimes Y)) & \xrightarrow{F(g_{X,Y})} F(G(X) \otimes Y) \xrightarrow{f_{G(X),Y}} & F(G(X)) \otimes Y
 \end{array}$$

2.19. LEMMA. *The category $\text{End}_{\mathcal{G}}(\mathcal{G})$ is strict monoidal.*

PROOF. All we need to verify is that the horizontal composition of morphisms of \mathcal{G} -invariant functors preserves the \mathcal{G} -invariant structure of the composition. This follows from the commutative diagram:

$$\begin{array}{ccccc}
 & F(G(XY)) & \xrightarrow{F(g_{X,Y})} & F(G(X)Y) & \xrightarrow{f_{G(X),Y}} & F(G(X))Y \\
 & \parallel & & \swarrow & & \parallel \\
 (F \circ G)(XY) & \xrightarrow{F(d_{XY})} & (F \circ G)(X)Y & \xrightarrow{f_{G(X),Y}} & (F \circ G)(X)Y & \xrightarrow{c_{G(X),Y}} & (F \circ G)(X)Y \\
 & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
 & F(G'(XY)) & \xrightarrow{F(g'_{X,Y})} & F(G'(X)Y) & \xrightarrow{f'_{G'(X),Y}} & F'(G'(X))Y \\
 & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
 & F'(G'(XY)) & \xrightarrow{F'(g'_{X,Y})} & F'(G'(X)Y) & \xrightarrow{f'_{G'(X),Y}} & F'(G'(X))Y \\
 & \parallel & & \swarrow & & \parallel \\
 (F' \circ G')(XY) & \xrightarrow{(f' \circ g')_{X,Y}} & (F' \circ G')(X)Y & \xrightarrow{f'_{G'(X),Y}} & (F' \circ G')(X)Y & \xrightarrow{f'_{G'(X),Y}} & (F' \circ G')(X)Y
 \end{array}$$


■

Now we extend the (left) action $L : \mathcal{N}_l(\mathcal{G}) \rightarrow \text{End}(\mathcal{G})$ to a functor $L : \mathcal{N}_l(\mathcal{G}) \rightarrow \text{End}_{\mathcal{G}}(\mathcal{G})$ by the assignment $(A, a) \mapsto (L(A), L(a))$, where the structure $L(a)$ of \mathcal{G} -invariant functor on $L(A)$ is defined as follows:

$$\begin{array}{ccc}
 L(A)(X \otimes Y) & \xrightarrow{L(a)_{X,Y}} & L(A)(X) \otimes Y \\
 \parallel & & \parallel \\
 A \otimes (X \otimes Y) & \xrightarrow{a_{X,Y}} & (A \otimes X) \otimes Y
 \end{array}$$

2.20. PROPOSITION. *Let \mathcal{G} be a unital magmoidal category. Then the functor $L : \mathcal{N}_l^1(\mathcal{G}) \rightarrow \text{End}_{\mathcal{G}}(\mathcal{G})$ is a monoidal equivalence.*

PROOF. First we note that the monoidal structure $L_{X,Y}$ of the functor $L : \mathcal{N}_l^1(\mathcal{G}) \rightarrow \text{End}(\mathcal{G})$ is compatible with the \mathcal{G} -invariant structures (a morphism in $\text{End}_{\mathcal{G}}(\mathcal{G})$) thus making the extended functor $L : \mathcal{N}_l^1(\mathcal{G}) \rightarrow \text{End}_{\mathcal{G}}(\mathcal{G})$ monoidal. This follows from the commutative diagram:



To see that L is an equivalence it is enough to note that the assignment $(F, f) \mapsto (F(1), \bar{f})$ defines a functor quasi-inverse to L . Here

$$\bar{f}_{X,Y} : F(1) \otimes (X \otimes Y) \longrightarrow (F(1) \otimes X) \otimes Y$$

is defined by

$$\begin{array}{ccccc} F(1) \otimes (X \otimes Y) & \xleftarrow{F_{1,X \otimes Y}} & F(1 \otimes (X \otimes Y)) & \xrightarrow{F(l_{X \otimes Y})} & F(X \otimes Y) \\ \downarrow \bar{f}_{X,Y} & & & & \downarrow F_{X,Y} \\ (F(1) \otimes X) \otimes Y & \xleftarrow{F_{1,X \otimes Y}} & F(1 \otimes X) \otimes Y & \xrightarrow{F(l_X) \otimes Y} & F(X) \otimes Y \end{array}$$

■

3. Multiplicants of functors

Now let $F : \mathcal{G} \longrightarrow \mathcal{H}$ be a functor between semigroupal (monoidal) categories. Define $\mathcal{M}_l(F)$ to be the category of pairs (A, a) , where $A \in \mathcal{G}$ and a is a natural family of isomorphisms $a_X : F(A \otimes X) \longrightarrow F(A) \otimes F(X)$. A morphism from (A, a) to (B, b) is a morphism $f : A \longrightarrow B$ in \mathcal{G} such that the following diagram commutes for each $X \in \mathcal{G}$

$$\begin{array}{ccc} F(A \otimes X) & \xrightarrow{a_X} & F(A) \otimes F(X) \\ F(f \otimes X) \downarrow & & \downarrow F(f) \otimes F(X) \\ F(B \otimes X) & \xrightarrow{b_X} & F(A) \otimes F(X) \end{array}$$

Note that the assignment $(A, a) \mapsto A$ defines a functor $\mathcal{M}_l(F) \longrightarrow \mathcal{G}$ (the *forgetful* functor).

We can define a tensor product on $\mathcal{M}_l(F)$ by $(A, a) \otimes (B, b) = (A \otimes B, a|b)$, where the natural family $a|b$ is given by the diagram

$$\begin{array}{ccccc} F(A \otimes (B \otimes X)) & \xrightarrow{a_{B \otimes X}} & F(A) \otimes F(B \otimes X) & \xrightarrow{F(A) \otimes b_X} & F(A) \otimes (F(B) \otimes F(X)) \\ F(\phi_{A,B,X}) \downarrow & & & & \psi_{F(A),F(B),F(X)} \downarrow \\ F((A \otimes B) \otimes X) & \xrightarrow{(a|b)_X} & F(A \otimes B) \otimes F(X) & \xrightarrow{a_B \otimes F(X)} & (F(A) \otimes F(B)) \otimes F(X) \end{array}$$


3.1. THEOREM. *The category $\mathcal{M}_l(F)$ is semigroupal. The forgetful functor $\mathcal{M}_l(F) \rightarrow \mathcal{G}$ is a strict semigroupal functor. The composition of the forgetful functor with F is a semigroupal functor $\bar{F} : \mathcal{M}_l(F) \rightarrow \mathcal{H}$ with semigroupal structure:*

$$\bar{F}_{(A,a),(B,b)} = a_B : F(A \otimes B) \rightarrow F(A) \otimes F(B).$$

PROOF. The following diagram proves that the tensor product $f \otimes g$ of morphisms $f : (A, a) \rightarrow (C, c)$, $g : (B, b) \rightarrow (D, d)$ is a morphism from $(A, a) \otimes (B, b)$ to $(C, c) \otimes (D, d)$:

$$\begin{array}{ccccc} & & F(C \otimes (D \otimes X)) & \xrightarrow{F(\phi_{C,D,X})} & F((C \otimes D) \otimes X) \\ & \nearrow F(f \otimes (g \otimes I)) & \downarrow F(\phi_{A,B,X}) & & \nearrow F((f \otimes g) \otimes I) \\ F(A \otimes (B \otimes X)) & \xrightarrow{F(\phi_{A,B,X})} & F((A \otimes B) \otimes X) & & \downarrow (c|d)_X \\ \downarrow a_{B \otimes X} & & \downarrow c_{D \otimes X} & & \downarrow \\ F(A) \otimes F(B \otimes X) & \nearrow F(f) \otimes F(g \otimes I) & F(C) \otimes F(D \otimes X) & \xrightarrow{(a|b)_X} & F(A \otimes B) \otimes F(X) \\ & \downarrow I \otimes d_X & \downarrow I \otimes d_X & & \nearrow F(f \otimes g) \otimes I \\ & & F(C) \otimes F(D) \otimes F(X) & & \downarrow c_{D \otimes I} \\ \downarrow I \otimes b_X & & \downarrow \phi_{F(C),F(D),F(X)} & & \downarrow \\ F(A) \otimes (F(B) \otimes F(X)) & \nearrow F(f) \otimes (F(g) \otimes I) & F(C) \otimes (F(D) \otimes F(X)) & \xrightarrow{\phi_{F(C),F(D),F(X)}} & (F(C) \otimes F(D)) \otimes F(X) \\ & \downarrow F(f) \otimes (F(g) \otimes I) & \downarrow \phi_{F(A),F(B),F(X)} & & \nearrow (F(f) \otimes F(g)) \otimes I \\ & & F(A) \otimes (F(B) \otimes F(X)) & \xrightarrow{\phi_{F(A),F(B),F(X)}} & (F(A) \otimes F(B)) \otimes F(X) \end{array}$$

It follows from the diagram



that the square

$$\begin{array}{ccc}
 F((A(BC))X) & \xrightarrow{(a|(b|c))_X} & F(A(BC))F(X) \\
 F(\phi X) \downarrow & & \downarrow F(\phi)F(X) \\
 F(((AB)C)X) & \xrightarrow{((a|b)|c)_X} & F((AB)C)F(X)
 \end{array}$$

commutes, which means that the associativity constraint

$\phi_{A,B,C} : A \otimes (B \otimes C) \longrightarrow (A \otimes B) \otimes C$ of \mathcal{G} is a morphisms of $\mathcal{M}_l(F)$:

$$\phi_{(A,a),(B,b),(C,c)} = \phi_{A,B,C}.$$

Thus $\mathcal{M}_l(F)$ is semigroupal and the forgetful functor $\mathcal{M}_l(F) \longrightarrow \mathcal{G}$ is strict semigroupal. It follows from the definition of $\mathcal{M}_l(F)$ that \overline{F} is semigroupal. ■

The semigroupal category $\mathcal{M}_l(F)$ becomes monoidal if we assume that the functor F preserves monoidal unit, i.e. there is given an isomorphism $\epsilon : F(1) \longrightarrow 1$. It can be verified that the object $(1, \iota) \in \mathcal{M}_l(F)$ is a monoidal unit. Here ι_X is the composition:

$$F(1 \otimes X) \xrightarrow{F(\iota_X)} F(X) \xleftarrow{\iota_{F(X)}} 1 \otimes F(X) \xleftarrow{\epsilon \otimes I} F(1) \otimes F(X)$$

The semigroupal functors $\mathcal{M}_l(F) \longrightarrow \mathcal{G}, \mathcal{H}$ allow us to define $\mathcal{M}_l(F)$ -actions on the categories \mathcal{G} and \mathcal{H} . In the next proposition we show that the functor F preserves these actions.

3.3. PROPOSITION. *The functor $F : \mathcal{G} \longrightarrow \mathcal{H}$ has a natural structure of a $\mathcal{M}_l(F)$ -functor.*

PROOF. The $\mathcal{M}_l(F)$ -action on \mathcal{G} is given by $(X, x)(Y) = X \otimes Y$ for $Y \in \mathcal{G}$, while the $\mathcal{M}_l(F)$ -action on \mathcal{H} is simply $(X, x)(Z) = F(X) \otimes Z$ for $Z \in \mathcal{H}$. The composition

$$F((X, x)(Y)) = F(X \otimes Y) \xrightarrow{xy} F(X)F(Y) = (X, x)F(Y)$$

defines the structure of a $\mathcal{M}_l(F)$ -functor on F . The coherence for this structure follows from the definition of tensor product in $\mathcal{M}_l(F)$. ■


3.4. FUNCTORIALITY. Here we define a correspondence between isomorphisms $c : F \longrightarrow G$ of functors $F, G : \mathcal{G} \longrightarrow \mathcal{H}$ between monoidal categories and monoidal functors $\mathcal{M}_l(c) : \mathcal{M}_l(F) \longrightarrow \mathcal{M}_l(G)$ between their multiplicants. For an object (A, a) of $\mathcal{M}_l(F)$, define $\mathcal{M}_l(c)(A, a)$ to be (A, a^c) where a^c is defined by the diagram:

$$\begin{array}{ccc} F(A \otimes X) & \xrightarrow{a_X} & F(A) \otimes F(X) \\ c_{A \otimes X} \downarrow & & \downarrow c_A \otimes c_X \\ G(A \otimes X) & \xrightarrow{a_X^c} & G(A) \otimes G(X) \end{array}$$

3.5. PROPOSITION. *The functor $\mathcal{M}_l(c) : \mathcal{M}_l(F) \longrightarrow \mathcal{M}_l(G)$ defined above is strict monoidal and fits into a commutative diagram of monoidal functors:*

$$\begin{array}{ccc} \mathcal{M}_l(F) & \xrightarrow{\mathcal{M}_l(c)} & \mathcal{M}_l(G) \\ \downarrow & \searrow & \downarrow \\ \mathcal{G} & & \mathcal{H} \end{array} \tag{10}$$

PROOF. The following commutative diagram shows that $(a|b)^c = (a^c|b^c)$ for $(A, a), (B, b) \in \mathcal{M}_l(F)$:



This proves that the functor $\mathcal{M}_l(c)$ is strict monoidal:

$$\mathcal{M}_l(c)((A, a) \otimes (B, b)) = \mathcal{M}_l(c)(A \otimes B, a|b) = (A \otimes B, (a|b)^c)$$

coincides with

$$\mathcal{M}_l(c)(A, a) \otimes \mathcal{M}_l(c)(B, b) = (A, a^c) \otimes (B, b^c) = (A \otimes B, a^c|b^c).$$


Commutativity of the diagram (10) follows from the definition of $\mathcal{M}_l(c)$. ■

Note that for successive isomorphisms of functors $F \xrightarrow{c} G \xrightarrow{d} H$ the composition of monoidal functors $\mathcal{M}_l(c)$ and $\mathcal{M}_l(d)$ coincides with $\mathcal{M}_l(cd)$. This follows from the fact that $a^{cd} = (a^c)^d$ for $(A, a) \in \mathcal{M}_l(F)$.

3.6. COMPOSITION PROPERTIES. For a successive (set-theoretic) maps of monoids

$$A \xrightarrow{f} B \xrightarrow{g} C$$

(left) multiplicants form a successive pair of spans (in the category of monoids):




The composition of spans is given by the pullback (fibered product) of the two middle maps:

$$M_l(f) \times_B M_l(g) = \{(a, f(a)) \mid a \in M_l(f), f(a) \in M_l(g)\}.$$

Note that the “projection” $(a, f(a)) \mapsto a$ defines a homomorphism $M_l(f) \times_B M_l(g) \longrightarrow M_l(gf)$. Indeed, for such a and any $x \in A$ we have that

$$gf(ax) = g(f(a)f(x)) = gf(a)gf(x).$$

Moreover, this homomorphism fits into a commutative diagram (morphism of spans):



Thus multiplicands define an oplax functor from the category of monoids and set-theoretic maps into the bicategory of spans of monoids.

Here we categorify this construction by defining a monoidal functor $\mathcal{M}_l(F) \times_{\mathcal{H}} \mathcal{M}_l(G) \longrightarrow \mathcal{M}_l(GF)$ for a pair of composable functors $\mathcal{G} \xrightarrow{F} \mathcal{H} \xrightarrow{G} \mathcal{J}$ between monoidal categories. Here by $\mathcal{M}_l(F) \times_{\mathcal{H}} \mathcal{M}_l(G)$ we mean the pseudo-pullback in the 2-category of monoidal categories: with objects (A, a, B, b, x) , where $(A, a) \in \mathcal{M}_l(F)$, $(B, b) \in \mathcal{M}_l(G)$, and $x : F(A) \longrightarrow B$ being an isomorphism in \mathcal{H} . A morphism $(A, a, B, b, x) \longrightarrow (A', a', B', b', x')$ is a pair (f, g) of morphisms $f : (A, a) \longrightarrow (A', a')$, $g : (B, b) \longrightarrow (B', b')$ in $\mathcal{M}_l(F)$, $\mathcal{M}_l(G)$ respectively, such that the square:

$$\begin{array}{ccc} F(A) & \xrightarrow{x} & B \\ F(f) \downarrow & & \downarrow g \\ F(A') & \xrightarrow{x'} & B' \end{array}$$

commutes. The tensor product in $\mathcal{M}_l(F) \times_{\mathcal{H}} \mathcal{M}_l(G)$ has the form:

$$(A, a, B, b, x) \otimes (A', a', B', b', x') = (A \otimes A', a|a', B \otimes B', b|b', x|x'),$$

where $x|x'$ is the composition

$$F(A \otimes A') \xrightarrow{a_{A'}} F(A) \otimes F(A') \xrightarrow{x \otimes x'} B \otimes B'.$$

Define a functor $\mathcal{M}_l(F, G) : \mathcal{M}_l(F) \times_{\mathcal{H}} \mathcal{M}_l(G) \longrightarrow \mathcal{M}_l(GF)$ by $(A, a, B, b, x) \mapsto (A, \tilde{a})$, where \tilde{a} is the composite:

$$\begin{aligned} GF(A \otimes X) &\xrightarrow{G(a_x)} G(F(A) \otimes F(X)) \xrightarrow{G(x \otimes F(X))} G(B \otimes F(X)) \\ &\xrightarrow{b_{F(X)}} G(B) \otimes GF(X) \xrightarrow{G(x)^{-1} \otimes GF(X)} GF(A) \otimes GF(X) \end{aligned}$$

3.7. PROPOSITION. *The functor $\mathcal{M}_l(F, G)$ is strict monoidal.*

PROOF. We need to show that

$$\begin{aligned} &\mathcal{M}_l(F, G)((A, a, B, b, x) \otimes (A', a', B', b', x')) = \\ &= \mathcal{M}_l(F, G)(A \otimes A', a|a', B \otimes B', b|b', x|x') = (A \otimes A', \widetilde{a|a'}) \end{aligned}$$

coincides with

$$\begin{aligned} &\mathcal{M}_l(F, G)(A, a, B, b, x) \otimes \mathcal{M}_l(F, G)(A', a', B', b', x') = \\ &= (A, \tilde{a}) \otimes (A', \tilde{a}') = (A \otimes A', \tilde{a}|\tilde{a}'). \end{aligned}$$


3.8. SUBCATEGORIES OF MULTIPLICANTS. Let \mathcal{G} , \mathcal{H} , \mathcal{C} be monoidal categories equipped with monoidal functors $F' : \mathcal{G} \longrightarrow \mathcal{C}$, $F'' : \mathcal{H} \longrightarrow \mathcal{C}$. Let $F : \mathcal{G} \longrightarrow \mathcal{H}$ be a functor and let $\gamma : F'' \circ F \longrightarrow F'$ be an isomorphism of functors. Define the full subcategory $\mathcal{M}_l(\gamma) = \mathcal{M}_l(F, F', F'', \gamma)$ in $\mathcal{M}_l(F)$ of objects (A, a) such that the following diagram is commutative

$$\begin{array}{ccc}
 F'(A \otimes X) & \xleftarrow{\gamma_{A \otimes X}} & F''F(A \otimes X) \\
 \downarrow F'_{A,X} & & \downarrow F''(a_X) \\
 & & F''(F(A) \otimes F(X)) \\
 & & \downarrow F'_{A,X} \\
 F'(A) \otimes F'(X) & \xleftarrow{\gamma_A \otimes \gamma_X} & F''F(A) \otimes F''F(X)
 \end{array}$$

3.9. THEOREM. *The subcategory $\mathcal{M}_l(\gamma)$ is closed under the tensor product in $\mathcal{M}_l(F)$. The natural isomorphism γ lifts to a monoidal natural isomorphism $\bar{\gamma}$ which fills the square*

$$\begin{array}{ccc}
 \mathcal{M}_l(\gamma) & \longrightarrow & \mathcal{H} \\
 \downarrow & & \downarrow F'' \\
 \mathcal{G} & \xrightarrow{F'} & \mathcal{C}
 \end{array}$$

PROOF. The fact that the tensor product $(A \otimes B, a|b)$ of two objects (A, a) , (B, b) of $\mathcal{M}_l(\gamma)$ belongs to $\mathcal{M}_l(\gamma)$ follows from the commutative diagram:



It follows from the definition of $\mathcal{M}_l(\gamma)$ that the natural transformation $\bar{\gamma}$ given by $\bar{\gamma}(A, a) = \gamma_A : F''F(A) \longrightarrow F'(A)$ is monoidal. ■

4. Examples, categories of modules

Here k denotes the ground field.

4.1. NUCLEI AND MULTIPLICANTS OF CATEGORIES OF MODULES. We start with the following technical statement.

4.2. LEMMA. *Let M be a left module over an associative algebra R and $m_{X_1, \dots, X_n} \in \text{End}(M \otimes X_1 \otimes \dots \otimes X_n)$ be a family of linear operators, which is natural in left R -modules X_i . Then m_{X_1, \dots, X_n} is given by multiplication by an element $m \in \text{End}(M) \otimes R \otimes \dots \otimes R$. Families of invertible operators correspond to invertible elements in the algebra $\text{End}(M) \otimes R \otimes \dots \otimes R$.*

PROOF. The R -module R is a generator in $R\text{-Mod}$. Thus any transformation m_{X_1, \dots, X_n} natural in $X_i \in R\text{-Mod}$ is defined by its specialisation $m_{R, \dots, R} \in \text{End}(M \otimes R \otimes \dots \otimes R)$. Naturality of m implies that $m_{R, \dots, R}$ commutes with $1 \otimes \text{End}_R(R) \otimes \dots \otimes \text{End}_R(R)$. Thus $m_{R, \dots, R}$ is given by multiplication by an element $m \in \text{End}(M) \otimes R \otimes \dots \otimes R$. ■

4.3. EXAMPLE. *Nucleus of a category of modules.*

Let R be an algebra and $\Delta : R \longrightarrow R \otimes R$ a homomorphism of algebras defining a magmoidal structure on the category $R\text{-Mod}$ of left R -modules. By lemma 4.2 the nucleus $\mathcal{N}_l(R\text{-Mod})$ is the category of pairs (M, m) where M is a left R -module and m is an invertible element of the tensor product $\text{End}(M) \otimes R \otimes R$ satisfying

$$(\Delta \otimes I)\Delta(r)m = m(I \otimes \Delta)\Delta(r) \tag{11}$$

for all $r \in R$. Here $\text{End}(M)$ is the endomorphism ring of the vector space M . The tensor product on the category $\mathcal{N}_l(R\text{-Mod})$ is defined by $(M, m) \otimes (N, n) = (M \otimes N, m|n)$ where $m|n \in \text{End}(M) \otimes \text{End}(N) \otimes R \otimes R$ is given by

$$(m|n) = (m \otimes 1)(I \otimes \Delta \otimes I)(m)(1 \otimes n)(I \otimes I \otimes \Delta)(m)^{-1}. \tag{12}$$

The tensor product is semigroupal with the associativity constraint:

$$\phi_{(M,m),(N,n),(L,l)} = (I \otimes \rho_N \otimes \rho_L)(m).$$

Here $\rho_N : R \longrightarrow \text{End}(N)$, $\rho_L : R \longrightarrow \text{End}(L)$ are the corresponding R -module structures. If $\varepsilon : R \longrightarrow k$ is a homomorphism of algebras (*counit*) such that

$$(\varepsilon \otimes I)\Delta = I = (I \otimes \varepsilon)\Delta,$$

the magmoidal category $R\text{-Mod}$ has a unit object $(k, 1)$. The monoidal subcategory $\mathcal{N}_l^1(R\text{-Mod})$ consists of those pairs (M, m) , where m satisfies the *normalisation* condition:

$$(I \otimes \varepsilon \otimes I)(m) = 1 = (I \otimes I \otimes \varepsilon)(m).$$

In the case when Δ is coassociative (when (R, Δ) is a bialgebra) the equation (11) means that m is R -invariant with respect to the diagonal inclusion of R into $\text{End}(M) \otimes R \otimes R$. The monoidal functor $R\text{-Mod} \longrightarrow \mathcal{N}_l(R\text{-Mod})$ corresponding to the canonical (trivial) associativity constraint on $R\text{-Mod}$, sends an R -module M into a pair $(M, 1)$.

4.4. EXAMPLE. *Multiplicant of a homomorphism.*

Let H_1, H_2 be bialgebras and let $f : H_1 \longrightarrow H_2$ be a homomorphism of algebras. It defines a functor $f^* : H_2\text{-Mod} \longrightarrow H_1\text{-Mod}$ between monoidal categories. Here we describe its (left) multiplicant $\mathcal{M}_l(f^*)$. Objects of $\mathcal{M}_l(f^*)$ are pairs (M, m) where M is H_2 -module and m is an invertible element of $\text{End}(M) \otimes H_2$ satisfying

$$m\Delta(f(h)) = (f \otimes f)\Delta(h)m \tag{13}$$

for all $h \in H_1$. The tensor product of pairs is given by the formula $(M, m) \otimes (N, n) = (M \otimes N, m|n)$ where $m|n \in \text{End}(M) \otimes \text{End}(N) \otimes H_2$ is defined by

$$(m|n) = (m \otimes 1)^{-1}(1 \otimes n)(I \otimes \Delta)(m).$$

The forgetful functor $\mathcal{M}_l(f^*) \rightarrow H_2\text{-Mod}$ sends a pair (M, m) to M . The functor $f^* : \mathcal{M}_l(f^*) \rightarrow H_1\text{-Mod}$, which sends (M, m) into $f^*(M)$, is semigroupal with respect to the constraint:

$$f^*_{(M,m),(N,n)} = (I \otimes \rho_N)(m) \in \text{End}(M) \otimes \text{End}(N).$$

Let $F_i : H_i\text{-Mod} \rightarrow \text{Vect}$ be the forgetful functors. An isomorphism $\gamma : F_1 \rightarrow F_2 f^*$ is given by multiplication with an invertible $x \in H_2$. The subcategory $\mathcal{M}_l(\gamma)$ of the multiplicant $\mathcal{M}_l(f^*)$ consists of the pairs (M, m) such that

$$m = (\rho_M \otimes I)(\Delta(x)^{-1}(x \otimes x)),$$

where $\rho_M : H_2 \rightarrow \text{End}(M)$ is the H_2 -action on M . Note that this together with (13) is equivalent to the condition:

$$(\rho_M \otimes I)(\Delta(g(h))) = (\rho_M \otimes I)((g \otimes g)\Delta(h)), \quad h \in H_1,$$

where $g : H_1 \rightarrow H_2$ is given by $g(h) = xf(h)x^{-1}$.

4.5. ALGEBRAIC CONSTRUCTIONS. For an associative algebra R consider the category $\mathcal{E}(R)$ of pairs (V, v) where V is a vector space and v is an element of $\text{End}(V) \otimes R$ with morphisms being vector space maps compatible with second components: a morphism $(V, v) \rightarrow (U, u)$ is a map $f : V \rightarrow U$ such that the equality

$$u(f \otimes 1) = (f \otimes 1)v \tag{14}$$

is valid in $\text{Hom}(V, U) \otimes R$. Denote by $\mathcal{A}(R)$ the full subcategory of $\mathcal{E}(R)$ of those pairs (V, v) for which v is invertible in $\text{End}(V) \otimes R$.

4.6. LEMMA. *Let R be a finite dimensional associative algebra. Then the category $\mathcal{E}(R)$ is equivalent to the category of modules over the free algebra $T(R^\vee)$ generated by the dual space of R . The category $\mathcal{A}(R)$ is equivalent to the category of modules over the quotient of the free algebra $T(R^\vee \oplus R^\vee)$ by the ideal generated by $\{\delta'(l) - l(1), \delta''(l) - l(1); l \in R^\vee\}$ where δ', δ'' are the composites of $\delta : R^\vee \rightarrow R^\vee \otimes R^\vee$, which is the dual map to the multiplication $R \otimes R \rightarrow R$, with two inclusions of $R^\vee \otimes R^\vee$ into $(R^\vee \oplus R^\vee)^{\otimes 2}$.*

PROOF. An element v of $\text{End}(V) \otimes R$ corresponds to a map $f_v : R^\vee \rightarrow \text{End}(V)$ sending $l \in R^\vee$ into $(I \otimes l)(v)$, and thus a $T(R^\vee)$ -module structure on V . Clearly compatibility condition (14) guarantees that a morphisms of pairs is $T(R^\vee)$ -linear. In the case when v is invertible the maps $f_v, f_{v^{-1}}$ satisfy the conditions:

$$\begin{aligned} (f_v \otimes f_{v^{-1}})\delta(l) &= (I \otimes \delta(l)(v_{12}v_{13}^{-1})) = (I \otimes l)(vv^{-1}) = l(1), \\ (f_{v^{-1}} \otimes f_v)\delta(l) &= (I \otimes \delta(l)(v_{12}^{-1}v_{13})) = (I \otimes l)(v^{-1}v) = l(1). \end{aligned}$$

■

Let $R = k(G)$ be an algebra of functions on a finite group G with standard multiplication Δ given by $\Delta(l)(f \otimes g) = l(fg)$ for $l \in k(G)$ and $f, g \in G$. By $p_g \in k(G)$ we denote the δ -function concentrated at $g \in G$. Note that δ -functions $p_g, g \in G$ form a basis in $k(G)$ so we can write $m \in \text{End}(M) \otimes k(G) \otimes k(G)$ as $\sum_{f,g \in G} m(f, g) \otimes p_f \otimes p_g$. The element m is invertible if $m(f, g) \in \text{End}(M)$ are invertible for any f, g . The $k(G)$ -invariance of m (11) means that $m(f, g)$ commutes with the image of $k(G)$ in $\text{End}(M)$. Thus we can identify the nucleus $\mathcal{N}_l(k(G)\text{-Mod})$ with the category $N_l(k(G))\text{-Mod}$ of left modules over the algebra $N_l(k(G)) = k(G) \otimes k[F(G \times G)]$ which is a tensor product of $k(G)$ and the group algebra of the free group $F(G \times G)$ generated by the cartesian product $G \times G$. Generators of $F(G \times G)$ will be denoted by $u(f, g)$. The pair (M, m) corresponds to the $N_l(k(G))$ action on M sending $u(f, g)$ into $m(f, g)$. The formula (12) for the tensor product of two pairs gives

$$(m|n)(f, g) = \sum_{h \in G} m(h, f)m(hf, g)m(h, fg)^{-1} \otimes p_h n(f, g)$$

which corresponds to the following comultiplication on $N_l(k(G))$

$$\Delta(p_h) = \sum_{fg=h} p_f \otimes p_g,$$

$$\Delta(u(f, g)) = \sum_{h \in G} u(h, f)u(hf, g)u(h, fg)^{-1} \otimes p_h u(f, g).$$

Note that $k(G)$ is a sub-bialgebra of $N_l(k(G))$. The coproduct on $N_l(k(G))$ is not coassociative but rather quasi-associative in the sense of [1]:

$$(I \otimes \Delta)(x) = \Phi(\Delta \otimes I)\Delta(x)\Phi^{-1}$$

where $\Phi = \sum_{f,g \in G} u(f, g) \otimes p_f \otimes p_g$ is an invertible element of $N_l(k(G))^{\otimes 3}$ (*associator*).

It is well known that associativity constraints on the category $k(G)\text{-Mod}$ are in one-to-one correspondence with 3-cocycles $Z^3(G, k^*)$ of the group G with coefficients in invertible elements of the ground field k . For any 3-cocycle $\alpha \in Z^3(G, k^*)$ there is defined a homomorphism of quasi-bialgebras $N_l(k(G)) \rightarrow k(G)$ splitting the inclusion $k(G) \rightarrow N_l(k(G))$ which sends $u(f, g)$ into $\sum_{h \in G} \alpha(h, f, g)p_h$. This homomorphism maps Φ into the associator $\Phi_\alpha = \sum_{h \in G} \alpha(h, f, g)p_h \otimes p_f \otimes p_g$ in $k(G)$ corresponding to α .

References

- [1] V. Drinfeld, Quasi-Hopf algebras. (Russian) Algebra i Analiz 1 (1989), no. 6, 114–148; translation in Leningrad Math. J. 1 (1990), no. 6, 1419–1457.
- [2] A. Joyal, R. Street, Tortile Yang-Baxter operators in tensor categories. Journal of Pure and Applied Algebra, 71 (1991), 43-51.

- [3] A. Joyal, R. Street, Braided tensor categories. *Adv. Math.* 102 (1993), no. 1, 20–78.
- [4] R. D. Schafer, *An Introduction to Nonassociative Algebras*, 1995 Courier Dover Publications, 176 pp.
- [5] J. Stasheff, *H-spaces from a homotopy point of view*. *Lecture Notes in Mathematics*, Vol. 161 Springer-Verlag, Berlin-New York 1970, 95 pp.

*Department of Mathematics, Division of Information and Communication Sciences
Macquarie University, Sydney, NSW 2109, Australia*

Email: davydov@math.mq.edu.au

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/18/16/18-16.{dvi,ps,pdf}>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION. Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS. The typesetting language of the journal is \TeX , and \LaTeX 2e strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

\TeX TECHNICAL EDITOR. Michael Barr, McGill University: mbarr@barrs.org

TRANSMITTING EDITORS.

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown \(at\) btinternet.com](mailto:ronnie.profbrown@btinternet.com)

Aurelio Carboni, Università dell' Insubria: aurelio.carboni@uninsubria.it

Valeria de Paiva, Xerox Palo Alto Research Center: paiva@parc.xerox.com

Ezra Getzler, Northwestern University: [getzler@northwestern\(dot\)edu](mailto:getzler@northwestern.edu)

Martin Hyland, University of Cambridge: M.Hyland@dpms.cam.ac.uk

P. T. Johnstone, University of Cambridge: ptj@dpms.cam.ac.uk

G. Max Kelly, University of Sydney: maxk@maths.usyd.edu.au

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, University of Western Sydney: s.lack@uws.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@acsu.buffalo.edu

Jean-Louis Loday, Université de Strasbourg: loday@math.u-strasbg.fr

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Brooke Shipley, University of Illinois at Chicago: bshipley@math.uic.edu

James Stasheff, University of North Carolina: jds@math.unc.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca