

## EXTENSIONS IN THE THEORY OF LAX ALGEBRAS

*Dedicated to Walter Tholen on the occasion of his 60th birthday*

CHRISTOPH SCHUBERT AND GAVIN J. SEAL

ABSTRACT. Recent investigations of lax algebras—in generalization of Barr’s relational algebras—make an essential use of lax extensions of monad functors on  $\mathbf{Set}$  to the category  $\mathbf{Rel}(\mathbf{V})$  of sets and  $\mathbf{V}$ -relations (where  $\mathbf{V}$  is a unital quantale). For a given monad there may be many such lax extensions, and different constructions appear in the literature. The aim of this article is to shed a unifying light on these lax extensions, and present a symptomatic situation in which distinct monads yield isomorphic categories of lax algebras.

### 1. Introduction

In addition to a monad  $\mathbb{T}$  on  $\mathbf{Set}$  and a unital quantale  $\mathbf{V}$ , the definition of  $(\mathbb{T}, \mathbf{V})$ -algebras requires a lax extension of the monad functor  $T$  to  $\mathbf{V}\text{-Rel}$ , the 2-category of sets and  $\mathbf{V}$ -relations. Until recently, such extensions were obtained by using Barr’s original construction [2] (indeed, if  $\mathbf{2}$  denotes the two-chain  $\{\perp, \top\}$ , then  $\mathbf{2}\text{-Rel}$  is just the category  $\mathbf{Rel}$  of sets and relations). In [3], Clementino and Hofmann describe a process that essentially produces a lax extension to  $\mathbf{V}\text{-Rel}$  out of a lax extension to  $\mathbf{Rel}$ . Two other constructions were proposed by Seal: the first, in [19], was based on the assumption that  $\mathbb{T}$  was a taut monad, while the second, in [20], required that the monad be conveniently  $\mathbf{Sup}$ -enriched. In other directions, Schubert [18] showed that Barr’s method could be adapted to an order-based setting, and Hofmann [10] generalized the original construction by exploiting the structure of a particular Eilenberg-Moore algebra on  $\mathbf{V}$ .

These extensions all yield topological spaces as instances of lax algebras, but they are nonetheless very different in nature, and therefore in their realizations. Roughly put, the lax extensions obtained via Barr’s construction are  $\mathbf{Set}$ -based and do not take into account any order-related considerations, on the contrary of those introduced by Schubert, which are intrinsically  $\mathbf{Ord}$ -based; the lax extensions studied by Seal live in-between the previous two, as they are originally defined over  $\mathbf{Set}$ , but nonetheless exploit an existing underlying

---

The first author gratefully acknowledges a Ph.D.-scholarship from the Ernst A.-C. Lange Stiftung, Bremen. The second author gratefully acknowledges financial support by the Swiss National Science Foundation.

Received by the editors 2007-09-20 and, in revised form, 2008-11-06.

Published on 2008-11-07 in the Tholen Festschrift.

2000 Mathematics Subject Classification: 18C20, 18B30, 54A05.

Key words and phrases: lax algebra, Kleisli extension, initial extension, strata extension, tower extension.

© Christoph Schubert and Gavin J. Seal, 2008. Permission to copy for private use granted.

order of the monad. Thus, even in the simple case where  $\mathbf{V} = \mathbf{2}$ , these constructions lead to very different lax extensions (a striking illustration of this is given in Example 4.6(3)). The intent of this article is to study the *Kleisli extension* introduced in [20] for **Sup**-enriched monads, and its interaction with four other types of extensions: the *initial extension* of a **Set**-functor described below, the *canonical and op-canonical extensions* of a taut **Set**-functor defined in [19], the *strata extension* of a lax **Rel**-functor [3], and the *tower extension* of a topological category [21].

The multiplicity of lax extensions is a manifestation of a rich ambient categorical structure; indeed, for a fixed  $\mathbf{V}$ , lax extensions are the objects of a “topological quasicategory” over the quasicategory of **Set**-endofunctors (see 3.5 for details). This remark leads us to a central theme of our work: if  $\mathbb{T}$  is provided with a lax extension  $\overline{T}$ , then any monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  allows for a lax extension  $\overline{S}$  of  $S$  via the initial lift of  $\alpha : S \rightarrow \overline{T}$ . For example, the “op-canonical” extension of a taut functor described in [19] is obtained as an initial lift of the filter functor’s Kleisli extension.

More generally, if one considers for  $\overline{T}$  the Kleisli extension of  $T$ , the initial lift construction allows us to identify the category  $\mathbf{Alg}(\mathbb{S}, \mathbf{2})$  of  $(\mathbb{S}, \mathbf{2})$ -algebras with the conceptually simpler category  $\mathbf{KIMon}(\mathbb{T})$  of Kleisli monoids. This is the theme of Theorem 5.6. The strata extension of  $\overline{S}$  then provides a suitable ingredient to pass from  $\mathbf{Alg}(\mathbb{S}, \mathbf{2})$  to  $\mathbf{Alg}(\mathbb{S}, \mathbf{V})$ , while the tower extension of  $\mathbf{KIMon}(\mathbb{T})$  yields a category  $\mathbf{KIMon}(\mathbb{T}, \mathbf{V})$ . Theorem 6.10 exploits the previous result to show that these new categories are again isomorphic, emphasizing in the process the importance of the original  $\mathbf{V} = \mathbf{2}$  case.

The examples used throughout this work support the thesis that “topological-related” lax extensions tend to appear either as Kleisli extensions, or as initial lifts of these (Example 4.6(2)). A similar line of investigation has been followed by Colebunders and Lowen in [5], where the authors show that many important lax extensions are initial extensions of a certain “functional power monad”.

## 2. The setting

2.1. **QUANTALES.** Throughout this article,  $\mathbf{V} = (\mathbf{V}, \otimes, k)$  denotes a unital quantale with tensor  $\otimes$ , and unit  $k$ . In other words,  $\mathbf{V}$  is a complete lattice equipped with an associative binary operation  $\otimes$  that preserves suprema in each variable, and admits  $k$  as neutral element:

$$\left(\bigvee_{i \in I} u_i\right) \otimes v = \bigvee_{i \in I} (u_i \otimes v) \quad , \quad u \otimes \left(\bigvee_{i \in I} v_i\right) = \bigvee_{i \in I} (u \otimes v_i) \quad , \quad k \otimes v = v = v \otimes k \quad ,$$

for all  $u, v, u_i, v_i \in \mathbf{V}$  ( $i \in I$ ). The bottom and top elements of  $\mathbf{V}$  are denoted by  $\perp$  and  $\top$ , respectively. In this article, we always suppose that  $\mathbf{V}$  is *non-trivial*, that is,  $\mathbf{V}$  is not a singleton (or equivalently,  $\perp \neq k$ ). Moreover, the quantale is said to be *integral* if  $k = \top$ .

### 2.2. EXAMPLES.

1. The *two-chain*  $\mathbf{2} = (\{\perp, \top\}, \wedge, \top)$  is the simplest non-trivial quantale. Of course, any frame is a unital quantale with binary meet and neutral element  $\top$ , but  $\mathbf{2}$  already leads to interesting examples of lax algebras.

2. The *diamond lattice*  $\mathbf{2}^2 = (\{\perp, u, v, \top\}, \wedge, \vee)$  (with  $u$  and  $v$  incomparable) is obviously isomorphic to the powerset of  $\mathbf{2}$ , and provides another meaningful example of a non-trivial quantale.
3. The *extended real line*  $\mathbf{P}_+ = ([0, \infty]^{\text{op}}, +, 0)$  with its addition extended via  $x + \infty = \infty = \infty + x$  for all  $x \in [0, \infty]$  is a quantale. Notice that the opposite order has been taken on the chain  $[0, \infty]$ , so that the neutral element  $0$  is also the top element of the lattice.
4. The *non-integral three-chain*  $\mathbf{3} = (\{\perp, k, \top\}, \otimes, k)$  is the smallest non-integral quantale. The tensor is commutative, and necessarily defined by  $\perp \otimes v = \perp$  and  $k \otimes v = v$  for all  $v \in \mathbf{3}$ , while  $\top \otimes \top = \top$ .

2.3. **V-RELATIONS AND RELATIONS.** The objects of the category  $\mathbf{V}\text{-Rel}$  are sets, and its morphisms  $r : X \leftrightarrow Y$  are **V-relations**, that is, maps of the form  $r : X \times Y \rightarrow \mathbf{V}$ . Composition of  $r : X \leftrightarrow Y$  with  $s : Y \leftrightarrow Z$  is given by the “matrix multiplication” formula:

$$(s \cdot r)(x, z) = \bigvee_{y \in Y} r(x, y) \otimes s(y, z) .$$

The identity  $1_X : X \leftrightarrow X$  is the **V-relation** defined by  $1_X(x, y) = k$  if  $x = y$  and  $1_X(x, y) = \perp$  otherwise. The category  $\text{Rel}$  of sets and relations is identified with  $\mathbf{2}\text{-Rel}$ , and embeds into  $\mathbf{V}\text{-Rel}$  via composition of  $r : X \times Y \rightarrow \mathbf{2}$  with

$$\lambda : \mathbf{2} \rightarrow \mathbf{V} , \quad \perp \mapsto \perp , \quad \top \mapsto k .$$

A **V-relation** with values in  $\{\perp, k\}$  is called a *relation in V-Rel*, while the term *relation* alone means a **2-relation**.

Similarly,  $\text{Set}$  can be embedded into  $\mathbf{V}\text{-Rel}$  by sending each map  $f : X \rightarrow Y$  to its *graph* (or more precisely its **V-graph**), defined by

$$f(x, y) = \begin{cases} k & \text{if } f(x) = y \\ \perp & \text{otherwise} . \end{cases}$$

Note that we will not insist on distinguishing between a map and its graph, since the context will always determine which level we are working on.

Equipped with the pointwise order induced by  $\mathbf{V}$ , the hom-sets of  $\mathbf{V}\text{-Rel}$  become sup-semilattices, and we have

$$r \leq r' , \quad s \leq s' \implies s \cdot r \leq s' \cdot r'$$

for all  $r, r' : X \leftrightarrow Y$  and  $s, s' : Y \leftrightarrow Z$ . The *transpose*  $r^\circ : X \leftrightarrow Y$  of a **V-relation**  $r : X \leftrightarrow Y$  is defined by  $r^\circ(y, x) = r(x, y)$ . Since the quantale  $\mathbf{V}$  is not necessarily commutative, we do *not* have in general that  $(s \cdot r)^\circ = r^\circ \cdot s^\circ$  in  $\mathbf{V}\text{-Rel}$ , although this formula does hold if either  $r$  or  $s$  is a relation in  $\mathbf{V}\text{-Rel}$ .

By composing a map  $f : X \rightarrow Y$ , a  $\mathbf{V}$ -relation  $s : Y \leftrightarrow Z$ , and the transpose of  $h : W \rightarrow Z$ , we get the convenient formula

$$h^\circ \cdot s \cdot f(x, w) = s(f(x), h(w)) .$$

Notice also that  $1_X \leq f^\circ \cdot f$  and  $f \cdot f^\circ \leq 1_X$ , so the following equivalences hold

$$t \leq s \cdot f \iff t \cdot f^\circ \leq s \quad \text{and} \quad g \cdot r \leq t \iff r \leq g^\circ \cdot t$$

for any  $\mathbf{V}$ -relation  $t : X \leftrightarrow Z$  and map  $g : Y \rightarrow Z$ .

2.4. STRATA. Let  $\mathbf{Inf}$  denote the category of inf-semilattices and inf-preserving maps. For a  $\mathbf{V}$ -relation  $r : X \leftrightarrow Y$  and  $v \in \mathbf{V}$ , the  $v$ -stratum of  $r$  is the relation  $r_v : X \leftrightarrow Y$  defined by

$$r_v(x, y) = \top \iff v \leq r(x, y) .$$

In the case where  $s : X \leftrightarrow Y$  is a relation in  $\mathbf{V}\text{-Rel}$ , the relation  $s_k$  is simply the preimage of  $s$  via the embedding  $\mathbf{Rel} \hookrightarrow \mathbf{V}\text{-Rel}$ .

For any  $\mathcal{A} \subseteq \mathbf{V}$ , we have  $r_{\bigvee \mathcal{A}} = \bigwedge_{v \in \mathcal{A}} r_v$ , so the map

$$\phi_r : \mathbf{V}^{\text{op}} \rightarrow \mathbf{Rel}(X, Y) , \quad v \mapsto r_v$$

preserves arbitrary infima. On the other hand, every inf-map  $\phi : \mathbf{V}^{\text{op}} \rightarrow \mathbf{Rel}(X, Y)$  yields a  $\mathbf{V}$ -relation

$$r_\phi : X \leftrightarrow Y , \quad (x, y) \mapsto \bigvee \{ v \in \mathbf{V} \mid \phi(v)(x, y) = \top \} .$$

These correspondences describe an order-preserving isomorphism

$$\mathbf{V}\text{-Rel}(X, Y) \cong \mathbf{Inf}(\mathbf{V}^{\text{op}}, \mathbf{Rel}(X, Y)) .$$

Furthermore, for any maps  $f : X \rightarrow Y$ ,  $g : Y \rightarrow Z$ ,  $\mathbf{V}$ -relations  $r : X \leftrightarrow Y$ ,  $s : Y \leftrightarrow Z$  and  $u, v \in \mathbf{V}$ , we have

$$s_v \cdot f = (s \cdot f)_v , \quad g^\circ \cdot r_v = (g^\circ \cdot r)_v \quad \text{and} \quad s_u \cdot r_v \leq (s \cdot r)_{v \otimes u} .$$

2.5. REMARK. As pointed out to the authors by Walter Tholen, the previous isomorphism is a particular case of the order-preserving isomorphism

$$\mathbf{Set}(A, \mathbf{V}) \cong \mathbf{Inf}(\mathbf{V}^{\text{op}}, PA) .$$

Indeed, whenever  $A = X \times Y$ , we have  $\mathbf{Set}(A, \mathbf{V}) = \mathbf{V}\text{-Rel}(X, Y)$ , and  $PA \cong \mathbf{Rel}(X, Y)$ . Moreover, this isomorphism is the restriction to fixpoints of the adjunction

$$\phi \dashv \psi : \mathbf{Set}(A, \mathbf{V}) \rightarrow \mathbf{Set}(\mathbf{V}, PA) ,$$

where  $\mathbf{Set}(A, \mathbf{V})$  and  $\mathbf{Set}(\mathbf{V}, PA)$  are endowed with their respective pointwise order, and

$$\phi(f)(a) := \bigvee \{ v \in \mathbf{V} \mid a \in f(v) \} , \quad \psi(g)(v) := \{ a \in A \mid v \leq g(a) \}$$

for all maps  $f : \mathbf{V} \rightarrow PA$ ,  $g : A \rightarrow \mathbf{V}$ , and elements  $a \in A$ ,  $v \in \mathbf{V}$ .

2.6. **MONADS.** Recall that a monad  $\mathbb{T}$  on  $\mathbf{Set}$  is a triple  $(T, \eta, \mu)$ , where  $T : \mathbf{Set} \rightarrow \mathbf{Set}$  is a functor, and the *unit*  $\eta : \text{Id} \rightarrow T$  and *multiplication*  $\mu : T^2 \rightarrow T$  of  $\mathbb{T}$  are natural transformations satisfying

$$\mu \cdot T\eta = 1 = \mu \cdot \eta T \quad \text{and} \quad \mu \cdot T\mu = \mu \cdot \mu T .$$

A *monad morphism*  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  from  $\mathbb{S} = (S, \delta, \nu)$  to  $\mathbb{T} = (T, \eta, \mu)$  is a natural transformation  $\alpha : S \rightarrow T$  such that

$$\eta = \alpha \cdot \delta \quad \text{and} \quad \mu \cdot \alpha^2 = \alpha \cdot \nu \quad (\text{with } \alpha^2 = T\alpha \cdot \alpha S = \alpha T \cdot S\alpha) .$$

We say that  $\alpha$  is an *embedding* if the components  $\alpha_X$  are injections.

2.7. **EXAMPLES.** The following monads will be used throughout the text.

1. The *identity monad* is the obvious monad  $\mathbb{I} = (\text{Id}, 1, 1)$ .
2. The *powerset monad* is  $\mathbb{P} = (P, \iota, \bigcup)$  with unit given by  $\iota_X(x) := \{x\}$ . Note that for  $A \in PX$  and a map  $f : X \rightarrow Y$  we will often write

$$f[A] := \{f(x) \mid x \in A\},$$

instead of  $Pf(A)$ .

3. The *double-dualization monad* (also called the “*contravariant*” *double-powerset monad*)  $\mathbb{D}_2 = (D_2, \dot{-}, \Sigma)$  is described by  $D_2X := PPX$  for every set  $X$ , together with the following three equivalences:

$$B \in D_2f(\mathfrak{f}) \iff f^{-1}(B) \in \mathfrak{f}, \quad A \in \dot{x} \iff x \in A, \quad A \in \Sigma_X(\mathfrak{F}) \iff A^{\mathbb{D}_2} \in \mathfrak{F},$$

for  $f : X \rightarrow Y$ ,  $\mathfrak{f} \in D_2X$ ,  $x \in X$ ,  $\mathfrak{F} \in D_2D_2X$ , and where

$$A^{\mathbb{D}_2} := \{\mathfrak{f} \in D_2X \mid A \in \mathfrak{f}\} .$$

It will also be convenient to use the following notation:

$$f[\mathfrak{f}] := D_2f[\mathfrak{f}] .$$

4. An element  $\mathfrak{f} \in D_2X = PPX$  is *up-closed* if  $A \in \mathfrak{f}$  and  $A \subseteq B$  implies  $B \in \mathfrak{f}$  for all  $B \subseteq X$ . For any  $\mathfrak{f} \subseteq PX$ , the *up-closure* of  $\mathfrak{f}$  is defined by

$$\uparrow\mathfrak{f} := \{B \subseteq X \mid \exists A \in \mathfrak{f} : A \subseteq B\} .$$

The *up-set monad*  $\mathbb{U} = (U, \dot{-}, \Sigma)$  is just the restriction of  $\mathbb{D}_2$  to up-closed elements of  $D_2X$ . In this case,  $\Sigma_X(\mathfrak{F}) = \{A^{\mathbb{U}} \mid A^{\mathbb{U}} \in \mathfrak{F}\}$  for  $\mathfrak{F} \in UUX$ , where  $A^{\mathbb{U}} := \{f \in UX \mid A \in f\}$ .

5. The *filter monad*  $\mathbb{F} = (F, \dot{-}, \Sigma)$  is the restriction of the up-set monad to filters. Here, we must use  $A^{\mathbb{F}} := \{f \in FX \mid A \in f\}$  in lieu of  $A^{\mathbb{U}}$  in the definition of the components of  $\Sigma : FF \rightarrow F$ . Let us point out that we consider the improper filter  $f = PX$  as an element of  $FX$ .
6. The *ultrafilter monad*  $\mathbb{B} = (\beta, \dot{-}, \Sigma)$  is the restriction of the filter monad to ultrafilters. Here we use the sets  $A^{\mathbb{B}} := \{x \in \beta X \mid A \in x\}$  instead of  $A^{\mathbb{F}}$ .
7. The previous monads may all be considered as submonads of  $\mathbb{D}_2$ . Indeed, the power-set monad  $\mathbb{P}$  embeds into  $\mathbb{F}$  via the principal filter natural transformation  $\tau : P \rightarrow F$ , defined componentwise by  $\tau_X(A) := \uparrow\{A\}$  for  $A \in PX$ , and the identity monad  $\mathbb{I}$  obviously embeds both into  $\mathbb{P}$  and  $\mathbb{B}$ . Therefore, we have the following diagram of monad embeddings:

$$\begin{array}{ccccccc} & & & \mathbb{P} & & & \\ & \nearrow & & \searrow & & & \\ \mathbb{I} & & & & & \mathbb{F} & \longrightarrow \mathbb{U} \longrightarrow \mathbb{D}_2 . \\ & \searrow & & \nearrow & & & \\ & & & \mathbb{B} & & & \end{array}$$

2.8. MONADS AS KLEISLI TRIPLES. For our purpose, the alternate description of monads from ([15], Exercise 1.3.12) will be useful. A *Kleisli triple*  $(T, \eta, -^{\mathbb{T}})$  on  $\mathbf{Set}$  consists of

- (i) a map  $T : \text{ob Set} \rightarrow \text{ob Set}$ ,
- (ii) for each set  $X$ , a map  $\eta_X : X \rightarrow TX$ ,
- (iii) an operation  $-^{\mathbb{T}}$  which sends  $f : X \rightarrow TY$  to  $f^{\mathbb{T}} : TX \rightarrow TY$ ,

subject to the conditions

$$(\eta_X)^{\mathbb{T}} = 1_{TX} \quad , \quad f^{\mathbb{T}} \cdot \eta_X = f \quad \text{and} \quad g^{\mathbb{T}} \cdot f^{\mathbb{T}} = (g^{\mathbb{T}} \cdot f)^{\mathbb{T}} .$$

Each Kleisli triple  $(T, \eta, -^{\mathbb{T}})$  yields a monad  $\mathbb{T} = (T, \eta, \mu)$  by setting

$$Tf := (\eta_Y \cdot f)^{\mathbb{T}} \quad \text{and} \quad \mu_X := (1_{TX})^{\mathbb{T}} ,$$

and every monad  $\mathbb{T} = (T, \eta, \mu)$  defines a Kleisli triple via

$$f^{\mathbb{T}} := \mu_Y \cdot Tf .$$

Since these processes are mutually inverse, we will freely switch between the two descriptions.

Given Kleisli triples  $(S, \delta, -^{\mathbb{S}})$  and  $(T, \eta, -^{\mathbb{T}})$ , a family  $(\alpha_X : SX \rightarrow TX)_{X \in \text{ob Set}}$  defines a monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  if and only if the equalities

$$\alpha_X \cdot \delta_X = \eta_X \quad \text{and} \quad (\alpha_Y \cdot f)^{\mathbb{T}} \cdot \alpha_X = \alpha_Y \cdot f^{\mathbb{S}}$$

hold for all sets  $X$  and maps  $f : X \rightarrow SY$ .

2.9. **KLEISLI CATEGORIES.** We recall that the *Kleisli category*  $\mathbf{Set}_{\mathbb{T}}$  of a monad  $\mathbb{T} = (T, \eta, -^{\mathbb{T}})$  has as objects sets, and as morphisms  $f : X \rightarrow Y$  maps  $f : X \rightarrow TY$ . The composite of  $f : X \rightarrow Y$  with  $g : Y \rightarrow Z$  is given via **Set**-composition as

$$g \circ f := g^{\mathbb{T}} \cdot f = \mu_Z \cdot Tg \cdot f ,$$

and the identity at  $X$  is  $\eta_X : X \rightarrow X$ . Observe that the conditions on a Kleisli triple are equivalent to the left and right unit laws of the identity, and to the associativity of composition, respectively.

Every monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  induces a functor  $\mathbf{Set}_{\mathbb{S}} \rightarrow \mathbf{Set}_{\mathbb{T}}$  by composing morphisms with  $\alpha_X$ . The conditions for monad morphisms between Kleisli triples are then equivalent to preservation of identities and composition, respectively.

The Kleisli category  $\mathbf{Set}_{\mathbb{P}}$  of the powerset monad is the category **Rel** of sets and relations. Using the isomorphism  $\mathbf{Rel} \cong \mathbf{Rel}^{\text{op}}$ , we will exploit the description of a relation  $r : X \leftrightarrow Y$  by its preimage-mapping

$$r^{\flat} : Y \rightarrow PX , \quad y \mapsto \{ x \mid r(x, y) = \top \}$$

(notice the transposition). This description is functorial since for  $s : Y \leftrightarrow Z$ , we have

$$(s \cdot r)^{\flat} = (r^{\flat})^{\mathbb{P}} \cdot s^{\flat} = r^{\flat} \circ s^{\flat} .$$

When  $f : X \leftrightarrow Y$  is the graph of a map, the preimage-mapping is related to well-known operations on  $f$ :

$$(f^{\flat})^{\mathbb{P}} = f^{-1} : PY \rightarrow PX \quad \text{and} \quad (f^{\circ})^{\flat} = \iota_Y \cdot f : X \rightarrow PY .$$

In particular, we have  $((f^{\circ})^{\flat})^{\mathbb{P}} = Pf : PX \rightarrow PY$ .

### 3. Lax extensions and lax algebras

3.1. **LAX EXTENSIONS.** A *lax extension* of a **Set**-functor  $T : \mathbf{Set} \rightarrow \mathbf{Set}$  along the embedding  $\mathbf{Set} \hookrightarrow \mathbf{V}\text{-Rel}$ , (or simply a *lax extension* of  $T$  to  $\mathbf{V}\text{-Rel}$ ), is a map

$$\bar{T} : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel} \quad , \quad (r : X \leftrightarrow Y) \mapsto (\bar{T}r : TX \leftrightarrow TY)$$

satisfying for all  $r : X \leftrightarrow Y$ ,  $s : Y \leftrightarrow Z$ , and  $f : X \rightarrow Y$  the conditions

$$(i) \quad s \leq r \implies \bar{T}s \leq \bar{T}r ,$$

$$(ii) \quad \bar{T}s \cdot \bar{T}r \leq \bar{T}(s \cdot r) ,$$

$$(iii) \quad Tf \leq \bar{T}f \text{ and } (Tf)^{\circ} \leq \bar{T}f^{\circ} .$$

The last condition implies in particular that  $1_{TX} \leq \bar{T}1_X$ , so a lax extension of  $T : \mathbf{Set} \rightarrow \mathbf{Set}$  is simply a lax functor satisfying the extension conditions (iii). Notice that a lax extension is a structure on a functor and not a mere property. A *morphism* of lax extensions  $\alpha : (S, \bar{S}) \rightarrow (T, \bar{T})$  is a natural transformation  $\alpha : S \rightarrow T$  between the underlying  $\mathbf{Set}$ -functors which extends to an *oplax transformation*  $\bar{S} \rightarrow \bar{T}$ ; that is,  $\alpha_Y \cdot \bar{S}r \leq \bar{T}r \cdot \alpha_X$ , or equivalently

$$\bar{S}r \leq \alpha_Y^\circ \cdot \bar{T}r \cdot \alpha_X,$$

for all  $r : X \rightarrow Y$ . Often, we simply write  $\alpha : \bar{S} \rightarrow \bar{T}$  instead of  $\alpha : (S, \bar{S}) \rightarrow (T, \bar{T})$ .

If  $\bar{T} : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel}$  is a lax extension of  $T : \mathbf{Set} \rightarrow \mathbf{Set}$ , then it satisfies

$$\bar{T}(s \cdot f) = \bar{T}s \cdot \bar{T}f = \bar{T}s \cdot Tf \quad \text{and} \quad \bar{T}(g^\circ \cdot s) = \bar{T}g^\circ \cdot \bar{T}s = (Tg)^\circ \cdot \bar{T}s,$$

for all  $s : Y \rightarrow Z$ ,  $f : X \rightarrow Y$ , and  $g : W \rightarrow Z$ . Indeed, we have

$$\bar{T}(s \cdot f) \leq \bar{T}(s \cdot f) \cdot \bar{T}f^\circ \cdot Tf \leq \bar{T}(s \cdot f \cdot f^\circ) \cdot Tf \leq \bar{T}s \cdot Tf \leq \bar{T}s \cdot \bar{T}f \leq \bar{T}(s \cdot f),$$

and the other set of equalities follows in a similar way.

**3.2. ASSOCIATED PREORDERS.** Any lax functor  $T : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel}$  yields a preorder on the sets  $TX$  via

$$\mathfrak{x} \leq_T \mathfrak{y} \iff k \leq T1_X(\mathfrak{x}, \mathfrak{y})$$

for all  $\mathfrak{x}, \mathfrak{y} \in TX$ . A lax extension  $\bar{T} : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel}$  of a functor  $T : \mathbf{Set} \rightarrow \mathbf{Set}$  takes this process one step further and induces a factorization of  $T$  through the category  $\mathbf{Ord}$  of preordered sets and monotone maps. Indeed, a map  $f : X \rightarrow Y$  is sent to a monotone map  $Tf : TX \rightarrow TY$ , since

$$\bar{T}1_X \leq \bar{T}(f^\circ \cdot 1_Y \cdot f) = (Tf)^\circ \cdot \bar{T}1_Y \cdot Tf.$$

If  $\alpha : \bar{S} \rightarrow \bar{T}$  is a morphism of lax extensions, then each  $\alpha_X : SX \rightarrow TX$  is monotone. Let us stress moreover that  $\leq_{\bar{T}}$  is a *relation*, while  $\bar{T}1_X$  is a  $\mathbf{V}$ -relation so that  $\leq_{\bar{T}} \neq \bar{T}1_X$  even if  $\leq_{\bar{T}}$  is seen as a  $\mathbf{V}$ -relation (this distinction will be important in the definition of an antitone  $\mathbf{V}$ -relation in 5.4).

When the sets  $TX$  are equipped with this order, the  $\mathbf{V}$ -relation  $\bar{T}r$  reverses it in its first variable and preserves it in its second; indeed, if  $\mathfrak{x}' \leq_{\bar{T}} \mathfrak{x}$  and  $\mathfrak{y} \leq_{\bar{T}} \mathfrak{y}'$ , then

$$\bar{T}r(\mathfrak{x}, \mathfrak{y}) \leq \bar{T}1_X(\mathfrak{x}', \mathfrak{x}) \otimes \bar{T}r(\mathfrak{x}, \mathfrak{y}) \otimes \bar{T}1_Y(\mathfrak{y}, \mathfrak{y}') \leq \bar{T}1_Y \cdot \bar{T}r \cdot \bar{T}1_X(\mathfrak{x}', \mathfrak{y}') = \bar{T}r(\mathfrak{x}', \mathfrak{y}')$$

for all  $\mathfrak{x}, \mathfrak{x}' \in TX$ ,  $\mathfrak{y}, \mathfrak{y}' \in TY$ .



3.3. INITIAL EXTENSIONS. Let  $\alpha : S \rightarrow T$  be a natural transformation of **Set**-functors, and  $\bar{T}$  a lax extension of  $T$  to **V-Rel**. The *initial extension* of  $S$  along  $\alpha$  is the lax extension given by

$$\alpha^*\bar{T}r := \alpha_Y^\circ \cdot \bar{T}r \cdot \alpha_X ,$$

for any **V**-relation  $r : X \leftrightarrow Y$ . The morphism of lax extensions  $\alpha : (S, \alpha^*\bar{T}) \rightarrow (T, \bar{T})$  is *initial* in the following sense: if  $\beta : R \rightarrow S$  is a natural transformation and  $\bar{R}$  is a lax extension of  $R$ , then  $\beta : \bar{R} \rightarrow \alpha^*\bar{T}$  is a morphism of lax extensions if and only if  $\alpha \cdot \beta : \bar{R} \rightarrow \bar{T}$  is one.

In presence of the initial extension  $\alpha^*\bar{T}$  of  $S$ , the maps  $\alpha_X : SX \rightarrow TX$  become *order-embeddings* with respect to the associated preorders:

$$\mathfrak{x} \leq_{\alpha^*\bar{T}} \mathfrak{x}' \iff \alpha_X(\mathfrak{x}) \leq_{\bar{T}} \alpha_X(\mathfrak{x}')$$

for all  $\mathfrak{x}, \mathfrak{x}' \in SX$ .

3.4. PROPOSITION. Let  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  be a morphism of **Set**-monads  $\mathbb{S} = (S, \delta, \nu)$  and  $\mathbb{T} = (T, \eta, \mu)$ . If  $T$  has a lax extension  $\bar{T}$  to **V-Rel** for which  $\eta : \text{Id} \rightarrow \bar{T}$  is oplax, then  $\delta : \text{Id} \rightarrow \bar{S}$  is also oplax for the initial extension  $\bar{S} = \alpha^*\bar{T}$  of  $S$  along  $\alpha$ . Similarly, if  $\mu : \bar{T}\bar{T} \rightarrow \bar{T}$  is oplax for  $\bar{T}$ , then  $\nu : \bar{S}\bar{S} \rightarrow \bar{S}$  is oplax for  $\bar{S} = \alpha^*\bar{T}$ .

PROOF. Oplaxness of  $\delta$  follows immediately from  $\eta = \alpha \cdot \delta$ . Since  $\alpha : \bar{S} \rightarrow \bar{T}$  is oplax,  $\alpha^2 : \bar{S}\bar{S} \rightarrow \bar{T}\bar{T}$  is too. Thus, oplaxness of  $\nu$  follows from  $\mu \cdot \alpha^2 = \alpha \cdot \nu$ . ■

3.5. REMARK. For a source  $\alpha = (\alpha_i : S \rightarrow T_i)_{i \in I}$ , where each  $T_i$  carries a lax extension  $\bar{T}_i$  to **V-Rel**, we can define an initial lift via the lax extension  $\bar{S}$  of  $S$  defined by

$$\bar{S}r := \bigwedge_{i \in I} (\alpha_i)_Y^\circ \cdot \bar{T}_i r \cdot (\alpha_i)_X ,$$

for any **V**-relation  $r : X \leftrightarrow Y$ . In other words, the quasicategory<sup>1</sup> of lax extensions to **V-Rel** is topological over the quasicategory of **Set**-functors.

3.6. EXAMPLES.

1. A lax extension of the identity functor  $\text{Id} : \mathbf{Set} \rightarrow \mathbf{Set}$  to **V-Rel** is given by the identity  $\text{Id} : \mathbf{V-Rel} \rightarrow \mathbf{V-Rel}$ . In this case, the preorder associated with this extension is the discrete order on  $X$ .
2. A lax extension of the powerset functor  $P : \mathbf{Set} \rightarrow \mathbf{Set}$  to **V-Rel** is given by

$$\bar{P}r(A, B) = \bigvee \{ v \in V \mid A \subseteq r_v^b[B] \}$$

for any  $A \in PX$ ,  $B \in PY$ , and **V**-relation  $r : X \leftrightarrow Y$ . The preorder associated with this extension is just subset inclusion. Moreover, the previous lax extension of the identity functor is the initial extension induced by the unit  $\iota : \text{Id} \rightarrow P$  of  $\mathbb{P}$ .

<sup>1</sup>Due to size restrictions on the hom-sets, the term “quasicategory” is used in lieu of “category”. This distinction is made here because topologicity is classically defined for locally small categories only [1].

3. A lax extension of the up-set functor  $U : \mathbf{Set} \rightarrow \mathbf{Set}$  to  $\mathbf{V}\text{-Rel}$  is given by

$$\bar{U}r(\mathfrak{f}, \mathfrak{g}) = \bigvee \{ v \in V \mid \mathfrak{f} \supseteq r_v^b[\mathfrak{g}] \}$$

for  $r : X \leftrightarrow Y$ ,  $\mathfrak{f} \in UX$ , and  $\mathfrak{g} \in UY$ . The preorder obtained on  $UX$  is the “finer than” relation, which is opposite to subset inclusion. One obtains lax extensions of the filter and ultrafilter functors by restricting this lax extension accordingly. The extensions obtained are the initial lifts along the corresponding embeddings (see Example 2.7(7)). Similarly, the extension  $\bar{P}$  described above is the initial lift of the restriction  $\bar{F}$  along the principal filter transformation  $\tau : \mathbb{P} \rightarrow \mathbb{F}$ .

The resulting lax extension of the ultrafilter functor is the well-known extension described originally in [2], or in the present form in [3].

4. Another lax extension of the filter functor  $F$  is given by

$$\tilde{F}r(A, B) = \bigvee \{ v \in V \mid \mathfrak{g} \supseteq (r_v^\circ)^b[\mathfrak{f}] \}$$

for any  $\mathbf{V}$ -relation  $r : X \leftrightarrow Y$ ,  $A \in PX$ ,  $B \in PY$ . This extension differs from the one above in that it uses and induces the opposite order on the sets  $FX$ . The corresponding initial extension of the powerset functor (along the principal filter transformation) is

$$\tilde{P}r(A, B) = \bigvee \{ v \in V \mid B \subseteq (r_v^\circ)^b[A] \} .$$

In [19], these extensions were called the “canonical” extensions of their respective functors.

3.7. STRATA EXTENSIONS. Reviewing the lax extensions to  $\mathbf{V}\text{-Rel}$  of Examples 3.6, it becomes clear that they are formed from the corresponding extension to  $\mathbf{Rel}$  according to a certain pattern. These lax extensions are all instances of a construction that allows us to extend any lax functor  $T : \mathbf{Rel} \rightarrow \mathbf{Rel}$  to a lax functor  $T_{\mathbf{V}} : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel}$  as follows. For a set  $X$ , let  $T_{\mathbf{V}}X := TX$  and

$$T_{\mathbf{V}}r(\mathfrak{x}, \mathfrak{y}) := \bigvee \{ v \in \mathbf{V} \mid Tr_v(\mathfrak{x}, \mathfrak{y}) = \top \} ,$$

for any  $\mathbf{V}$ -relation  $r : X \leftrightarrow Y$ . We call  $T_{\mathbf{V}}$  the *strata extension* of  $T$ . This extension process was extensively studied in [3] and [18], to which we refer for further details. For any  $\mathbf{V}$ -relation  $r : X \leftrightarrow Y$  and  $v \in \mathbf{V}$ , we obviously have

$$Tr_v \leq (T_{\mathbf{V}}r)_v , \tag{1}$$

and  $T_{\mathbf{V}}$  is the least extension of a lax functor  $T : \mathbf{Rel} \rightarrow \mathbf{Rel}$  to  $\mathbf{V}\text{-Rel}$  for which this inequality holds. By identifying a  $\mathbf{2}$ -relation  $s : X \leftrightarrow Y$  with its image in  $\mathbf{V}\text{-Rel}$ , we may write  $Ts = Ts_k$  in  $\mathbf{Rel}$ . This equality can be transported to  $\mathbf{V}\text{-Rel}$ , where we then have

$$Ts = Ts_k \leq (T_{\mathbf{V}}s)_k \leq T_{\mathbf{V}}s .$$

Let us point out *en passant* that these inequalities are in fact equalities if  $\mathbf{V}$  is integral, or if  $T$  preserves every empty  $\mathbf{2}$ -relation  $\perp : X \leftrightarrow Y$ . In any case, this shows that if  $\bar{T} : \mathbf{Rel} \rightarrow \mathbf{Rel}$  is a lax extension of a  $\mathbf{Set}$ -functor  $T$ , then  $\bar{T}_{\mathbf{V}}$  is a lax extension of  $T$  to  $\mathbf{V}\text{-Rel}$ .

Finally, if  $\alpha : S \rightarrow T$  is a natural transformation between  $\mathbf{Set}$ -functors, and  $\bar{T}$  is a lax  $\mathbf{Rel}$ -extension of  $T$ , then we have

$$\alpha^*(\bar{T}_{\mathbf{V}}) = (\alpha^*\bar{T})_{\mathbf{V}} .$$

In other words, initial lifts commute with strata extensions.

**3.8. PROPOSITION.** *The preorders induced on the sets  $TX$  by a lax functor  $T : \mathbf{Rel} \rightarrow \mathbf{Rel}$  and its strata extension  $T_{\mathbf{V}}$  coincide.*

**PROOF.** Note that we already have  $(\leq_T) \leq (\leq_{T_{\mathbf{V}}})$  by (1) in 3.7 above. For the other inequality, assume that  $\mathfrak{x} \leq_{T_{\mathbf{V}}} \mathfrak{y}$  holds for  $\mathfrak{x}, \mathfrak{y} \in TX$ , that is,  $k \leq \bigvee \{v \in \mathbf{V} \mid T((1_X)_v)(\mathfrak{x}, \mathfrak{y}) = \top\}$ . Non-triviality of  $\mathbf{V}$  implies that there is some  $u$  in  $\mathbf{V} \setminus \{\perp\}$  with  $T((1_X)_u)(\mathfrak{x}, \mathfrak{y}) = \top$ . Since  $u \neq \perp$ , we have  $(1_X)_u \leq 1_X$ , and therefore  $T1_X(\mathfrak{x}, \mathfrak{y}) = \top$ , so that  $\mathfrak{x} \leq_T \mathfrak{y}$ . ■

**3.9. LAX ALGEBRAS.** Let  $\mathbb{T} = (T, \eta, \mu)$  be a monad on  $\mathbf{Set}$  equipped with a lax extension  $\bar{T}$  of  $T$ . The category  $\mathbf{Alg}(\mathbb{T}, \mathbf{V})$  of  $(\mathbb{T}, \mathbf{V})$ -algebras, or *lax algebras*, has as objects pairs  $(X, a)$ , where  $X$  is a set, and its structure  $a : TX \leftrightarrow X$  is a *reflexive* and *transitive*  $\mathbf{V}$ -relation:

$$1_X \leq a \cdot \eta_X \quad \text{and} \quad a \cdot \bar{T}a \leq a \cdot \mu_X .$$

Morphisms  $f : (X, a) \rightarrow (Y, b)$  are  $\mathbf{Set}$ -maps  $f : X \rightarrow Y$  satisfying:

$$f \cdot a \leq b \cdot Tf ,$$

and composing as in  $\mathbf{Set}$ . It will sometimes be useful to identify the lax extension that is being used; in such cases, we will write  $\mathbf{Alg}(\mathbb{T}, \bar{T}, \mathbf{V})$  instead of  $\mathbf{Alg}(\mathbb{T}, \mathbf{V})$ .

Using the—not necessarily associative—*Kleisli convolution*  $b * a$  of  $\mathbf{V}$ -relations  $a : TX \leftrightarrow Y$ ,  $b : TY \leftrightarrow Z$  defined by

$$b * a := b \cdot \bar{T}a \cdot \mu_X^\circ ,$$

we can rewrite the reflexivity and transitivity conditions above as

$$\eta_X^\circ \leq a \quad \text{and} \quad a * a \leq a . \tag{2}$$

If  $(X, a)$  is a lax algebra, then  $a \cdot (\leq_{\bar{T}}) = a \cdot \bar{T}1_X = a$ , since

$$a = a \cdot 1_{TX} \leq a \cdot (\leq_{\bar{T}}) \leq a \cdot \bar{T}1_X \leq a \cdot \bar{T}a \cdot T\eta_X \leq a \cdot \mu_X \cdot T\eta_X = a$$

by applying  $\bar{T}$  to the reflexivity, and combining with the transitivity of  $a$ . Therefore,  $a$  reverses the order  $\leq_{\bar{T}}$  in its first variable:

$$\mathfrak{x} \leq_{\bar{T}} \mathfrak{y} \implies k \leq \bar{T}1_X(\mathfrak{x}, \mathfrak{y}) \implies a(\mathfrak{y}, z) \leq \bar{T}1_X(\mathfrak{x}, \mathfrak{y}) \otimes a(\mathfrak{y}, z) \leq a \cdot \bar{T}1_X(\mathfrak{x}, z) = a(\mathfrak{x}, z)$$

for all  $\mathfrak{x}, \mathfrak{y} \in TX$ ,  $z \in X$ .

3.10. **EXAMPLES.** Two of the original examples of lax algebras, called *relational algebras* in [2], are given by the categories **Ord** of preordered sets, and **Top** of topological spaces:

$$\mathbf{Ord} \cong \mathbf{Alg}(\mathbb{I}, \mathbf{2}) \quad , \quad \mathbf{Top} \cong \mathbf{Alg}(\mathbb{B}, \mathbf{2}) .$$

Other examples (as well as a proof of these isomorphisms) will be given further on.

3.11. **REMARK.** Since the structure relation  $a : TX \rightarrow X$  of a  $(\mathbb{T}, \mathbf{V})$ -algebra reverses the preorder on  $TX$  in its first variable, it is reasonable to expect that it also preserves a non-trivial preorder in its second. In fact, this depends on the lax extension. To see this, notice first that

$$\eta_X^\circ \cdot \bar{T}a = a \cdot \eta_X \cdot \eta_X^\circ \cdot \bar{T}a \leq a \cdot \bar{T}a \leq a \cdot \mu_X ,$$

so the inequality  $\eta_X^\circ \cdot \bar{T}a \cdot \eta_{TX} \leq a$  always holds. In the case where the unit  $\eta : 1 \rightarrow \bar{T}$  is oplax, this inequality becomes an equality, and  $a$  induces the *dual specialization preorder* on  $X$  via

$$x \leq_a y \iff k \leq a(\eta_X(x), y) .$$

The structure  $a$  is then monotone in its second variable with respect to this preorder:

$$x \leq_a y \implies a(\mathfrak{x}, x) \leq \bar{T}a(\eta_{TX}(\mathfrak{x}), \eta_X(x)) \otimes a(\eta_X(x), y) \leq a(\mathfrak{x}, y)$$

for all  $x, y \in X$ ,  $\mathfrak{x} \in TX$ . Notice also that the Kleisli extension introduced further on does make  $\eta$  into an oplax transformation (Proposition 4.8).

3.12. **FUNCTORIALITY OF  $\mathbf{Alg}(-, \mathbf{V})$ .** We have seen that every monad  $\mathbb{T}$  equipped with a lax extension  $\bar{T} : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel}$  of the underlying functor  $T$  gives rise to a category  $\mathbf{Alg}(\mathbb{T}, \mathbf{V})$ . For a monad  $\mathbb{S}$  equipped with a lax extension  $\bar{S}$ , a *morphism*  $\alpha : (\mathbb{S}, \bar{S}) \rightarrow (\mathbb{T}, \bar{T})$  is a monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  which is also a morphism of lax extensions. In this case,  $\alpha$  induces a 2-functor

$$F_\alpha = \mathbf{Alg}(\alpha, \mathbf{V}) : \mathbf{Alg}(\mathbb{T}, \mathbf{V}) \rightarrow \mathbf{Alg}(\mathbb{S}, \mathbf{V})$$

sending  $(X, a)$  to  $(X, a \cdot \alpha_X)$ , and mapping morphisms identically (see [4], Section 3.7). The correspondence

$$(\alpha : \mathbb{S} \rightarrow \mathbb{T}) \mapsto (F_\alpha : \mathbf{Alg}(\mathbb{T}, \mathbf{V}) \rightarrow \mathbf{Alg}(\mathbb{S}, \mathbf{V}))$$

is functorial, so that  $F_{\alpha \cdot \beta} = F_\beta F_\alpha$  and  $F_1 = \text{Id}$  for all morphisms  $\beta : (\mathbb{R}, \bar{R}) \rightarrow (\mathbb{S}, \bar{S})$  and  $\alpha : (\mathbb{S}, \bar{S}) \rightarrow (\mathbb{T}, \bar{T})$ .

3.13. **INDUCED ADJUNCTIONS.** Let  $S, T$  be two **Set**-functors admitting lax extensions  $\bar{S}, \bar{T}$ , and

$$\alpha \dashv \beta : \bar{T} \rightarrow \bar{S}$$

a *natural adjunction*; that is, a pair  $(\alpha : \bar{S} \rightarrow \bar{T}, \beta : \bar{T} \rightarrow \bar{S})$  of morphisms of lax extensions whose components form an adjunction  $\alpha_X \dashv \beta_X : TX \rightarrow SX$  for every set  $X$ , so that

$$1_X \leq_{\bar{S}} \beta_X \cdot \alpha_X \quad \text{and} \quad \alpha_X \cdot \beta_X \leq_{\bar{T}} 1_X .$$

In the case where  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ ,  $\beta : \mathbb{T} \rightarrow \mathbb{S}$  are also monad morphisms, then there is an induced adjunction

$$F_\alpha \dashv F_\beta : \mathbf{Alg}(\mathbb{S}, \mathbf{V}) \rightarrow \mathbf{Alg}(\mathbb{T}, \mathbf{V}) .$$

Indeed, since  $a$  reverses the order in its first variable, we obtain the inequalities  $a \leq a \cdot (\alpha_X \cdot \beta_X)$  and  $b \cdot (\beta_X \cdot \alpha_X) \leq b$ , for any lax algebra structures  $a : TX \dashv\vdash X$ ,  $b : SX \dashv\vdash X$ . These inequalities yield the unit  $\eta : \text{Id} \rightarrow F_\beta F_\alpha$  and co-unit  $\varepsilon : F_\alpha F_\beta \rightarrow \text{Id}$  of the adjunction.

**3.14. PROPOSITION.** *Let  $\bar{T}$  be a lax extension of  $T$ , and  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  a retraction (so there exists a monad morphism  $\beta : \mathbb{T} \rightarrow \mathbb{S}$  with  $\alpha \cdot \beta \simeq_{\bar{T}} 1$ ). If  $\bar{S}$  is the initial extension induced by  $\alpha$ , then  $\mathbf{Alg}(\mathbb{S}, \mathbf{V})$  and  $\mathbf{Alg}(\mathbb{T}, \mathbf{V})$  are concretely equivalent categories.*

**PROOF.** By composing each side of the equality  $\bar{S}r = \alpha_Y^\circ \cdot \bar{T}r \cdot \alpha_X$  with  $\beta_Y^\circ$  on the left and  $\beta_X$  on the right, we obtain  $\beta_Y^\circ \cdot \bar{S}r \cdot \beta_X = \bar{T}r$ , so that  $\beta$  is a morphism of lax extensions. Moreover, we have

$$1_X \leq \alpha_X^\circ \cdot \bar{T}1_X \cdot \alpha_X = \alpha_X^\circ \cdot \beta_X^\circ \cdot \alpha_X^\circ \cdot \bar{T}1_X \cdot \alpha_X = \alpha_X^\circ \cdot \beta_X^\circ \cdot \bar{S}1_X ,$$

so that  $1 \leq_{\bar{S}} \beta \cdot \alpha$ , and  $\beta \cdot \alpha \leq_{\bar{S}} 1$  by composing each side of  $\alpha_X \cdot \beta_X \simeq 1_X$  with  $\bar{S}1_X$ . Therefore we have  $\beta \cdot \alpha \simeq_{\bar{S}} 1$ , so that  $\alpha \sim \beta$  induces a concrete equivalence  $F_\beta \sim F_\alpha$ . ■

## 4. Kleisli extensions

**4.1. Sup-ENRICHED MONADS.** Let us recall that the Eilenberg-Moore category  $\mathbf{Set}^{\mathbb{P}}$  of the powerset monad is the category  $\mathbf{Sup}$  of sup-semilattices and sup-maps. A *Sup-enriched monad* on  $\mathbf{Set}$  is a pair  $(\mathbb{T}, \tau)$  made up of a monad  $\mathbb{T}$  and a monad morphism  $\tau : \mathbb{P} \rightarrow \mathbb{T}$ . Any such monad morphism induces a concrete functor  $\mathbf{Set}^{\mathbb{T}} \rightarrow \mathbf{Sup}$ . In particular, every set  $TX$  carries the structure of a sup-semilattice, with

$$\bigvee \mathcal{A} := \mu_X \cdot \tau_{TX}(\mathcal{A}) ,$$

for all  $\mathcal{A} \subseteq TX$ ; moreover, the multiplications  $\mu_X$  and each  $Tf : TX \rightarrow TY$  preserve these suprema. Thus, every  $\mathbf{Sup}$ -enriched monad  $(\mathbb{T}, \tau)$  induces a factorization of  $\mathbb{T}$  through  $\mathbf{Sup}$ . Conversely, given a monad  $\mathbb{T} = (T, \eta, \mu)$  factoring through  $\mathbf{Sup}$ , one can define a monad morphism  $\tau : \mathbb{P} \rightarrow \mathbb{T}$  (as in [20]) via

$$\tau_X(A) := \bigvee_{x \in A} \eta_X(x) .$$

Since the operations described above are inverse of each other,  $\mathbf{Sup}$ -enriched monads and monads factoring through  $\mathbf{Sup}$  are equivalent concepts.

A *morphism of Sup-enriched monads*  $\alpha : (\mathbb{S}, \sigma) \rightarrow (\mathbb{T}, \tau)$  is a monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  such that  $\tau = \alpha \cdot \sigma$ . As we show below, this is equivalent to stating that the components of  $\alpha$  are sup-maps.

We say that a **Sup**-enriched monad  $(\mathbb{T}, \tau)$  is *coherent* if the operation  $-^{\mathbb{T}}$  is monotone with respect to the order induced by  $\tau$ ; that is, for any  $g, h : X \rightarrow TY$ , we must have

$$g \leq h \implies g^{\mathbb{T}} \leq h^{\mathbb{T}} .$$

For any **Sup**-enriched monad  $(\mathbb{T}, \tau)$ , coherence is equivalent to the fact that the Kleisli-category  $\mathbf{Set}_{\mathbb{T}}$  is a 2-category with respect to the pointwise order, *i.e.*, the Kleisli-composition  $g \circ f = g^{\mathbb{T}} \cdot f$  is monotone in each variable.

**4.2. PROPOSITION.** *Let  $\mathbb{S} = (S, \delta, \nu)$ ,  $\mathbb{T} = (T, \eta, \mu)$  be monads on  $\mathbf{Set}$ , and  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  a monad morphism. The following statements are equivalent for **Sup**-enrichments  $\sigma$  of  $\mathbb{S}$  and  $\tau$  of  $\mathbb{T}$ :*

- (a)  $\alpha$  is a morphism  $(\mathbb{S}, \sigma) \rightarrow (\mathbb{T}, \tau)$ ;
- (b) each  $\alpha_X : SX \rightarrow TX$  preserves suprema.

**PROOF.** Since  $\mathbf{Sup} \cong \mathbf{Set}^{\mathbb{P}}$ ,  $\alpha_X$  preserves suprema if and only if the diagram

$$\begin{array}{ccc} PSX & \xrightarrow{P\alpha_X} & PTX \\ \nu_X \cdot \sigma_{SX} \downarrow & & \downarrow \mu_X \cdot \tau_{TX} \\ SX & \xrightarrow{\alpha_X} & TX \end{array} \quad (3)$$

commutes. Thus, if  $\alpha : (\mathbb{S}, \sigma) \rightarrow (\mathbb{T}, \tau)$  is a morphism, then

$$\alpha_X \cdot \nu_X \cdot \sigma_{SX} = \mu_X \cdot \alpha_{TX} \cdot \sigma_{TX} \cdot P\alpha_X = \mu_X \cdot \tau_{TX} \cdot P\alpha_X ,$$

as required. Conversely, if the diagram above commutes, we have

$$\begin{aligned} \alpha_X \cdot \sigma_X &= \alpha_X \cdot \nu_X \cdot S\delta_X \cdot \sigma_X = \mu_X \cdot \tau_{TX} \cdot P(\alpha_X \cdot \delta_X) \\ &= \mu_X \cdot \tau_{TX} \cdot P\eta_X = \mu_X \cdot T\eta_X \cdot \tau_X = \tau_X . \end{aligned}$$

■

### 4.3. EXAMPLES.

1. Although there are two trivial monads on  $\mathbf{Set}$ , there is only one which is **Sup**-enriched, namely the monad  $\mathbb{T}_1$  for which  $T_1X = \{*\}$  for all sets  $X$ .
2. The powerset monad  $\mathbb{P}$  is coherent **Sup**-enriched via  $1_{\mathbb{P}}$ , and the supremum operation is given by set union. Moreover,  $(\mathbb{P}, 1_{\mathbb{P}})$  is an initial object in the quasicategory of **Sup**-enriched monads and their morphisms.
3. The filter monad  $\mathbb{F}$  and the up-set monad  $\mathbb{U}$  are coherent **Sup**-enriched via the principal filter natural transformations  $\tau : \mathbb{P} \rightarrow \mathbb{F}$  and  $\tau : \mathbb{P} \rightarrow \mathbb{U}$ , respectively. In both cases, supremum is given by intersection.

4. The monad  $\mathbb{D}_2$  becomes a **Sup**-enriched monad in at least two different ways. Indeed, there are monad morphisms  $\diamond, \square : \mathbb{P} \rightarrow \mathbb{D}_2$ , defined componentwise for  $A \in PX$  by

$$\diamond_X(A) = \{ B \subseteq X \mid A \cap B \neq \emptyset \}, \quad \square_X(A) = \{ B \subseteq X \mid A \subseteq B \},$$

(see [15], Exercise 3.2.18; the notations  $\square$  and  $\diamond$  come from coalgebraic modal logic, where these natural transformations play a vital role). This demonstrates that a factorization through **Sup** is a structure on the monad, and not a property. The suprema induced on  $D_2X$  by  $\square$  are given by intersection, while the ones induced by  $\diamond$  are given by union. However, it can be seen that neither  $(\mathbb{D}_2, \diamond)$  nor  $(\mathbb{D}_2, \square)$  is coherent.

Since the natural transformation  $\tau : \mathbb{P} \rightarrow \mathbb{U}$  above is just the restriction of  $\square$ , the embeddings

$$(\mathbb{P}, 1_{\mathbb{P}}) \rightarrow (\mathbb{F}, \tau) \rightarrow (\mathbb{U}, \tau) \rightarrow (\mathbb{D}_2, \square)$$

form a chain of **Sup**-enriched monads (although only the first three are coherent).

- 4.4. **THE KLEISLI EXTENSION.** Let  $(\mathbb{T}, \tau)$  be a **Sup**-enriched monad. If  $r : X \leftrightarrow Y$  is a relation, we denote by  $r^\tau : TY \rightarrow TX$  the composite

$$r^\tau := (\tau_X \cdot r^b)^\mathbb{T} = \mu_X \cdot T(\tau_X \cdot r^b).$$

For a relation  $s : Y \leftrightarrow Z$  and a map  $g$ , we obtain the following useful formulas:

$$r^\tau \cdot \eta_X = \tau_X \cdot r^b, \quad (s \cdot r)^\tau = r^\tau \cdot s^\tau, \quad (g^\circ)^\tau = Tg. \quad (4)$$

The first equation is obvious; the other two follow from the fact that  $\tau$  is a monad morphism.

The *Kleisli extension*  $T^\tau : \mathbf{Rel} \rightarrow \mathbf{Rel}$  of  $T$  with respect to  $\tau$  is defined by

$$T^\tau r(\mathfrak{x}, \mathfrak{y}) = \top \iff \mathfrak{x} \leq r^\tau(\mathfrak{y}).$$

for a relation  $r : X \leftrightarrow Y$  and  $\mathfrak{x} \in TX$ ,  $\mathfrak{y} \in TY$ . Using strata extensions, we can define  $T_{\mathbf{V}}^\tau := (T^\tau)_{\mathbf{V}} : \mathbf{V}\text{-Rel} \rightarrow \mathbf{V}\text{-Rel}$  as

$$T_{\mathbf{V}}^\tau r(\mathfrak{x}, \mathfrak{y}) = \bigvee \{ v \in \mathbf{V} \mid \mathfrak{x} \leq (r_v)^\tau(\mathfrak{y}) \},$$

for a **V**-relation  $r : X \leftrightarrow Y$  and  $\mathfrak{x} \in TX$ ,  $\mathfrak{y} \in TY$ . This formula was used in [20] to define  $T_{\mathbf{V}}^\tau$  in one step. There, it is shown that *if  $\tau$  is a coherent **Sup**-enrichment*, then  $T_{\mathbf{V}}^\tau$  is a lax extension of  $T$ , and that whenever **V** is non-trivial (as we assumed in this work), then the preorder  $\leq_{T_{\mathbf{V}}^\tau}$  associated with this extension is identical to the original order on  $TX$ . In particular, all  $\mu_X$  and  $Tf$  preserve this order.

- 4.5. **REMARK.** The previous definition of the Kleisli extension does not require that  $(\mathbb{T}, \tau)$  be *coherent **Sup**-enriched*, so that  $T_{\mathbf{V}}^\tau$  is not necessarily a lax extension of  $T$ . However, one can still exploit this definition to put forth meaningful mechanisms that underly the coherent case (see in particular Example 4.6(3) below, or Corollary 4.13).

## 4.6. EXAMPLES.

1. For the trivial **Sup**-enriched monad  $\mathbb{T}_1$ , we have  $\tau_X = !_X : PX \rightarrow \{*\}$  and  $T_{\mathbf{V}}^!r(*, *) = \top$  for any **V**-relation  $r : X \leftrightarrow Y$ . In particular,  $(T_1, T_{\mathbf{V}}^!)$  is a final object in the quasicategory of lax extensions.
2. The Kleisli extensions of the powerset, filter and up-set functors are the lax extensions described in 3.6.
3. Example 4.3(4) shows that there are at least *two* lax extensions obtained as Kleisli extensions of the up-set functor  $U$ . Indeed, the monad morphism  $\square : \mathbb{P} \rightarrow \mathbb{D}_2$  restricts to the principal filter natural transformation  $\tau : \mathbb{P} \rightarrow \mathbb{U}$ , and  $\diamond : \mathbb{P} \rightarrow \mathbb{D}_2$  also restricts to a *coherent* **Sup**-enrichment  $\sigma : \mathbb{P} \rightarrow \mathbb{U}$ . This demonstrates that the lax extensions presented here are fundamentally different from the extensions obtained by the original construction of Barr in [2]. Indeed, the latter are lax functors if and only if the original **Set**-functor satisfies the Beck-Chevalley condition of ([3], Section 1.3). However, it can be seen that the up-set functor  $U$  does not satisfy this condition, so Barr's construction fails to yield a lax extension of  $U$ .

4.7. PROPOSITION. *The Kleisli extension  $T^\tau : \text{Rel} \rightarrow \text{Rel}$  preserves composition:*

$$T^\tau s \cdot T^\tau r = T^\tau (s \cdot r)$$

for all relations  $r : X \leftrightarrow Y$ ,  $s : Y \leftrightarrow Z$ .

PROOF. This follows immediately from the second equation of (4) in 4.4. ■

4.8. PROPOSITION. *If  $(\mathbb{T}, \tau)$  is a coherent **Sup**-enriched monad, then  $\eta : (\text{Id}, \text{Id}) \rightarrow (T, T_{\mathbf{V}}^\tau)$  is a morphism of lax extensions, that is:  $\eta : 1 \rightarrow \overline{T}$  is an oplax transformation.*

PROOF. Take a **V**-relation  $r : X \leftrightarrow Y$ , and for  $x \in X$ ,  $y \in Y$  set  $v := r(x, y)$ . By definition,  $\{x\} \subseteq r_v^b(y)$ , so that  $\eta_X(x) = \tau_X(\{x\}) \leq \tau_X \cdot r_v^b(y) = (\tau_X \cdot r_v^b)^\top \cdot \eta_Y(y)$ . We conclude that  $r(x, y) = v \leq T_{\mathbf{V}}^\tau r(\eta_X(x), \eta_Y(y))$ , so  $r \leq \eta_Y^\circ \cdot T_{\mathbf{V}}^\tau r \cdot \eta_X$ , as required. ■

4.9. PROPOSITION. *If  $\alpha : (\mathbb{S}, \sigma) \rightarrow (\mathbb{T}, \tau)$  is a morphism of coherent **Sup**-enriched monads, then  $\alpha : S^\sigma \rightarrow T^\tau$  is a morphism of lax extensions.*

Moreover, if the  $\alpha_X$  are order-embeddings, then we have  $S^\sigma = \alpha^*(T^\tau)$ ; that is, the initial extension induced by  $\alpha$  is the Kleisli extension of  $S$ .

PROOF. Observe that we have  $\alpha_X \cdot (r_v)^\sigma = (r_v)^\tau \cdot \alpha_Y$  for any **V**-relation  $r : X \leftrightarrow Y$  and  $v \in \mathbf{V}$ . Therefore,

$$\mathfrak{x} \leq (r_v)^\sigma(\mathfrak{y}) \implies \alpha_X(\mathfrak{x}) \leq \alpha_X \cdot (r_v)^\sigma(\mathfrak{y}) = (r_v)^\tau \cdot \alpha_Y(\mathfrak{y})$$

for all  $\mathfrak{x} \in SX$ ,  $\mathfrak{y} \in SY$ . This implies immediately that  $\alpha$  is a morphism between the Kleisli extensions. If moreover  $\alpha_X$  is an order-embedding, the implication above is an equivalence, so that  $\alpha$  is initial. ■



4.10. **OP-CANONICAL EXTENSIONS OF TAUT MONADS.** For any taut monad  $\mathbb{S}$  (see [16]), the component at a set  $X$  of the *support* monad morphism  $\text{supp} : \mathbb{S} \rightarrow \mathbb{F}$  maps  $\mathfrak{x} \in SX$  to the filter

$$\text{supp}_X(\mathfrak{x}) := \{ A \subseteq X \mid \mathfrak{x} \in SA \} ,$$

where  $SA$  is identified with a subset of  $SX$ . One obtains a lax extension  $\bar{S} = \text{supp}^*(F^\tau)$ , so that

$$\bar{S}r(\mathfrak{x}, \mathfrak{y}) := \bigvee \{ v \in \mathbf{V} \mid \forall B \subseteq Y \ (B \in \text{supp}_Y(\mathfrak{y}) \implies r_v^b[B] \in \text{supp}_X(\mathfrak{x})) \} ,$$

for all  $r : X \leftrightarrow Y$ ,  $\mathfrak{x} \in SX$ , and  $\mathfrak{y} \in SY$ . This lax extension is just the “op-canonical” extension of a taut functor introduced in [19]. The natural inclusion  $\varepsilon : \mathbb{F} \rightarrow \mathbb{U}$  is an embedding of coherent **Sup**-enriched monads  $(\mathbb{F}, \tau) \rightarrow (\mathbb{U}, \tau)$ , and we have  $F^\tau = \varepsilon^*(U^\tau)$  by Proposition 4.9. Thus, the op-canonical extension of  $\mathbb{S}$  arises in fact from the **Sup**-enrichment  $\tau$  of  $\mathbb{U}$ :

$$\bar{S} = \text{supp}^*(F^\tau) = \text{supp}^*(\varepsilon^*(U^\tau)) = (\varepsilon \cdot \text{supp})^*(U^\tau) .$$

Note that Proposition 3.14 states that the functor  $F_{\text{supp}} : \text{Alg}(\mathbb{F}, \mathbf{V}) \rightarrow \text{Alg}(\mathbb{S}, \mathbf{V})$  induced by the support morphism does not yield new examples of lax algebras if the monad  $\mathbb{S}$  “contains”  $\mathbb{F}$ , that is, if  $\text{supp} : \mathbb{S} \rightarrow \mathbb{F}$  is a retraction.

4.11. **CANONICAL EXTENSIONS OF TAUT MONADS.** The “canonical” extension [19] of a taut monad  $\mathbb{S}$  is described by

$$\tilde{S}r := (\bar{S}(r^\circ))^\circ ,$$

for all  $r : X \leftrightarrow Y$ , where  $\bar{S}$  is the “op-canonical” extension given above. This canonical extension arises as an initial lift of  $U^\sigma$  along  $\varepsilon \cdot \text{supp}$ , where  $\sigma : \mathbb{P} \rightarrow \mathbb{U}$  is the codomain-restriction of the natural transformation  $\diamond : \mathbb{P} \rightarrow \mathbb{D}_2$  (see Example 4.6 (3)). To verify our claim, it sufficient to see that

$$D_2^\diamond(r) = (D_2^\square(r^\circ))^\circ ,$$

since one obtains  $U^\sigma(r) = (U^\tau(r^\circ))^\circ$  by restriction of the double-dualization monad to the up-set one (see Corollary 4.13 below).

Thus, let us point out first that every Kleisli morphism  $f : Y \rightarrow D_2X$  of  $\mathbb{D}_2$  corresponds uniquely to a map  $f_\bullet : PX \rightarrow PY$  via

$$f_\bullet(A) = \{ y \in Y \mid A \in f(y) \} , \quad f(y) = \{ A \subseteq X \mid y \in f_\bullet(A) \} .$$

This operation  $-_\bullet$  is related to the monad operations via  $f^{\mathbb{D}_2} = f_\bullet^{-1}$  and  $(\dot{-})_\bullet = 1_P$ . It is also functorial, since

$$(f \circ g)_\bullet = (f^{\mathbb{D}_2} \cdot g)_\bullet = g_\bullet \cdot f_\bullet$$

holds for all  $g : Z \rightarrow D_2Y$ , and we have  $f \subseteq f'$  pointwise if and only if  $f_\bullet \subseteq f'_\bullet$  pointwise.

4.12. **PROPOSITION.** *For any relation  $r$ , we have  $D_2^\diamond(r) = (D_2^\square(r^\circ))^\circ$ .*

PROOF. Let us write  $[r] := (\square_X \cdot r^b)_\bullet : PX \rightarrow PY$  and  $\langle r \rangle := (\diamond_X \cdot r^b)_\bullet : PX \rightarrow PY$  for a relation  $r : X \leftrightarrow Y$ . We remark that

$$[r](A) = \{y \mid \forall x : r(x, y) = \top \Rightarrow x \in A\} \quad \text{and} \quad \langle r \rangle(A) = \{y \mid \exists x \in A : r(x, y) = \top\}$$

hold; that is,  $[r]$  and  $\langle r \rangle$  are just the “necessarily” and “possibly”, respectively, modalities associated to the relation  $r^\circ$ . In any case, we have an adjunction

$$\langle r^\circ \rangle \dashv [r] ,$$

and  $r^\square = (\square_X \cdot r^b)^{\mathbb{D}_2} = [r]^{-1}$  is left adjoint to  $\langle r^\circ \rangle^{-1} = (\diamond_X \cdot (r^\circ)^b)^{\mathbb{D}_2} = (r^\circ)^\diamond$ . Thus, recalling the orders induced by  $\square$  and  $\diamond$ , we have, for  $\mathfrak{x} \in D_2X$  and  $\mathfrak{y} \in D_2Y$ :

$$\mathfrak{x} \leq r^\square(\mathfrak{y}) \iff r^\square(\mathfrak{y}) \subseteq \mathfrak{x} \iff \mathfrak{y} \subseteq (r^\circ)^\diamond(\mathfrak{x}) \iff D_2^\diamond(r^\circ)(\mathfrak{y}, \mathfrak{x}) = \top ,$$

so that  $D_2^\square(r) = (D_2^\diamond(r^\circ))^\circ$ , or equivalently  $D_2^\diamond(r) = (D_2^\square(r^\circ))^\circ$ . ■

4.13. COROLLARY. *For any relation  $r$ , we have  $U^\sigma(r) = (U^\tau(r^\circ))^\circ$ .*

PROOF. This is a direct consequence of the previous Proposition, since  $\sigma : \mathbb{P} \rightarrow \mathbb{U}$  is the codomain-restriction of  $\diamond : \mathbb{P} \rightarrow \mathbb{D}_2$ , and the principal filter natural transformation  $\tau : \mathbb{P} \rightarrow \mathbb{U}$  is the codomain-restriction of  $\square : \mathbb{P} \rightarrow \mathbb{D}_2$ . ■

4.14. REMARK. For future use, notice that a map  $f : Y \rightarrow D_2X$  factorizes through  $UX \hookrightarrow D_2X$  if and only if  $f_\bullet$  is monotone, and through  $FX \hookrightarrow D_2X$  if and only if  $f_\bullet$  preserves finite infima.

4.15. INITIAL LIFTS OF KLEISLI EXTENSIONS. Fix a monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ , where  $\mathbb{T} = (T, \eta, \mu)$  is coherent **Sup**-enriched via  $\tau$ , and  $\mathbb{S} = (S, \delta, \nu)$ . We will now study extensions of the form  $\alpha^*(T_V^\tau)$ . Since initial lifts commute with strata extensions, it actually suffices to study  $\mathbf{V} = \mathbf{2}$ , and from now on we write  $\bar{S} := \alpha^*(T^\tau)$ . In this context, each  $\nu_X$  is monotone: in the commutative diagram

$$\begin{array}{ccccc} S S X & \xrightarrow{\alpha_{S X}} & T S X & \xrightarrow{T \alpha_X} & T T X \\ \nu_X \downarrow & & & & \downarrow \mu_X \\ S X & \xrightarrow{\alpha_X} & T X & & \end{array}$$

the upper-right path is monotone, so  $\nu_X$  is monotone by initiality of  $\alpha_X$ . Observe that  $\delta$  and  $\nu$  become oplax transformations with respect to  $\bar{S}$ .

We introduce a natural transformation  $\alpha^\vee : PS \rightarrow T$  via

$$\alpha_X^\vee := \bigvee P \alpha_X = \mu_X \cdot \tau_{T X} \cdot P \alpha_X = \alpha_X^\mathbb{T} \cdot \tau_{S X} ,$$

or equivalently,  $\alpha_X^\vee(\mathcal{A}) = \bigvee \alpha_X[\mathcal{A}]$  for all  $\mathcal{A} \subseteq S X$ . Each  $\alpha_X^\vee$  preserves suprema, and therefore has a right adjoint which we denote by  $\alpha_X^\downarrow : T X \rightarrow P S X$ . Clearly,

$$\alpha_X^\downarrow(\mathfrak{f}) = \{\mathfrak{x} \in S X \mid \alpha_X(\mathfrak{x}) \leq \mathfrak{f}\} .$$

The maps  $\alpha_X^\downarrow$  allow for a convenient description of  $\bar{S} = \alpha^*(T^\tau)$ . Indeed, for  $r : X \leftrightarrow Y$  we have

$$(\bar{S}r)^b = \alpha_X^\downarrow \cdot r^\tau \cdot \alpha_Y.$$

In particular, the order relation  $\leq_{\bar{S}}$  on  $SX$  satisfies  $(\leq_{\bar{S}})^b = \alpha_X^\downarrow \cdot \alpha_X$ .

4.16. **SUP-GENERATING MORPHISMS.** We say that  $\mathfrak{r} \in TX$  is  $\alpha$ -*approachable* if it is in the image of  $\alpha_X^\vee$ . We call  $\alpha$  *sup-generating* in  $(\mathbb{T}, \tau)$  if each  $\alpha_X^\vee$  is surjective, that is, if

$$\alpha^\vee \cdot \alpha^\downarrow = 1,$$

or equivalently,

$$\forall \mathfrak{f} \in TX \exists \mathcal{A} \subseteq SX : \mathfrak{f} = \bigvee \alpha_X[\mathcal{A}],$$

holds. When  $\mathbb{S}$  is a submonad of  $\mathbb{T}$  and the embedding is sup-generating, we simply say that  $\mathbb{S}$  is *sup-generating* in  $\mathbb{T}$ .

Observe that a morphism of **Sup**-enriched monads is sup-generating if and only if every  $\alpha_X$  is surjective. Indeed, for  $\alpha : (\mathbb{S}, \sigma) \rightarrow (\mathbb{T}, \tau)$ ,  $\alpha_X^\vee$  is just the diagonal in (3) of 4.2, and  $\nu_X \cdot \sigma_{SX}$  is always surjective.

4.17. **THE FIRST INTERPOLATION CONDITION.** We say that a morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  *interpolates* a relation  $r : SX \leftrightarrow Y$  (in the **Sup**-enriched monad  $(\mathbb{T}, \tau)$ ) if

$$\alpha_X^\downarrow \cdot \alpha_X^\vee \cdot r^b \leq (\leq_{\bar{S}X}^b \cdot \nu_X)^\mathbb{P} \cdot (\bar{S}r)^b \cdot \delta_Y$$

holds (where  $\bar{S} = \alpha^*(T^\tau)$ ). The above condition expands to  $\alpha_X^\downarrow \cdot \alpha_X^\vee \cdot r^b \leq (\alpha_X^\downarrow \cdot \alpha_X \cdot \nu_X)^\mathbb{P} \cdot \alpha_{SX}^\downarrow \cdot r^\tau \cdot \eta_Y$  and can be written pointwise as

$$\alpha_X(\mathfrak{r}) \leq \bigvee \{ \alpha_X(\mathfrak{h}) \mid r(\mathfrak{h}, y) = \top \} \implies \exists \mathfrak{x} \in SSX : \mathfrak{r} \leq \nu_X(\mathfrak{x}) \ \& \ \alpha_{SX}(\mathfrak{x}) \leq r^\tau \cdot \eta_Y(y)$$

for all  $\mathfrak{r} \in SX$ ,  $y \in Y$ ; note that  $\alpha_{SX}(\mathfrak{x}) \leq r^\tau \cdot \eta_Y(y)$  is equivalent to  $\bar{S}r(\mathfrak{x}, \delta_Y(y)) = \top$ . If  $\mathbb{S}$  is a submonad of  $\mathbb{T}$ , the previous condition naturally has a simpler expression, and may be represented graphically by

$$\mathfrak{r} \leq \mu_X \cdot r^\tau \cdot \eta_Y(y) \implies \exists \mathfrak{x} : \begin{array}{c} \mathfrak{x} \leq r^\tau \cdot \eta_Y(y) \\ \downarrow \\ \mathfrak{r} \leq \mu_X(\mathfrak{x}) \end{array}$$

The morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  satisfies the *first interpolation condition* in  $(\mathbb{T}, \tau)$  if it interpolates every relation  $r : SX \leftrightarrow Y$ .

4.18. **THE SECOND INTERPOLATION CONDITION.** A morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  satisfies the *second interpolation condition* (in the **Sup**-enriched monad  $(\mathbb{T}, \tau)$ ) if the following inequality

$$\alpha_X^{\mathbb{T}} \cdot r^\tau \cdot \alpha_Y \leq \alpha_X^\vee \cdot P\nu_X \cdot (\overline{S}r)^{\flat}$$

holds for all relations  $r : SX \leftrightarrow Y$  (where  $\overline{S} = \alpha^*(T^\tau)$ ). In pointwise notation, this condition becomes

$$\mu_X \cdot T\alpha_X \cdot r^\tau \cdot \alpha_Y(\eta) \leq \bigvee \{ \alpha_X \cdot \nu_X(\mathfrak{X}) \mid \mathfrak{X} \in SSX : \alpha_{SX}(\mathfrak{X}) \leq r^\tau \cdot \alpha_Y(\eta) \} ,$$

for all  $\eta \in SY$ . Note that this can be read as an equality, since

$$\alpha_X^\vee \cdot P\nu_X \cdot \alpha_{SX}^\downarrow \cdot r^\tau \cdot \alpha_Y \leq \alpha_X^{\mathbb{T}} \cdot r^\tau \cdot \alpha_Y$$

always holds. Indeed, we have

$$\alpha_X^\vee \cdot P\nu_X = (\alpha_X \cdot 1_{SX}^{\mathbb{S}})^{\mathbb{T}} \cdot \tau_{SSX} = (\alpha_X^{\mathbb{T}} \cdot \alpha_{SX})^{\mathbb{T}} \cdot \tau_{SSX} = \alpha_X^{\mathbb{T}} \cdot \alpha_{SX}^\vee \quad (5)$$

so that  $\alpha_X^\vee \cdot P\nu_X \cdot \alpha_{SX}^\downarrow \leq \alpha_X^{\mathbb{T}}$ , and we can conclude by composing each side with  $r^\tau \cdot \alpha_Y$  on the right.

Let us point out that if  $\alpha$  is sup-generating, then it satisfies the second interpolation condition. This is immediate from (5) upon composing with  $\alpha_{SX}^\downarrow$ .

4.19. **INTERPOLATING MORPHISMS.** A monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  is *interpolating* in  $(\mathbb{T}, \tau)$  if it satisfies the *first and second* interpolation conditions. If  $\mathbb{S}$  is a submonad of  $\mathbb{T}$  and the embedding is interpolating, we may simply say that  $\mathbb{S}$  is *interpolating* in  $\mathbb{T}$ .

Notice that  $\alpha$  is interpolating whenever it is a morphism of coherent **Sup**-enriched monads  $\alpha : (\mathbb{S}, \sigma) \rightarrow (\mathbb{T}, \tau)$ . Indeed, if  $\mathfrak{X} := r^\sigma(\eta) \in SSX$ , then we get

$$\alpha_X \cdot \nu_X(\mathfrak{X}) = \mu_X \cdot T\alpha_X \cdot r^\tau \cdot \alpha_Y(\eta) \quad \text{and} \quad \alpha_{SX}(\mathfrak{X}) = r^\tau \cdot \alpha_Y(\eta) .$$

Therefore, the second interpolation condition is verified, and the first follows by setting  $\eta := \delta_Y(y)$ .

4.20. **EXAMPLES.**

1. Any **Sup**-enriched monad  $\mathbb{T} = (T, \eta, \mu)$  comes with a monad morphism  $\eta : \mathbb{I} \rightarrow \mathbb{T}$ . By the fact that  $\tau_X = \bigvee P\eta_X$  (see 4.1), both interpolation conditions may easily be seen to hold. That is,  $\eta$  is always interpolating.
2. If  $\mathbb{S} = \mathbb{P}$  is the powerset monad embedded in  $\mathbb{T} = \mathbb{F}$  via the principal filter morphism  $\tau : \mathbb{P} \rightarrow \mathbb{F}$ , then the interpolation conditions are immediate since  $\mathbb{P}$  is **Sup**-enriched.
3. In the case where  $\mathbb{T} = \mathbb{F}$  is the filter monad, and  $\mathbb{S} = \mathbb{B}$  the ultrafilter monad, we simply have for  $\mathfrak{f} \in FY$  and  $r : X \leftrightarrow Y$  that  $r^\tau(\mathfrak{f})$  is the filter on  $X$  given by

$$r^\tau(\mathfrak{f}) = r^{\flat}[\mathfrak{f}] = \uparrow \{ r^{\flat}[B] \mid B \in \mathfrak{f} \} .$$

To see that  $\mathbb{B}$  is interpolating in  $\mathbb{F}$ , it is sufficient to verify the first interpolation condition, since  $\mathbb{B}$  is sup-generating in  $\mathbb{F}$ . For ultrafilters  $\mathfrak{x}, \mathfrak{y}$  on  $X$  and a relation  $a : \beta X \leftrightarrow X$ , the inequality  $\mathfrak{x} \leq \Sigma_X \cdot a^\tau(\mathfrak{y})$  translates as

$$\forall B \in \mathfrak{y} (a^b[B] \subseteq A^{\mathbb{B}} \implies A \in \mathfrak{x}) ,$$

for all  $A \subseteq X$  (where  $A^{\mathbb{B}} = \{\mathfrak{z} \in \beta X \mid A \in \mathfrak{z}\}$ ). If there existed  $A \in \mathfrak{x}$  and  $B \in \mathfrak{y}$  with  $A^{\mathbb{B}} \cap a^b[B] = \emptyset$ , we would have  $a^b[B] \subseteq (A^{\mathbb{B}})^c = (A^c)^{\mathbb{B}}$  (here  $A^c$  denotes the set-complement of  $A$ ), so that  $A^c \in \mathfrak{x}$ , a contradiction. Therefore,  $A^{\mathbb{B}} \cap a^b[B] \neq \emptyset$  for all  $A \in \mathfrak{x}$  and  $B \in \mathfrak{y}$ , and there exists an ultrafilter  $\mathfrak{X}$  on  $\beta X$  that refines both  $\{A^{\mathbb{B}} \mid A \in \mathfrak{x}\}$  and  $a^\tau(\mathfrak{y})$ . This implies that  $\Sigma_X(\mathfrak{X}) = \mathfrak{x}$ , and we can conclude by setting  $\mathfrak{y} = \eta_X(y)$ .

## 5. Kleisli structures and lax algebras

5.1. **KLEISLI MONOIDS.** Let  $(\mathbb{T}, \tau)$  be a coherent **Sup**-enriched monad. The category  $\mathbf{KIMon}(\mathbb{T})$  of *Kleisli monoids* has as objects pairs  $(X, c)$ , where  $X$  is a set, and its structure  $c : X \rightarrow TX$  is an *extensive* and *idempotent* map:

$$\eta_X \leq c \quad \text{and} \quad c \circ c \leq c .$$

Morphisms  $f : (X, c) \rightarrow (Y, d)$  are **Set**-maps  $f : X \rightarrow Y$  satisfying:

$$Tf \cdot c \leq d \cdot f .$$

Contrasting the conditions on the structure on a Kleisli monoid with the reflexivity and transitivity conditions of a lax algebra as in (2) of 3.9, we note the strong parallel between the two concepts. This parallel will be analyzed in more detail in Theorem 5.6 and its subsequent results.

### 5.2. EXAMPLES.

1. If one takes for  $\mathbb{T}$  the powerset monad  $\mathbb{P}$ , then  $\mathbf{KIMon}(\mathbb{P})$  is the category of reflexive and transitive relations. That is,  $\mathbf{KIMon}(\mathbb{P})$  is concretely isomorphic to the category **Ord** of preordered sets.
2. Let  $\mathbb{T}$  be the up-set monad  $\mathbb{U}$ . Remark 4.14 tells us that a map  $i : X \rightarrow UX$  corresponds to a monotone map  $i_\bullet : PX \rightarrow PX$ . The discussion in Section 4.11 then shows that  $i$  is extensive and idempotent if and only if  $i_\bullet$  satisfies

$$A \supseteq i_\bullet(A) \quad \text{and} \quad i_\bullet \cdot i_\bullet(A) \supseteq i_\bullet(A)$$

for all  $A \in PX$ , that is, if and only if  $i_\bullet$  is an interior operator on  $X$ . One easily checks that  $\mathbf{KIMon}(\mathbb{U})$ -morphisms correspond to continuous maps  $f : (X, i_\bullet) \rightarrow (Y, j_\bullet)$ , that is, maps  $f : X \rightarrow Y$  such that

$$f^{-1}(j_\bullet(A)) \subseteq i_\bullet(f^{-1}(A))$$

for all  $A \subseteq Y$ . Hence,  $\mathbf{KIMon}(\mathbb{U})$  is concretely isomorphic to the category  $\mathbf{Int}$  of interior spaces, and therefore also concretely isomorphic to the category  $\mathbf{Cls}$  of closure spaces.

3. Let us consider now the filter monad  $\mathbb{F}$ . From Remark 4.14 and the previous example, it follows that  $\mathbf{KIMon}(\mathbb{F})$ -structures on a set  $X$  correspond to interior operators which preserve finite infima, that is, to topologies on  $X$ . Thus,  $\mathbf{KIMon}(\mathbb{F})$  is concretely isomorphic to the category  $\mathbf{Top}$  of topological spaces. The “monadic” presentation of topological spaces as Kleisli monoids is exactly Hausdorff’s original definition [9] of topological spaces by way of neighborhood systems, rephrased in categorical terms (and without the Hausdorff separation condition). See [12] for applications and further references.
4. Since there is a monad morphism  $\tau : \mathbb{P} \rightarrow \mathbb{T}$ , any Eilenberg-Moore algebra  $(X, a : TX \rightarrow X)$  is a sup-semilattice  $X$  with supremum given by  $a \cdot \tau_X$  (as in 4.1). Moreover, we have

$$a \cdot (\mu_X \cdot \tau_{TX}) = a \cdot Ta \cdot \tau_{TX} = (a \cdot \tau_X) \cdot Pa ,$$

so  $a$  is a sup-map and therefore has a right adjoint  $a^* : X \rightarrow TX$ . On one hand,  $a \cdot \eta_X \leq 1_X$  implies  $\eta_X \leq a^*$ . On the other,  $a \cdot \mu_X \leq a \cdot Ta$  yields  $\mu_X \leq a^* \cdot a \cdot Ta$ , so

$$a^* \circ a^* = \mu_X \cdot Ta^* \cdot a^* \leq a^* \cdot a \cdot T(a \cdot a^*) \cdot a^* \leq a^* .$$

Therefore,  $(X, a^*)$  is a Kleisli  $\mathbb{T}$ -algebra, and the Eilenberg-Moore category  $\mathbf{Set}^{\mathbb{T}}$  is a subcategory of  $\mathbf{KIMon}(\mathbb{T})$ . More precisely, there is a faithful functor  $\mathbf{Set}^{\mathbb{T}} \rightarrow \mathbf{KIMon}(\mathbb{T})$  (not necessarily full, as the examples below demonstrate) that is injective on objects.

This example presents the category  $\mathbf{Sup} \cong \mathbf{Set}^{\mathbb{P}}$  of sup-semilattices as a subcategory of  $\mathbf{Ord}$ , and the category  $\mathbf{Cnt} \cong \mathbf{Set}^{\mathbb{F}}$  of continuous lattices [6] as a subcategory of  $\mathbf{Top}$ . Moreover, the topology obtained on continuous lattices via the previous considerations is the Scott topology [8].

**5.3. SUPERCATEGORIES OF THE CATEGORY OF KLEISLI MONOIDS.** Let us introduce the following supercategories of  $\mathbf{KIMon}(\mathbb{T})$ : the objects of the category  $\mathbf{K}(\mathbb{T})$  are pairs  $(X, c : X \rightarrow TX)$ , and the morphisms  $f : (X, c) \rightarrow (Y, d)$  are maps  $f : X \rightarrow Y$  such that  $Tf \cdot c \leq d \cdot f$ . We also introduce the following full subcategories of  $\mathbf{K}(\mathbb{T}) = \mathbf{KIMon}_0(\mathbb{T})$ :

- $\mathbf{KIMon}_1(\mathbb{T})$  is the full subcategory of  $\mathbf{KIMon}_0(\mathbb{T})$  whose objects have an extensive structure;
- $\mathbf{KIMon}_2(\mathbb{T})$  is the full subcategory of  $\mathbf{KIMon}_0(\mathbb{T})$  whose objects have an idempotent structure.

Giving a concrete description of the various categories of the form  $\mathbf{KIMon}_i(\mathbb{T})$  for the monads mentioned in Examples 5.2 is straightforward. For instance,  $\mathbf{KIMon}_1(\mathbb{F})$  is concretely isomorphic to the category of pretopological spaces.

5.4. **SUPERCATEGORIES OF LAX ALGEBRAS.** Let  $\mathbb{T} = (T, \eta, \mu)$  be a monad on  $\mathbf{Set}$  together with a lax extension  $\bar{T}$  of  $T$ . We say a  $\mathbf{V}$ -relation  $a : TX \leftrightarrow Y$  is

- *antitone* if  $a \cdot (\leq_{\bar{T}}) \leq a$  holds, and
- *left unitary* if  $\eta_Y^\circ \cdot \bar{T}a \leq a \cdot \mu_X$  holds.

In the notations of 3.9, the left unitary condition may be rephrased as  $\eta_Y^\circ * a \leq a$ . Observe also that the structure  $\mathbf{V}$ -relation of a lax algebra  $(X, a)$  is antitone as well as left unitary.

Denote by  $\mathbf{A}(\mathbb{T}, \mathbf{V})$  the category whose objects are pairs  $(X, a)$  (with  $a : TX \leftrightarrow X$  a  $\mathbf{V}$ -relation), and whose morphisms  $f : (X, a) \rightarrow (Y, b)$  are the maps  $f : X \rightarrow Y$  satisfying  $f \cdot a \leq b \cdot Tf$ . We define the following full subcategories of  $\mathbf{A}(\mathbb{T}, \mathbf{V})$ :

- the objects of  $\mathbf{Alg}_0(\mathbb{T}, \mathbf{V})$  are those pairs  $(X, a)$  whose structure  $a$  is antitone and left unitary;
- $\mathbf{Alg}_1(\mathbb{T}, \mathbf{V})$  is the full subcategory of  $\mathbf{Alg}_0(\mathbb{T}, \mathbf{V})$  whose objects have a reflexive structure;
- $\mathbf{Alg}_2(\mathbb{T}, \mathbf{V})$  is the full subcategory of  $\mathbf{Alg}_0(\mathbb{T}, \mathbf{V})$  whose objects have a transitive structure.

5.5. **RELATING LAX ALGEBRAS AND KLEISLI MONOIDS.** Guided by the example of topological spaces, which can be described not only via neighborhood filters or via filter convergence, but also by the convergence of ultrafilters, we are going to compare  $\mathbf{KIMon}(\mathbb{T})$  to  $\mathbf{Alg}(\mathbb{S}, \mathbf{2})$ , where the monad  $\mathbb{S}$  is related to the coherent  $\mathbf{Sup}$ -enriched monad  $(\mathbb{T}, \tau)$  by way of a monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ . The following Theorem will follow directly from Proposition 5.13:

5.6. **THEOREM.** *Let  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  be a monad morphism with  $\mathbb{T}$  coherent  $\mathbf{Sup}$ -enriched via  $\tau$ , and set  $\bar{S} = \alpha^*(T^\tau)$ . If  $\alpha$  is sup-generating and interpolating then the dotted arrows in the following diagram are concrete isomorphisms:*

$$\begin{array}{ccccc}
 \mathbf{KIMon}(\mathbb{T}) & \hookrightarrow & \mathbf{KIMon}_1(\mathbb{T}) & & \\
 \downarrow & \searrow \cong & \downarrow & \searrow \cong & \\
 & \mathbf{Alg}(\mathbb{S}, \mathbf{2}) & \hookrightarrow & \mathbf{Alg}_1(\mathbb{S}, \mathbf{2}) & \\
 & \downarrow & \downarrow & \downarrow & \\
 \mathbf{KIMon}_2(\mathbb{T}) & \hookrightarrow & \mathbf{KIMon}_0(\mathbb{T}) = \mathbf{K}(\mathbb{T}) & & \\
 & \searrow \cong & \downarrow & \searrow \cong & \\
 & \mathbf{Alg}_2(\mathbb{S}, \mathbf{2}) & \hookrightarrow & \mathbf{Alg}_0(\mathbb{S}, \mathbf{2}) & \hookrightarrow \mathbf{A}(\mathbb{S}, \mathbf{2})
 \end{array}$$

The case of the inclusion  $\alpha : \mathbb{B} \rightarrow \mathbb{F}$  is treated in [11], which inspired the following presentation.

Until the end of this section, we fix a coherent  $\mathbf{Sup}$ -enriched monad  $(\mathbb{T}, \tau)$  with  $\mathbb{T} = (T, \eta, \mu)$ , a monad  $\mathbb{S} = (S, \delta, \nu)$ , a monad morphism  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ , and we set  $\bar{S} = \alpha^*(T^\tau)$ .

5.7. THE UNDERLYING ADJUNCTION. For sets  $X$  and  $Y$ , we define an adjunction

$$\text{Rel}(SX, Y) \begin{array}{c} \xrightarrow{\phi} \\ \xleftarrow{\perp} \\ \xrightarrow{\psi} \end{array} \text{Set}(Y, TX) \quad ,$$

as follows. By using the isomorphism  $\text{Rel}(SX, Y) \cong \text{Set}(Y, PSX)$  in  $\text{Ord}$ , we define

$$\begin{aligned} \phi : \text{Set}(Y, PSX) &\rightarrow \text{Set}(Y, TX) \quad , & r^b &\mapsto \alpha_X^\vee \cdot r^b \quad , \\ \psi : \text{Set}(Y, TX) &\rightarrow \text{Set}(Y, PSX) \quad , & c &\mapsto \alpha_X^\perp \cdot c \quad . \end{aligned}$$

The adjunction  $\phi \dashv \psi$  follows from  $\alpha_X^\vee \dashv \alpha_X^\perp$ . In pointwise notation,  $\phi$  and  $\psi$  can be expressed by

$$\phi(r)(y) = \bigvee \alpha_X[r^b(y)] \quad \text{and} \quad \psi(c)(\mathbf{x}, y) = \top \iff \alpha_X(\mathbf{x}) \leq c(y) \quad ,$$

for  $r : SX \leftrightarrow Y$ , and  $c : Y \rightarrow TX$ . As usual,  $\phi$  and  $\psi$  restrict to mutually inverse isomorphisms between the sets of fixpoints of  $\psi \cdot \phi$  and of  $\phi \cdot \psi$ , respectively.

5.8. FIXPOINTS OF THE ADJUNCTION. Obviously, the fixpoints of  $\phi \cdot \psi$  are those maps  $c : Y \rightarrow TX$  such that  $c(y)$  is  $\alpha$ -approachable for all  $y \in Y$ . Unwinding the fixpoint condition for  $\psi \cdot \phi$ , we see that the fixpoints are those relations  $r : SX \leftrightarrow Y$  such that  $r^b(y)$  is  $\alpha$ -closed for all  $y \in Y$ , where a subset  $\mathcal{A} \subseteq SX$  is  $\alpha$ -closed if

$$\alpha_X(\mathbf{x}) \leq \bigvee \alpha_X[\mathcal{A}] \implies \mathbf{x} \in \mathcal{A} \quad .$$

The latter terminology is borrowed from [11], where  $\alpha : \mathbb{B} \rightarrow \mathbb{F}$  is the inclusion, and the  $\alpha$ -closed subsets of  $\beta X$  are precisely the closed sets of the so-called Zariski topology, which arises from the free compact Hausdorff space on  $X$ .

While the  $\alpha$ -closed subsets are closed under arbitrary intersection, they are in general not closed under finite unions.

Observe that  $r^b(y)$  is  $1_{\mathbb{T}}$ -closed if and only if  $r$  is *continuous*, that is, if and only if

$$r(\bigvee \mathcal{A}, y) = \bigwedge_{x \in \mathcal{A}} r(x, y)$$

holds for any  $\mathcal{A} \subseteq TX$ . This concept of continuity was used in [20] and [18] to obtain Theorem 5.6 for the special case  $\alpha = 1_{\mathbb{T}}$ .

5.9. LEMMA. *A relation  $r : SX \leftrightarrow Y$  is a fixpoint of  $\psi \cdot \phi$  if and only if it is interpolated by  $\alpha$ , left unitary, and antitone.*

PROOF. Observe that for any antitone and left unitary relation  $r : SX \leftrightarrow Y$ , we have in fact  $r \cdot (\leq_{\overline{S}}) = r$  (see 3.2) and  $\delta_Y^\circ \cdot \overline{S}r \cdot \nu_X^\circ = r$ . Indeed, to see that the left unitary condition  $\delta_Y^\circ \cdot \overline{S}r \cdot \nu_X^\circ \leq r$  is in fact an equality, recall that  $\delta : 1 \rightarrow \overline{S}$  is oplax, so that  $r \leq \delta_Y^\circ \cdot \overline{S}r \cdot \delta_{SX}$ , hence

$$r = r \cdot \delta_{SX}^\circ \cdot \nu_X^\circ \leq \delta_Y^\circ \cdot \overline{S}r \cdot \delta_{SX} \cdot \delta_{SX}^\circ \cdot \nu_X^\circ \leq \delta_Y^\circ \cdot \overline{S}r \cdot \nu_X^\circ \quad .$$



Thus, by applying  $-^b$ , we observe that a relation  $r : SX \leftrightarrow Y$  is left unitary precisely when

$$r^b = P\nu_X \cdot (\overline{Sr})^b \cdot \delta_Y$$

holds, while  $r$  is antitone if and only if  $(\leq_{\overline{S}}^b)^{\mathbb{P}} \cdot r^b = r^b$ .

Suppose that  $r : SX \leftrightarrow Y$  is antitone and left unitary. We have:

$$\begin{aligned} r^b &= (\leq_{\overline{S}}^b)^{\mathbb{P}} \cdot r^b && (r \text{ antitone}) \\ &= (\leq_{\overline{S}}^b)^{\mathbb{P}} \cdot P\nu_X \cdot (\overline{Sr})^b \cdot \delta_Y && (r \text{ left unitary}) \\ &= (\leq_{\overline{S}}^b \cdot \nu_X)^{\mathbb{P}} \cdot (\overline{Sr})^b \cdot \delta_Y . \end{aligned}$$

Therefore, if  $r$  is interpolated by  $\alpha$ , we have  $\alpha_X^\perp \cdot \alpha_X^\vee \cdot r^b \leq (\leq_{\overline{S}}^b \cdot \nu_X)^{\mathbb{P}} \cdot (\overline{Sr})^b \cdot \delta_Y = r^b$ , that is,  $r$  is a fixpoint of  $\psi \cdot \phi$ .

Conversely, any fixpoint  $r$  of  $\psi \cdot \phi$  is of the form  $\psi(c)$  for some  $c$ , hence clearly antitone. Similarly, if  $\overline{Sr}(\mathfrak{X}, \delta_X(y)) = \top$  holds, we can apply  $\mu_X \cdot T\alpha_X$  to each side of the inequality  $\alpha_{SX}(\mathfrak{X}) \leq r^\tau \cdot \eta_X(y)$ , and conclude that  $r(\nu_X(\mathfrak{X}), y) = \top$  by the fixpoint condition. Finally, since any fixpoint is left unitary and antitone, we see that it is interpolated by  $\alpha$  due to the displayed equation above.  $\blacksquare$

5.10. PROPOSITION. *The ordered sets of all  $c : Y \rightarrow TX$  which are pointwise  $\alpha$ -approachable and of all  $r : SX \leftrightarrow Y$  which are antitone, left unitary, and interpolated by  $\alpha$  are isomorphic.*

PROOF. This is immediate from the previous lemma.  $\blacksquare$

5.11. LEMMA. *The map  $\phi$  satisfies*

$$\phi(\delta_X^\circ) = \eta_X \quad \text{and} \quad \phi(s * r) \leq \phi(r) \circ \phi(s)$$

for all  $r : SX \leftrightarrow Y$  and  $s : SY \leftrightarrow Z$ . As a consequence, the conditions

$$\delta_X^\circ \leq \psi(\eta_X) \quad \text{and} \quad \psi(d) * \psi(c) \leq \psi(c \circ d)$$

hold for all  $c : Y \rightarrow TX$ ,  $d : Z \rightarrow TY$ .

Moreover,  $\phi$  satisfies  $\phi(s * r) = \phi(r) \circ \phi(s)$  if and only if  $\alpha$  satisfies the second interpolation condition.

PROOF. By recalling that  $(\delta_X^\circ)^b = \iota_{SX} \cdot \delta_X$ , we obtain  $\phi(\delta_X^\circ) = \alpha_X^\top \cdot \tau_{SX} \cdot \iota_{SX} \cdot \delta_X = \alpha_X^\top \cdot \eta_{SX} \cdot \delta_X = \alpha_X \cdot \delta_X = \eta_X$ . To show  $\phi(s * r) \leq \phi(r) \circ \phi(s)$ , observe that

$$(\alpha_X^\vee \cdot r^b)^\top = (\alpha_X^\top \cdot \tau_{SX} \cdot r^b)^\top = \alpha_X^\top \cdot (\tau_{SX} \cdot r^b)^\top = \alpha_X^\top \cdot r^\tau , \quad (6)$$

and recall from 4.18 that  $\alpha_X^{\mathbb{T}} \cdot r^\tau \cdot \alpha_Y \geq \alpha_X^\vee \cdot P\nu_X \cdot (\overline{S}r)^\flat$  holds. Therefore, we get

$$\begin{aligned} (\alpha_X^\vee \cdot r^\flat)^\mathbb{T} \cdot \alpha_Y^\vee &= \alpha_X^{\mathbb{T}} \cdot (\tau_{SX} \cdot r^\flat)^\mathbb{T} \cdot \alpha_Y^{\mathbb{T}} \cdot \tau_{SY} \\ &= (\alpha_X^{\mathbb{T}} \cdot r^\tau \cdot \alpha_Y)^\mathbb{T} \cdot \tau_{SY} \\ &\geq (\alpha_X^\vee \cdot P\nu_X \cdot (\overline{S}r)^\flat)^\mathbb{T} \cdot \tau_{SY} \\ &= \alpha_X^{\mathbb{T}} \cdot (\tau_{SX} \cdot P\nu_X \cdot (\overline{S}r)^\flat)^\mathbb{T} \cdot \tau_{SY} \\ &= \alpha_X^{\mathbb{T}} \cdot \tau_{SX} \cdot (P\nu_X \cdot (\overline{S}r)^\flat)^\mathbb{P} \\ &= \alpha_X^\vee \cdot (P\nu_X \cdot (\overline{S}r)^\flat)^\mathbb{P} \end{aligned}$$

so that  $\phi(r) \circ \phi(s) = (\alpha_X^\vee \cdot r^\flat)^\mathbb{T} \cdot \alpha_Y^\vee \cdot s^\flat \geq \alpha_X^\vee \cdot P\nu_X \cdot ((\overline{S}r)^\flat)^\mathbb{P} \cdot s^\flat = \alpha_X^\vee \cdot (s \cdot \overline{S}r \cdot \nu_X^\circ)^\flat = \phi(s * r)$ .

If the second interpolation condition holds, then the last two inequalities become equalities. To show the converse claim, observe that  $\alpha_Y = \alpha_Y^{\mathbb{T}} \cdot \eta_{SY} = \alpha_Y^{\mathbb{T}} \cdot \tau_{SY} \cdot \iota_{SY} = \alpha_Y^\vee \cdot \iota_{SY}$  and  $\iota_{SY} = (1_{SY})^\flat$ . Putting these equalities together, using (6) above, and assuming that  $\phi(r) \circ \phi(s) = \phi(s * r)$ , we obtain:

$$\begin{aligned} \alpha_X^{\mathbb{T}} \cdot r^\tau \cdot \alpha_Y &= (\alpha_X^\vee \cdot r^\flat)^\mathbb{T} \cdot (\alpha_Y^\vee \cdot 1_{SY}^\flat) = \phi(r) \circ \phi(1_{SY}) = \phi(1_{SY} * r) \\ &= \alpha_X^\vee \cdot (1_{SY} \cdot \overline{S}r \cdot \nu_X^\circ)^\flat = \alpha_X^\vee \cdot ((\nu_X^\circ)^\flat)^\mathbb{P} \cdot (\overline{S}r)^\flat = \alpha_X^\vee \cdot P\nu_X \cdot (\overline{S}r)^\flat, \end{aligned}$$

so the second interpolation condition holds. The claims for  $\psi$  follow from the adjunction  $\phi \dashv \psi$ .  $\blacksquare$

**5.12. THE MAIN ADJUNCTION.** We are now in a position to prove Theorem 5.6. Define concrete functors

$$\Psi : \mathbf{K}(\mathbb{T}) \rightarrow \mathbf{A}(\mathbb{S}, \mathbf{2}) \quad \text{and} \quad \Phi : \mathbf{A}(\mathbb{S}, \mathbf{2}) \rightarrow \mathbf{K}(\mathbb{T})$$

by applying  $\psi$  and  $\phi$  to the structures; that is,  $\Psi(X, c) = (X, \psi(c))$  and  $\Phi(X, r) = (X, \phi(r))$ . The fact that  $\Psi$  and  $\Phi$  send morphisms to morphisms follows easily from the definitions. These functors yield the isomorphisms of Theorem 5.6. The situation is analyzed in detail in the following result, of which 5.6 is an immediate consequence.

**5.13. PROPOSITION.** *The functor  $\Psi$  is right adjoint to  $\Phi$ . Moreover,  $\Psi$  and  $\Phi$  restrict as the solid arrows in the diagram below, leading to adjunctions between  $\mathbf{Alg}_i(\mathbb{S}, \mathbf{2})$  and  $\mathbf{KIMon}_i(\mathbb{T})$  for  $i = 0, 1$  as in the right vertical face of the diagram below.*

$$\begin{array}{ccc} \mathbf{KIMon}(\mathbb{T}) \hookrightarrow & & \mathbf{KIMon}_1(\mathbb{T}) \\ \downarrow & \begin{array}{c} \swarrow \Psi \\ \searrow \Phi \end{array} & \downarrow \\ & \mathbf{Alg}(\mathbb{S}, \mathbf{2}) \hookrightarrow & \mathbf{Alg}_1(\mathbb{S}, \mathbf{2}) \\ \downarrow & & \downarrow \\ \mathbf{KIMon}_2(\mathbb{T}) \hookrightarrow & & \mathbf{KIMon}_0(\mathbb{T}) \\ \downarrow & \begin{array}{c} \swarrow \Psi \\ \searrow \Phi \end{array} & \downarrow \\ & \mathbf{Alg}_2(\mathbb{S}, \mathbf{2}) \hookrightarrow & \mathbf{Alg}_0(\mathbb{S}, \mathbf{2}) \end{array}$$

If  $\alpha$  satisfies the second interpolation condition, we get restrictions of  $\Phi$  as the dashed arrows above, hence adjunctions between  $\mathbf{KIMon}_2(\mathbb{T})$  and  $\mathbf{Alg}_2(\mathbb{S}, \mathbf{2})$  as well as between  $\mathbf{KIMon}(\mathbb{T})$  and  $\mathbf{Alg}(\mathbb{S}, \mathbf{2})$ .

If  $\alpha$  is sup-generating, then  $\Psi$  and all its restrictions in the diagram above are full reflective embeddings.

If  $\alpha$  is interpolating, then  $\Phi$  and all its restrictions as above are full coreflective embeddings.

PROOF. The adjunction  $\Phi \dashv \Psi$  follows from  $\phi \dashv \psi$ . The other properties are immediate from 5.9–5.11 (recall that  $\alpha$  satisfies the second interpolation condition provided it is sup-generating). ■

5.14. COROLLARY. *There are concrete isomorphisms*

$$\begin{aligned} \mathbf{KIMon}_0(\mathbb{T}) &\cong \mathbf{Alg}_0(\mathbb{T}, \mathbf{2}) \quad , & \mathbf{KIMon}_2(\mathbb{T}) &\cong \mathbf{Alg}_2(\mathbb{T}, \mathbf{2}) \quad , \\ \mathbf{KIMon}_1(\mathbb{T}) &\cong \mathbf{Alg}_1(\mathbb{T}, \mathbf{2}) \quad , & \mathbf{KIMon}(\mathbb{T}) &\cong \mathbf{Alg}(\mathbb{T}, \mathbf{2}) \quad . \end{aligned}$$

PROOF. The monad morphism  $1_{\mathbb{T}}$  is sup-generating and interpolating. ■

5.15. COROLLARY. *If  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  is interpolating in  $(\mathbb{T}, \tau)$ , then  $\mathbf{Alg}_i(\mathbb{S}, \mathbf{2})$  is a coreflective subcategory of  $\mathbf{Alg}_i(\mathbb{T}, \mathbf{2})$  for  $i = 0, 1, 2$ , and  $\mathbf{Alg}(\mathbb{S}, \mathbf{2})$  is a concrete coreflective subcategory of  $\mathbf{Alg}(\mathbb{T}, \mathbf{2})$ . Moreover, if  $\alpha$  is also sup-generating, then these coreflections are isomorphisms.*

PROOF. Simply compose the coreflections from Proposition 5.13 with the isomorphisms from Corollary 5.14. ■

5.16. REMARK. In the case where  $\alpha$  is interpolating, the coreflection  $\mathbf{Alg}(\mathbb{T}, \mathbf{2}) \rightarrow \mathbf{Alg}(\mathbb{S}, \mathbf{2})$  is simply the functor  $F_\alpha$  (considered in 3.12) which sends  $(X, a : TX \leftrightarrow X)$  to  $(X, a \cdot \alpha_X : SX \leftrightarrow X)$ . The concrete coreflective embedding  $\mathbf{Alg}(\mathbb{S}, \mathbf{2}) \rightarrow \mathbf{Alg}(\mathbb{T}, \mathbf{2})$  then sends a structure  $a : SX \leftrightarrow X$  to  $\hat{a} : TX \leftrightarrow X$ , where

$$\hat{a}(f, y) = \bigwedge \{ a(\mathfrak{x}, y) \mid \mathfrak{x} \in SX : \alpha_X(\mathfrak{x}) \leq f \} ,$$

for all  $f \in TX$ ,  $y \in X$ .

5.17. EXAMPLES.

1. The category  $\mathbf{RRel}$  of reflexive relations and relation-preserving maps can be described either as  $\mathbf{Alg}_1(\mathbb{I}, \mathbf{2})$ ,  $\mathbf{KIMon}_1(\mathbb{P})$ , or  $\mathbf{Alg}_1(\mathbb{P}, \mathbf{2})$ , depending on whether a relation  $r$  on a set  $X$  is seen as a map

$$r : X \times X \rightarrow \mathbf{2} \quad , \quad r : X \rightarrow PX \quad , \quad \text{or even} \quad r : PX \times X \rightarrow \mathbf{2}$$

respectively. Similarly, the category  $\mathbf{Ord}$  of preordered sets can be obtained as any of the three categories

$$\mathbf{Alg}(\mathbb{I}, \mathbf{2}) \cong \mathbf{KIMon}(\mathbb{P}) \cong \mathbf{Alg}(\mathbb{P}, \mathbf{2}) .$$

2. The category  $\mathbf{PrTop}$  of pretopological spaces and continuous maps can be presented via convergence of ultrafilters as  $\mathbf{Alg}_1(\mathbb{B}, \mathbf{2})$ , via neighborhood systems as  $\mathbf{KIMon}_1(\mathbb{F}, \mathbf{2})$ , or via convergence of filters as  $\mathbf{Alg}_1(\mathbb{F}, \mathbf{2})$ . The corresponding descriptions of the category  $\mathbf{Top}$  of topological spaces are

$$\mathbf{Alg}(\mathbb{B}, \mathbf{2}) \cong \mathbf{KIMon}(\mathbb{F}) \cong \mathbf{Alg}(\mathbb{F}, \mathbf{2}) .$$

3. The concretely isomorphic categories  $\mathbf{Int}$  and  $\mathbf{Cls}$  of interior and closure spaces respectively can be described either as  $\mathbf{Alg}(\mathbb{U}, \mathbf{2})$  or as  $\mathbf{KIMon}(\mathbb{U}, \mathbf{2})$ . The objects of  $\mathbf{Alg}_1(\mathbb{U}, \mathbf{2}) \cong \mathbf{KIMon}_1(\mathbb{U})$  are *preclosure spaces*, that is, sets  $X$  equipped with a monotone and extensive map  $PX \rightarrow PX$ . The resulting category is denoted by  $\mathbf{PrCls}$ .
4. Thanks to the previous examples, Corollary 5.15 may be applied to the diagram of 2.7(7) to get either of the two chains of coreflective embeddings below:

$$\begin{array}{ccccc} \mathbf{RRel} & \longrightarrow & \mathbf{PrTop} & \longrightarrow & \mathbf{PrCls} \\ \mathbf{Ord} & \longrightarrow & \mathbf{Top} & \longrightarrow & \mathbf{Cls} . \end{array}$$

5. By example 5.2(4) and Theorem 5.6, every  $\mathbb{T}$ -algebra is a  $(\mathbb{T}, \mathbf{2})$ -algebra (for the Kleisli extension  $\overline{T}$ ), so that the latter concept is indeed a generalization of the former. This fact is *a priori* not obvious because in general  $Ta \neq \overline{T}a$  for  $\mathbf{Set}$ -maps  $a : TX \rightarrow X$ , and should be contrasted with the case of Barr's extension that satisfies  $\overline{T}f = Tf$  for any map  $f : X \rightarrow Y$ .

## 6. Extending $\mathbf{Alg}(\mathbb{T}, \mathbf{2})$ to $\mathbf{Alg}(\mathbb{T}, \mathbf{V})$

6.1. TOWER EXTENSIONS. Let us first recall the original definition of a tower extension from [21]. Consider a topological functor  $G : \mathbf{C} \rightarrow \mathbf{Set}$ , and denote by  $G_X$  the fiber of  $G$  over a set  $X$ . By topologicity of  $G$ , every  $G_X$  is a complete lattice with respect to the order

$$A \leq B \iff 1_X : A \rightarrow B \text{ is a } \mathbf{C}\text{-morphism,}$$

for all  $A, B \in \mathbf{C}$  with  $GA = X = GB$ . If  $\mathbf{V}$  is a complete lattice, the *tower extension*  $\mathbf{C}[\mathbf{V}]$  of  $\mathbf{C}$  is the concrete category defined as follows:

- (i) objects are  $\mathbf{V}$ -*towers* of  $\mathbf{C}$ -objects, that is, pairs  $(X, \phi)$ , with  $X$  a set and  $\phi : \mathbf{V}^{\text{op}} \rightarrow G_X$  an inf-map;
- (ii) morphisms  $f : (X, \phi) \rightarrow (Y, \psi)$  are given by maps  $f : X \rightarrow Y$  such that for each  $v \in \mathbf{V}$ ,  $f : \phi(v) \rightarrow \psi(v)$  is a  $\mathbf{C}$ -morphism.

For  $v \in \mathbf{V}$ , we call  $\phi(v)$  the *component* of  $\phi$  at  $v$ . Observe that we have  $\mathbf{C}[\mathbf{1}] \cong \mathbf{Set}$  and  $\mathbf{C}[\mathbf{2}] \cong \mathbf{C}$ .

6.2. PROPOSITION. Consider a monad  $\mathbb{T} = (T, \eta, \mu)$  on  $\mathbf{Set}$  with a lax extension  $\bar{T}$  of  $T$  to  $\mathbf{Rel}$ . The concrete functors

$$\mathbf{A}(\mathbb{T}, \mathbf{V}) \rightarrow \mathbf{A}(\mathbb{T}, \mathbf{2})[\mathbf{V}], \quad (X, a) \mapsto (X, (v \mapsto a_v)),$$

and

$$\mathbf{A}(\mathbb{T}, \mathbf{2})[\mathbf{V}] \rightarrow \mathbf{A}(\mathbb{T}, \mathbf{V}), \quad (X, \phi) \mapsto (X, r_\phi),$$

are mutually inverse and restrict to an isomorphism  $\mathbf{A}_0(\mathbb{T}, \mathbf{V}) \cong \mathbf{A}_0(\mathbb{T}, \mathbf{2})[\mathbf{V}]$ .

PROOF. Using the laws from 2.4, one checks easily that the assignments above define functors. The fact that they are mutually inverse also follows from 2.4. ■

6.3. PROPOSITION. Let  $\mathbb{T} = (T, \eta, \mu)$  be a monad on  $\mathbf{Set}$  with a lax  $\mathbf{Rel}$ -extension  $\bar{T}$  of  $T$ . For any  $(X, a) \in \mathbf{A}(\mathbb{T}, \mathbf{V})$ , we have

- $a : TX \rightarrow X$  is reflexive if and only if  $a_k \in \mathbf{Rel}(TX, X)$  is reflexive.
- $a$  is transitive if and only if  $a_v * a_u \leq a_{u \otimes v}$  holds for all  $u, v \in \mathbf{V}$ .

PROOF. The first statement is obvious.

Assume now that  $(X, a)$  is transitive. It suffices to show that  $a_v \cdot (T_{\mathbf{V}}a)_u \leq a_{u \otimes v} \cdot \mu_X$ . For  $x \in X$ , let  $\mathfrak{X} \in (a_v \cdot (T_{\mathbf{V}}a)_u)^b(x)$ , so there exists  $\mathfrak{x} \in TX$  with  $v \leq a(\mathfrak{x}, x)$  and  $u \leq T_{\mathbf{V}}a(\mathfrak{X}, \mathfrak{x})$ . Therefore,

$$u \otimes v \leq \bigvee_{\mathfrak{x} \in TX} T_{\mathbf{V}}a(\mathfrak{X}, \mathfrak{x}) \otimes a(\mathfrak{x}, x) = a \cdot T_{\mathbf{V}}a(\mathfrak{X}, x) \leq a \cdot \mu_X(\mathfrak{X}, x).$$

Since  $(a \cdot \mu_X)_w = a_w \cdot \mu_X$  holds, we are done.

Conversely, assume that  $a_v * a_u \leq a_{u \otimes v}$  holds for all  $u, v \in \mathbf{V}$ . It suffices to show  $T_{\mathbf{V}}a(\mathfrak{X}, \mathfrak{x}) \otimes a(\mathfrak{x}, x) \leq a(\mu_X(\mathfrak{X}), x)$  for all  $x \in X$ ,  $\mathfrak{x} \in TX$ ,  $\mathfrak{X} \in TTX$ . Fix such  $x, \mathfrak{x}, \mathfrak{X}$  and set  $v := a(\mathfrak{x}, x)$ ,  $\mathcal{A} := \{u \in \mathbf{V} \mid Ta_u(\mathfrak{X}, \mathfrak{x}) = \top\}$ . We have  $T_{\mathbf{V}}a(\mathfrak{X}, \mathfrak{x}) \otimes a(\mathfrak{x}, x) = \bigvee_{u \in \mathcal{A}} u \otimes v$ . For any  $u \in \mathcal{A}$ , we have  $a_v \cdot Ta_u \leq a_{u \otimes v} \cdot \mu_X$ , so  $u \otimes v \leq a(\mu_X(\mathfrak{X}), x)$ , and the claim is proved. ■

6.4. REFLEXIVE AND TRANSITIVE TOWERS. The previous result motivates the following definitions. We say that  $(X, \phi) \in \mathbf{A}(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  is

- *reflexive* if  $\phi(k)$  is reflexive,
- *transitive* if  $\phi(u) * \phi(v) \leq \phi(v \otimes u)$  holds for all  $u, v \in \mathbf{V}$ , and
- *lax* if it is both reflexive and transitive.

Any lax tower  $(X, \phi)$  in  $\mathbf{A}(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  is componentwise left unitary, hence an element of  $\mathbf{Alg}_0(\mathbb{T}, \mathbf{2})[\mathbf{V}]$ . Indeed, we have:

$$\eta_X^\circ * \phi(v) \leq \phi(k) * \phi(v) \leq \phi(k \otimes v) = \phi(v).$$

6.5. PROPOSITION. *The functors of Proposition 6.2 restrict to concrete isomorphisms between*

- $\mathbf{Alg}_1(\mathbb{T}, \mathbf{V})$  and the full subcategory of  $\mathbf{Alg}_0(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  whose objects are the reflexive towers;
- $\mathbf{Alg}_2(\mathbb{T}, \mathbf{V})$  and the full subcategory of  $\mathbf{Alg}_0(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  whose objects are the transitive towers;
- $\mathbf{Alg}(\mathbb{T}, \mathbf{V})$  and the full subcategory of  $\mathbf{A}(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  whose objects are the lax towers.

PROOF. This is immediate. ■

6.6. COROLLARY. *If  $\mathbf{V}$  is integral, then  $\mathbf{Alg}_1(\mathbb{T}, \mathbf{V})$  and  $\mathbf{Alg}_1(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  are concretely isomorphic.*

PROOF. The statement follows from the previous result and the fact that a tower  $(X, \phi)$  in  $\mathbf{Alg}_0(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  is reflexive if and only if  $\phi(v)$  is reflexive for each  $v \leq k$ . ■

6.7. COROLLARY. *If  $\mathbf{V}$  is a frame, then  $\mathbf{Alg}(\mathbb{T}, \mathbf{V})$  and  $\mathbf{Alg}(\mathbb{T}, \mathbf{2})[\mathbf{V}]$  are concretely isomorphic.*

PROOF. Assume that  $\otimes = \wedge$  and each  $\phi(v)$  is transitive. Take any  $u, v \in \mathbf{V}$ . For  $w = u \wedge v$  we have  $\phi(u) \leq \phi(w)$  and  $\phi(v) \leq \phi(w)$ , so that  $\phi(u) * \phi(v) \leq \phi(w) * \phi(w) \leq \phi(w) = \phi(v \wedge u)$ . The conclusion follows from the previous results. ■

6.8. REMARK. Observe that if  $(X, \phi)$  is transitive, then  $\phi(v)$  is transitive whenever  $v$  is idempotent in  $\mathbf{V}$ . In particular,  $\phi(k)$  is transitive for any transitive tower. This simple remark turns out to be useful in a number of results where an idempotent structure is sought in the multitude of  $\phi(v)$ ,  $v \in \mathbf{V}$ .

6.9. TOWER EXTENSIONS OF KLEISLI MONOIDS. Let  $\mathbf{KIMon}(\mathbb{T}, \mathbf{V})$  denote the full subcategory of  $\mathbf{KIMon}(\mathbb{T})[\mathbf{V}]$  whose objects are lax towers. Thus, an object of  $\mathbf{KIMon}(\mathbb{T}, \mathbf{V})$  is a pair  $(X, (a^v : X \rightarrow TX)_{v \in \mathbf{V}})$  where  $X$  is a set and the family  $(a^v)$  satisfies

$$a^{\vee W} = \bigwedge_{w \in W} a^w, \quad \eta_X \leq a^k, \quad a^v \circ a^u \leq a^{u \otimes v},$$

for all  $W \subseteq \mathbf{V}$  and  $u, v \in \mathbf{V}$ .

6.10. THEOREM. *Let  $(\mathbb{T}, \tau)$  be a coherent  $\mathbf{Sup}$ -enriched monad, and  $\alpha : \mathbb{S} \rightarrow \mathbb{T}$  an interpolating and sup-generating monad morphism. With respect to the extension  $\overline{\mathbb{S}} = \alpha^*(T_V^\tau)$ , there is a concrete isomorphism*

$$\mathbf{Alg}(\mathbb{S}, \mathbf{V}) \cong \mathbf{KIMon}(\mathbb{T}, \mathbf{V}).$$

PROOF. By Lemma 5.11, the isomorphism  $\mathbf{Alg}(\mathbb{S}, \mathbf{2}) \cong \mathbf{KIMon}(\mathbb{T})$  from Theorem 5.6 is compatible with the operations  $*$  and  $\circ$  on the fibers of  $\mathbf{Alg}(\mathbb{S}, \mathbf{2})$  and  $\mathbf{KIMon}(\mathbb{T})$ , respectively. Hence, this isomorphism lifts to the lax tower extensions. ■

6.11. **COROLLARY.** *Let  $(\mathbb{T}, \tau)$  be a coherent **Sup**-enriched monad. With respect to the extension  $T_{\mathbf{V}}^{\tau}$ , we have a concrete isomorphism  $\mathbf{Alg}(\mathbb{T}, \mathbf{V}) \cong \mathbf{KIMon}(\mathbb{T}, \mathbf{V})$ .*

**PROOF.** One just uses the fact that  $1_{\mathbb{T}}$  is interpolating and sup-generating. (Note that a similar result has been obtained in [20] under the additional assumption that  $\mathbf{V}$  is constructively complete distributive). ■

6.12. **REMARK.** The previous results suggest that categories fibered in “complete lattices with unital convolutions” provide an adequate setting in which to study laxly indexed towers. The latter are closely related to *continuous relational presheaves* on the one-object quantaloid  $\mathbf{V}$  (see [17]).

6.13. **EXAMPLES.**

1. The category **PrApp** of preapproach spaces [13] can be described as any one of the concretely isomorphic categories below

$$\mathbf{Alg}_1(\mathbb{B}, \mathbf{P}_+) \cong \mathbf{KIMon}_1(\mathbb{F}, \mathbf{P}_+) \cong \mathbf{Alg}_1(\mathbb{F}, \mathbf{P}_+),$$

and the full subcategory **App** of approach spaces [14] can be described as

$$\mathbf{Alg}(\mathbb{B}, \mathbf{P}_+) \cong \mathbf{KIMon}(\mathbb{F}, \mathbf{P}_+) \cong \mathbf{Alg}(\mathbb{F}, \mathbf{P}_+).$$

2. As in [21], the categories of “fuzzy neighborhood convergence spaces” and of “fuzzy neighborhood spaces” are obtained as

$$\begin{aligned} \mathbf{Alg}_1(\mathbb{B}, [0, 1]) &\cong \mathbf{KIMon}_1(\mathbb{F}, [0, 1]) \cong \mathbf{Alg}_1(\mathbb{F}, [0, 1]), \\ \mathbf{Alg}(\mathbb{B}, [0, 1]) &\cong \mathbf{KIMon}(\mathbb{F}, [0, 1]) \cong \mathbf{Alg}(\mathbb{F}, [0, 1]). \end{aligned}$$

where  $[0, 1]$  carries the quantale structure given by  $\wedge$ . Analogous results hold for  $\mathbb{U}$ , giving rise to various extensions of the category of interior (or closure) spaces.

3.  $\mathbf{Alg}(\mathbb{B}, \mathbf{2}^2)$  is the category **BiTop** of bitopological spaces. This follows from 6.14 below. We can also describe **BiTop** as  $\mathbf{KIMon}(\mathbb{F}, \mathbf{2}^2)$ , or as  $\mathbf{Alg}(\mathbb{F}, \mathbf{2}^2)$ . Similar results hold for interior spaces, pretopological spaces, *etc.*
4. Tower extensions allow to effectively describe  $(\mathbb{T}, \mathbf{V})$ -algebras for simple lattices  $\mathbf{V}$ . For instance,  $\mathbf{Alg}(\mathbb{B}, \{0, 1, 2\})$ , where  $\{0, 1, 2\}$  carries  $\wedge$  as tensor, is concretely isomorphic to the category which has as objects triples  $(X, \tau_0, \tau_1)$ , where  $\tau_0 \subseteq \tau_1$  are topologies, and as morphisms maps which are separately continuous.
5. Another interesting example is given by the identity monad together with the non-integral three-chain  $\mathbf{3}$  of Example 2.2(4). Indeed, objects of  $\mathbf{Alg}(\mathbb{I}, \mathbf{3})$  are sets  $X$  with a map  $\phi : \mathbf{3} \rightarrow \mathbf{Rel}(X, X)$  such that  $\phi(\perp) = r_{\perp}$  satisfies  $r_{\perp}(x, x) = \top$  for all  $x \in X$ ,  $\phi(k) = r_k$  is a reflexive and transitive relation, and  $\phi(\top) = r_{\top}$  is a relation such that  $r_{\top} \leq r_k$ ,  $r_{\top} \cdot r_k \leq r_{\top}$ , and  $r_k \cdot r_{\top} \leq r_{\top}$ ; morphisms of  $\mathbf{Alg}(\mathbb{I}, \mathbf{3})$  are maps

that preserve the relations  $r_k$  and  $r_\top$ . Therefore,  $\mathbf{Alg}(\mathbb{I}, \mathbf{3})$  is concretely isomorphic to the category whose objects are preordered sets  $(X, \leq)$  equipped with an *auxiliary relation* (see [8]), that is, a relation  $\prec$  on  $X$  such that

$$x \prec y \implies x \leq y \quad \text{and} \quad x \leq x' \prec y' \leq y \implies x \prec y,$$

for all  $x, x', y, y' \in X$ , and whose morphisms are  $\prec$ -preserving monotone maps.

**6.14. PROPOSITION.** *Let  $I$  be a set. For any topological category  $G : \mathbf{C} \rightarrow \mathbf{Set}$ , the tower extension  $\mathbf{C}[PI]$  by the powerset of  $I$  is concretely isomorphic to the category  $I\text{-}\mathbf{C}$ , whose objects are pairs  $(X, (A_i)_I)$  (with  $X$  a set, and each  $A_i$  an element of the fibre  $G_A$ ), and morphisms  $f : (X, (A_i)) \rightarrow (Y, (B_i))$  are maps  $f : X \rightarrow Y$  such that  $f : A_i \rightarrow B_i$  is a  $\mathbf{C}$ -morphism for each  $i \in I$ .*

**PROOF.** Recall that  $PI$  is the free **Sup**-lattice on  $I$ , therefore maps  $s : I \rightarrow G_B$  correspond to inf-maps  $\bar{s} : PI^{\text{op}} \rightarrow G_B$  via  $A \mapsto \bigwedge s(a)$ , where  $\bigwedge$  denotes the infimum in the fiber, that is, the initial lift.

Thus, we only need to show that this assignment is compatible with morphisms in the following sense: if  $f : A_j \rightarrow B_j$  is a morphism for each  $j \in J$ ,  $J \subseteq I$ , then  $f : (\bigwedge_J A_j) \rightarrow (\bigwedge_J B_j)$  is also a morphism. To prove this, recall that each map  $f : X \rightarrow Y$  gives rise to a monotone map  $f_* : G_X \rightarrow G_Y$  by forming final lifts along  $f$ . Thus, for each  $i \in J$  we have  $f_*(\bigwedge A_j) \leq f_*(A_i) \leq B_i$  since  $f : A_i \rightarrow B_i$  is a morphism. Hence,  $f_*(\bigwedge A_j) \leq \bigwedge B_j$  also holds, that is,  $f$  is a morphism  $(\bigwedge A_j) \rightarrow (\bigwedge B_j)$ . ■

## References

- [1] J. Adámek, H. Herrlich, and G.E. Strecker. Abstract and concrete categories: The joy of cats. *Repr. Theory Appl. Categ.*, (17):1–507 (electronic), 2006. Reprint of the 1990 original.
- [2] M. Barr. Relational algebras. In *Reports of the Midwest Category Seminar, IV*, number 137 in Lecture Notes in Mathematics, pages 39–55. Springer, 1970.
- [3] M.M. Clementino and D. Hofmann. On extensions of lax monads. *Theory Appl. Categ.*, 13(3):41–60 (electronic), 2004.
- [4] M.M. Clementino, D. Hofmann, and W. Tholen. One setting for all: Metric, topology, uniformity, approach structure. *Appl. Categ. Structures*, 12(2):127–154, 2004.
- [5] E. Colebunders and R. Lowen. Lax monads and algebras via initial monad morphisms. 2007. Preprint.
- [6] A. Day. Filter monads, continuous lattices and closure systems. *Canad. J. Math.*, 27:50–59, 1975.



- [7] B. Fawcett and R.J. Wood. Constructive complete distributivity. I. *Math. Proc. Camb. Phil. Soc.*, 107(1):81–89, 1990.
- [8] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *Continuous Lattices and Domains*. Number 93 in Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2003.
- [9] F. Hausdorff. *Grundzüge der Mengenlehre*. Chelsea, 1949.
- [10] D. Hofmann. Topological theories and closed objects. *Adv. Math.*, 215(2):789–824, 2007.
- [11] D. Hofmann and W. Tholen. Kleisli operations for topological spaces. *Topology Appl.*, 153(15):2952–2961, 2006.
- [12] U. Höhle. *Many Valued Topology and its Applications*. Kluwer, Boston, 2001.
- [13] E. Lowen and R. Lowen. Topological quasitopos hulls of categories containing topological and metric objects. *Cahiers Top. et Géom. Diff.*, 30(3):213–228, 1989.
- [14] R. Lowen. *Approach Spaces*. Oxford Mathematical Monographs. Oxford University Press, New York, 1997. The missing link in the topology-uniformity-metric triad.
- [15] E.G. Manes. *Algebraic Theories*. Springer-Verlag, New York, 1976. Graduate Texts in Mathematics, No. 26.
- [16] E.G. Manes. Taut monads and  $T_0$ -spaces. *Theoret. Comput. Sci.*, 275(1–2):79–109, 2002.
- [17] K.I. Rosenthal. *The Theory of Quantaloids*, volume 348 of *Pitman Research Notes in Mathematics Series*. Longman, Harlow, 1996.
- [18] C. Schubert. *Lax Algebras—A Scenic Approach*. PhD thesis, Universität Bremen, 2006.
- [19] G.J. Seal. Canonical and op-canonical lax algebras. *Theory Appl. Categ.*, 14(10):221–243 (electronic), 2005.
- [20] G.J. Seal. A Kleisli-based approach to lax algebras. *Appl. Categ. Structures*, 2007. To appear.
- [21] D. Zhang. Tower extension of topological constructs. *Comment. Math. Univ. Carolin.*, 41(1):41–51, 2000.

*Fakultät für Informatik, Lehrstuhl X  
Technische Universität Dortmund  
44221 Dortmund, Germany*

*School of Computer Science  
McGill University  
H3A 2A7 Montreal, Canada*

Email: `christoph.schubert@uni-dortmund.de`  
`gseal@fastmail.fm`

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/21/7/21-07.{dvi,ps,pdf}>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

**SUBSCRIPTION INFORMATION.** Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to [tac@mta.ca](mailto:tac@mta.ca) including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca).

**INFORMATION FOR AUTHORS.** The typesetting language of the journal is  $\text{T}_{\text{E}}\text{X}$ , and  $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\text{e}$  strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

**MANAGING EDITOR.** Robert Rosebrugh, Mount Allison University: [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca)

**$\text{T}_{\text{E}}\text{X}$  NICAL EDITOR.** Michael Barr, McGill University: [barr@math.mcgill.ca](mailto:barr@math.mcgill.ca)

**ASSISTANT  $\text{T}_{\text{E}}\text{X}$  EDITOR.** Gavin Seal, McGill University: [gavin\\_seal@fastmail.fm](mailto:gavin_seal@fastmail.fm)

**TRANSMITTING EDITORS.**

Richard Blute, Université d' Ottawa: [rblute@uottawa.ca](mailto:rblute@uottawa.ca)

Lawrence Breen, Université de Paris 13: [breen@math.univ-paris13.fr](mailto:breen@math.univ-paris13.fr)

Ronald Brown, University of North Wales: [ronnie.profbrown \(at\) btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Aurelio Carboni, Università dell' Insubria: [aurelio.carboni@uninsubria.it](mailto:aurelio.carboni@uninsubria.it)

Valeria de Paiva, Cuill Inc.: [valeria@cuill.com](mailto:valeria@cuill.com)

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Martin Hyland, University of Cambridge: [M.Hyland@dpms.cam.ac.uk](mailto:M.Hyland@dpms.cam.ac.uk)

P. T. Johnstone, University of Cambridge: [ptj@dpms.cam.ac.uk](mailto:ptj@dpms.cam.ac.uk)

Anders Kock, University of Aarhus: [kock@imf.au.dk](mailto:kock@imf.au.dk)

Stephen Lack, University of Western Sydney: [s.lack@uws.edu.au](mailto:s.lack@uws.edu.au)

F. William Lawvere, State University of New York at Buffalo: [wlawvere@acsu.buffalo.edu](mailto:wlawvere@acsu.buffalo.edu)

Jean-Louis Loday, Université de Strasbourg: [loday@math.u-strasbg.fr](mailto:loday@math.u-strasbg.fr)

Ieke Moerdijk, University of Utrecht: [moerdijk@math.uu.nl](mailto:moerdijk@math.uu.nl)

Susan Niefield, Union College: [niefiels@union.edu](mailto:niefiels@union.edu)

Robert Paré, Dalhousie University: [pare@mathstat.dal.ca](mailto:pare@mathstat.dal.ca)

Jiri Rosicky, Masaryk University: [rosicky@math.muni.cz](mailto:rosicky@math.muni.cz)

Brooke Shipley, University of Illinois at Chicago: [bshipley@math.uic.edu](mailto:bshipley@math.uic.edu)

James Stasheff, University of North Carolina: [jds@math.unc.edu](mailto:jds@math.unc.edu)

Ross Street, Macquarie University: [street@math.mq.edu.au](mailto:street@math.mq.edu.au)

Walter Tholen, York University: [tholen@mathstat.yorku.ca](mailto:tholen@mathstat.yorku.ca)

Myles Tierney, Rutgers University: [tierney@math.rutgers.edu](mailto:tierney@math.rutgers.edu)

Robert F. C. Walters, University of Insubria: [robert.walters@uninsubria.it](mailto:robert.walters@uninsubria.it)

R. J. Wood, Dalhousie University: [rjwood@mathstat.dal.ca](mailto:rjwood@mathstat.dal.ca)