

SITE CHARACTERIZATIONS FOR GEOMETRIC INVARIANTS OF TOPOSES

OLIVIA CARMELLO

ABSTRACT. We discuss the problem of characterizing the property of a Grothendieck topos to satisfy a given ‘geometric’ invariant as a property of its sites of definition, and indicate a set of general techniques for establishing such criteria. We then apply our methodologies to specific invariants, notably including the property of a Grothendieck topos to be localic (resp. atomic, locally connected, equivalent to a presheaf topos), obtaining explicit site characterizations for them.

1. Introduction

In this paper we provide a set of general methodologies for obtaining bijective site characterizations for ‘geometric’ invariants of toposes, that is criteria of the kind ‘a topos $\mathbf{Sh}(\mathcal{C}, J)$ satisfies the property I if and only if the site (\mathcal{C}, J) satisfies a property $P_{(\mathcal{C}, J)}$ (explicitly written in the language of the site (\mathcal{C}, J))’, holding for any site (\mathcal{C}, J) or for appropriate classes of sites.

Throughout the past years, site characterizations have been established for several important geometric invariants of toposes, including the property of a topos to be atomic, locally connected, equivalent to a presheaf topos etc. (cf. in particular [Barr and Diaconescu, 1980] and [Barr and Paré, 1980], and [Johnstone, 2002] as a general reference); however, all of these characterizations are of form ‘A Grothendieck topos satisfies an invariant I if and only if *there exists* a site of definition (\mathcal{C}, J) of it satisfying a certain property $P_{(\mathcal{C}, J)}$ ’. As such, these characterizations are only partially satisfactory, since they allow to infer properties of toposes $\mathbf{Sh}(\mathcal{C}, J)$ starting from properties of sites (\mathcal{C}, J) but not conversely; in fact, not even the proofs of these results provide information which one can exploit to obtain site characterizations going in the other direction.

In order to obtain bijective site characterizations, the problem thus needs to be completely reconsidered and approached from a different angle; we do so in this paper, by adopting the point of view of separating sets of toposes. In fact, it turns out that most of the geometric invariants of toposes considered in the literature, notably including the property of a topos to be localic (resp. atomic, locally connected, equivalent to a presheaf topos, coherent), can be expressed in terms of the existence of a separating set of objects for the topos satisfying some invariant property. In this paper, amongst other things, we

Received by the editors 2012-02-29 and, in revised form, 2012-11-18.


Transmitted by Jiri Rosicky. Published on 2012-11-23. Revision published 2013-02-20; see footnotes. 2010 Mathematics Subject Classification: 03G30, 18C10, 18B25.

Key words and phrases: Grothendieck topos, site characterizations, geometric logic.

© Olivia Caramello, 2012. Permission to copy for private use granted.

show that expressing topos-theoretic invariants in terms of the existence of a separating set of objects of the topos satisfying some property paves the way for natural ‘unravelings’ of such invariants as properties of the sites of definition of the topos, and hence for criteria of the desired form.

Concerning the applicability of the results obtained in this paper, we remark that bijective site characterizations of the kind ‘a topos $\mathbf{Sh}(\mathcal{C}, J)$ satisfies the property I if and only if the site (\mathcal{C}, J) satisfies a property $P_{(\mathcal{C}, J)}$ are particularly relevant in connection to the philosophy ‘toposes as bridges’ introduced in [Caramello, 2010]. Specifically, in [Caramello, 2010] it is argued that Grothendieck toposes can effectively act as unifying spaces in Mathematics serving as ‘bridges’ for transferring information between distinct mathematical theories. The transfer of information between Morita-equivalent theories (i.e., theories classified by the same topos), represented by different sites of definition (\mathcal{C}, J) and (\mathcal{C}', J') of their classifying topos, takes place by expressing topos-theoretic invariant properties (resp. constructions) on the topos in terms of properties (resp. constructions) of its two different sites of definition. For each invariant, we thus have a ‘bridge’



whose ‘deck’ is the equivalence of toposes $\mathbf{Sh}(\mathcal{C}, J) \simeq \mathbf{Sh}(\mathcal{C}', J')$ and whose ‘arches’, represented by the dashed arrows, are given by site characterizations corresponding to the given invariant.

Now, in presence of any Morita-equivalence, any such criterion gives us the possibility to operate an automatic transfer of information between the two theories, leading to concrete mathematical results of various nature. Particular cases of these general results have already been applied by the author in several different contexts (cf. for example [Caramello, 2012c] and [Caramello, 2012b]), and in fact the primary aim of this paper is to make a systematic investigation of the problem of obtaining site characterizations which can be conveniently applied in connection to our general philosophy ‘toposes as bridges’ (notice that, on the other hand, the classical characterizations cannot be directly applied in connection to this philosophy, since the criteria that they give rise to only allow one to enter a given bridge (i.e., to pass from the property $P_{(\mathcal{C}, J)}$ of the site (\mathcal{C}, J) to the invariant I) and not to exit from it).

The paper is organized as follows. In section 2, we make some general remarks about the problem of obtaining site characterizations for geometric invariants of toposes, leading to a metatheorem giving sufficient conditions for a topos-theoretic invariant (of a general specified form) to admit bijective site characterizations holding for a given class of sites, while in the following sections we investigate the specific invariants mentioned above in more detail, obtaining natural site characterizations for them of the desired kind. Besides their technical interest, these results are meant to provide the reader with a general idea

of how the technique ‘toposes as bridges’ introduced in [Caramello, 2010] actually works in a variety of different cases.

This work should be considered as a companion to [Caramello, 2012b], where several syntactic characterizations of geometric invariants on toposes in terms of the theories classified by them were obtained.

1.1. TERMINOLOGY AND NOTATION Our terminology and notation is borrowed from [Johnstone, 2002], if not otherwise indicated.

Moreover, we will employ the following conventions.

Given a Grothendieck site (\mathcal{C}, J) , we denote by $a_J : [\mathcal{C}^{\text{op}}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}, J)$ the associated sheaf functor, by c_J the universal closure operation on subobjects of $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ corresponding to it and by η the unit of the adjunction between a_J and the canonical inclusion $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$. We denote by $l : \mathcal{C} \rightarrow \mathbf{Sh}(\mathcal{C}, J)$ the composite of the Yoneda embedding $Y : \mathcal{C} \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ with the associated sheaf functor $a_J : [\mathcal{C}^{\text{op}}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}, J)$.

To mean that c is an object of a category \mathcal{C} , we simply write $c \in \mathcal{C}$.

All the toposes considered in this paper will be Grothendieck toposes, if not otherwise stated.

2. Geometric invariants of toposes

Several topologically-inspired invariants of Grothendieck toposes have been considered in the literature. In fact, as emphasized by Grothendieck himself, a topos can be conveniently considered as a generalized space apt to be studied by adopting a topological intuition. Indeed, a topos $\mathbf{Sh}(\mathcal{C}, J)$ can be seen as a sort of completion of the site (\mathcal{C}, J) , on which one can define invariants which correspond to (in the sense of being logically equivalent, or implied by) natural ‘geometric’ properties of sites, thus representing analogues, in the topos-theoretic setting, of classical properties of topological spaces.

The natural topos-theoretic analogue of the notion of basis of a topological space is the notion of separating set of objects of a topos. Indeed, a basis of a topological space X can be considered as a full subcategory \mathcal{B} of the poset category $\mathcal{O}(X)$ of open sets of X which is J_X -dense, where J_X is the canonical topology on $\mathcal{O}(X)$; similarly, a separating set of a topos \mathcal{E} can be regarded as a full subcategory of \mathcal{E} which is $J_{\mathcal{E}}$ -dense, where $J_{\mathcal{E}}$ is the canonical topology on the topos \mathcal{E} . Note that if \mathcal{B} is a basis of a topological space X then

$$\mathbf{Sh}(X) \simeq \mathbf{Sh}(\mathcal{B}, J_X|_{\mathcal{B}}),$$

by Grothendieck’s Comparison Lemma; similarly, if \mathcal{C} is a separating set of a topos \mathcal{E} then

$$\mathcal{E} \simeq \mathbf{Sh}(\mathcal{C}, J_{\mathcal{E}}|_{\mathcal{C}}).$$

The notion of separating set is intimately related to that of site; in fact, the separating sets of a topos \mathcal{E} are (up to the obvious notion of isomorphism) precisely the sets of the form $L_{(\mathcal{C}, J)} := \{l(c) \mid c \in \mathcal{C}\}$, where (\mathcal{C}, J) is a site of definition of \mathcal{E} and $l : \mathcal{C} \rightarrow \mathbf{Sh}(\mathcal{C}, J)$

is the functor given by the composite of the Yoneda embedding with the associated sheaf functor.

We can thus naturally expect the properties of (sober) topological spaces which can be expressed in terms of the existence of a basis for the space satisfying a certain property P to be naturally generalizable to the topos-theoretic setting, by replacing bases with separating sets and the property P with an appropriate topos-theoretic analogue. For example, the property of a topos to be atomic (resp. locally connected, coherent) represents a natural topos-theoretic analogue of the property of a space to be discrete (resp. locally connected, coherent).

Of course, not all the natural properties of topological spaces can be expressed in terms of the existence of a basis satisfying a certain condition; for instance, many can be expressed as frame-theoretic properties P of the top element of their frame of open sets, in which case natural topos-theoretic analogues of them can be obtained by replacing the top element of the frame of open sets with the terminal object of the topos and the property P with an appropriate topos-theoretic analogue. For example, the property of a topos to be two-valued (resp. compact) represents a natural topos-theoretic analogue of the property of a topological space to be trivial (resp. compact).

In this paper we shall mostly be concerned with invariants of the first kind, that is of the form ‘to have a separating set of objects satisfying some property P ’, but the techniques that we shall elaborate will be also adaptable to invariants of the second kind.

The problem of finding effective site characterizations for invariants of one kind or another of course admits a satisfactory solution or not depending on the specific invariant under consideration; nonetheless, one can identify some properties which are responsible for such invariants to admit explicit site characterizations.

A general remark which, as we will see, turns out to be extremely useful in practice is the following: if the property P descends along epimorphisms (that is, for any epimorphism $f : A \rightarrow B$ in the topos, if A satisfies P then B satisfies P) then one can try to obtain an explicit site characterization of the invariant ‘to have a separating set of objects satisfying the property P ’, as follows. A topos $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of objects satisfying P if and only if every object of $\mathbf{Sh}(\mathcal{C}, J)$ of the form $l(c)$ is covered by an epimorphic family of arrows whose domains satisfy P . Clearly, if the property P descends along epimorphisms then we can suppose, without loss of generality, this family of arrows to consist entirely of monomorphisms, which can be supposed to be, up to isomorphism, of the form $a_J(S \rightarrow \mathcal{C}(-, c))$ for some (J -closed) sieve S on c . Therefore, provided that the property P is sufficiently well-behaved to the extent of admitting an ‘unraveling’ of the condition of an object of the kind $a_J(S)$ to satisfy P as an explicit condition on S written in the language of the site (\mathcal{C}, J) , we have an explicit site characterization for our invariant of the required form. Examples of application of this method are given in sections 3, 4, 5 and 7.

Of course, if the property P does not descend along epimorphisms, it still make sense to look for explicit site characterizations for the given invariant, holding for large classes of sites if not for all sites; an example is given by the property of a topos to be equivalent

to a presheaf topos, which is expressible as the requirement ‘to have a separating set of irreducible objects’, which we shall discuss in section 6. In fact, for any given topos-theoretic invariant I which can be expressed as the existence of a set of objects of the topos satisfying some invariant property P , the essential condition is that it be provable, for any site (\mathcal{C}, J) (or at least for some identifiable classes of them), that the topos $\mathbf{Sh}(\mathcal{C}, J)$ satisfies I if and only if there exists a set of objects of $\mathbf{Sh}(\mathcal{C}, J)$ satisfying property P all of which are (provably) isomorphic to an object of the form $a_J(S)$ for some (J -closed) sieve S on \mathcal{C} .

On the other hand, many geometrically motivated invariant properties of toposes can be naturally expressed in terms of the existence of a separating set for the topos satisfying a property which cannot be expressed as the requirement that all the objects in the separating set satisfy a certain condition. For example, the property of a topos to be coherent can be expressed as the existence of a separating set of compact objects which is closed under finite limits in the topos (cf. [Carmello, 2012b]). Invariants of this kind are in general more hardly tractable, from the point of view of site characterizations, than those discussed above; nonetheless, as witnessed by the past literature on the subject (cf. for example [Johnstone, 2002]), partial or even complete characterizations for them (holding for large classes of sites) can be achieved, by exploiting the particular ‘combinatorics’ of the sites of definition of the topos in relation to the given invariant.

In connection with the problem of ‘unraveling’ the condition on a set of object of the form $a_J(S)$ to satisfy an invariant property P as an explicit condition on S written in the language of the site (\mathcal{C}, J) , we remark that if P can be logically formulated in terms of the condition that certain monomorphisms m in the topos $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ be c_J -dense, where c_J is the universal closure operation on subobjects of $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ corresponding to the canonical geometric inclusion $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$, and such m are built out of objects of the form S (and arrows between them) through topos-theoretic constructions (that is, small limits, colimits, exponentials and subobject classifier) in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ then an ‘unraveling’ of the required form is possible.

Summarizing, we have the following metatheorem.

2.1. THEOREM. *Let I be an invariant property of toposes which is logically equivalent to the requirement that there should be a class of objects and arrows in the topos satisfying some (invariant) property P . Let \mathcal{K} be a class of sites (\mathcal{C}, J) such that*

- (i) *for any (\mathcal{C}, J) in \mathcal{K} , the topos $\mathbf{Sh}(\mathcal{C}, J)$ satisfies I if and only if there exists a class $\mathcal{A}_{(\mathcal{C}, J)}$ of objects of $\mathbf{Sh}(\mathcal{C}, J)$ of the form $a_J(S)$ and arrows $a_J(S) \rightarrow a_J(S')$ between them of the form $a_J(h)$ (where $h : S \rightarrow S'$ is an arrow in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$) which satisfies P , and*
- (ii) *there exists a class of monomorphisms m in the topos $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$, built out of objects of the form S and arrow between them through topos-theoretic constructions, such that $\mathcal{A}_{(\mathcal{C}, J)}$ satisfies P if and only if a logical condition entirely expressible in terms of the c_J -denseness of such monomorphisms m holds.*

Then I admits a bijective site characterization holding for all sites (\mathcal{C}, J) in \mathcal{K} .

PROOF. If condition (i) is satisfied then for any site (\mathcal{C}, J) in \mathcal{K} the property of the topos $\mathbf{Sh}(\mathcal{C}, J)$ to satisfy I can be expressed as a condition involving objects of the form $a_J(S)$ and arrows $a_J(S) \rightarrow a_J(S')$ between them of the form $a_J(h)$ (where $h : S \rightarrow S'$ is an arrow in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$), while by (ii) such condition can in turn be expressed as a logical condition involving the denseness of monomorphisms m in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ built out of objects of the form S and arrows between them through topos-theoretic constructions.

Now, if m is such a monomorphism then the condition that m should be c_J -dense admits an explicit formulation in terms of objects and arrows of the category \mathcal{C} as well as of the topology J ; indeed, small limits and colimits in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ admit explicit descriptions, they being computed pointwise, the subobject classifier of the topos and the exponential of two objects of the form S admit elementary descriptions in terms of the geometry of the category \mathcal{C} , and the condition of a monomorphism $A \rightarrowtail P$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ to be c_J -dense can be formulated explicitly as follows: for any $c \in \mathcal{C}$ and any $x \in P(c)$, $\{f : \text{dom}(f) \rightarrow c \mid P(f)(x) \in A(\text{dom}(f))\} \in J(c)$.

The hypotheses of the theorem precisely ensure that for any (\mathcal{C}, J) in \mathcal{K} the condition that the topos $\mathbf{Sh}(\mathcal{C}, J)$ satisfies property I can be reformulated as a logical condition entirely expressible in terms of the c_J -denseness of monomorphisms m of the above-mentioned form; it thus follows that, under these hypotheses, property I admits a bijective site characterization holding for all sites in \mathcal{K} , as required. ■

2.2. REMARKS.

- (a) The class of invariants which are logically equivalent to the requirement that there should be a class of objects and arrows in the topos satisfying some (invariant) property is very extensive. For instance, in addition to the usual topological properties of toposes, its completion up to Boolean combinations (that is, the class of invariants which are logically equivalent to a Boolean combination of invariants of this form) contains all the invariants which can be expressed in a first-order way in the language of Category Theory (or, more generally, of elementary topos theory), such as for example all the usual logical properties of toposes (to be Boolean, to be De Morgan etc.). This can be shown by an easy induction on the structure of the first-order sentence ϕ in the language of elementary topos theory which defines I ; indeed, if ϕ is of the form $(\exists x)\psi(x)$ then I can be expressed as the condition that there exists a class of objects (and arrows) in the topos satisfying the property of being a singleton whose only element satisfies property ψ , while if ϕ is of the form $(\forall x)\psi(x)$ then I can be expressed as the condition that there exists a class of objects (and arrows) in the topos satisfying the property of being the total class and that each of its elements satisfies property ψ (the case of Boolean combinations is obvious, and that of atomic sentences is equally clear since the validity of any such sentence is formally equivalent to the existence of a set of objects of the topos (vacuously) satisfying this property, since a set of objects of the topos always exists).

Also, any invariant which can be expressed as the condition that a certain specified class of objects and arrows in the topos satisfies a property P (for example, the

property of a topos to be compact, which can be expressed as the condition that the terminal object of the topos enjoys the property that any covering of it can be refined by a finite subcovering) belongs to this class of invariants; indeed, such an invariant is logically equivalent to the condition that there should exist a class of objects and arrows in the topos satisfying the joint property of being equal to the specified class and of satisfying property P .

- (b) Let I be an invariant property of toposes which is logically equivalent to the requirement that there should be a separating set of objects in the topos each of which satisfying some (invariant) property P . Then, if P descends along epimorphisms (that is, for any epimorphism in a Grothendieck topos, if its domain satisfies P then its codomain satisfies P as well), I satisfies condition (i) in the statement of Theorem 2.1 for any site (\mathcal{C}, J) . Indeed, for any site (\mathcal{C}, J) , if $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of objects satisfying property P then every object of the form $l(c)$ can be covered by a family of objects with property P , and the fact that P descends along epimorphisms ensures that we can suppose these objects to be subobjects of $l(c)$, that is objects (up to isomorphism) of the form $a_J(S)$ (for a J -closed sieve S in \mathcal{C} on c); the set $\mathcal{A}_{(\mathcal{C}, J)}$ of all such objects thus satisfies condition (i) in the statement of Theorem 2.1.
- (c) If property P can be reformulated as a logical condition entirely expressible in terms of the property that certain monomorphisms m in $\mathbf{Sh}(\mathcal{C}, J)$ built out of objects of the form $a_J(S)$ and arrows $a_J(S) \rightarrow a_J(S')$ between them of the form $a_J(h)$ (where $h : S \rightarrow S'$ is an arrow in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$) through *geometric constructions* (that is, constructions only involving finite limits or colimits in the topos $\mathbf{Sh}(\mathcal{C}, J)$) be isomorphisms then condition (ii) in the statement of Theorem 2.1 is satisfied. Indeed, finite limits and colimits are preserved by $a_J : [\mathcal{C}^{\text{op}}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}, J)$, the inclusion $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ preserves monomorphisms and for any monomorphism $m : A \rightarrow B$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is c_J -dense if and only if $a_J(m)$ is an isomorphism in $\mathbf{Sh}(\mathcal{C}, J)$.
- (d) The classical site characterizations for geometric invariants of toposes (such as the property of being localic, atomic, locally connected, equivalent to a presheaf topos, coherent etc.) of the form ‘A Grothendieck topos satisfies an invariant I if and only if *there exists* a site of definition (\mathcal{C}, J) of it satisfying a certain property $P_{(\mathcal{C}, J)}$ mentioned in section 1 can all be obtained, if the invariant can be expressed as the existence of a separating set of objects of the topos satisfying a certain property P , by considering the sites (\mathcal{C}, J) such that the set $\tilde{\mathcal{C}} := \{l(c) \mid c \in \mathcal{C}\}$ satisfies property P ; in this case, and more generally for obtaining characterizations of the kind ‘If (\mathcal{C}, J) satisfies a certain property then the topos $\mathbf{Sh}(\mathcal{C}, J)$ satisfies the given invariant’, only condition (ii) of Theorem 2.1 becomes relevant; in particular, provided that one can formulate the condition of the set $\tilde{\mathcal{C}}$ to satisfy property P as a logical condition entirely expressible in terms of the c_J -denseness of monomorphisms in $\mathbf{Sh}(\mathcal{C}, J)$ built out of objects of the form $a_J(S)$ and arrows $a_J(S) \rightarrow a_J(S')$ between them of the form $a_J(h)$

(where $h : S \rightarrow S'$ is an arrow in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$) through geometric constructions, such a characterization exists (cf. Remark 2.2(c)).

2.3. EXAMPLES.

(a) The property of a topos to be localic (respectively, to be locally connected, atomic, to have a separating set of well-supported objects) satisfies the hypotheses of Theorem 2.1. Indeed, the validity of condition (i) follows from the fact that all these invariants can be expressed in terms of the existence of a separating set of objects of the topos satisfying a property (respectively, to be subterminal, indecomposable, an atom, well-supported) which descends along epimorphisms (by Remark 2.2 - see sections 3, 4, 5, 6, 7 below for detailed proofs of these descent properties), while the fact that condition (ii) holds can be proved as follows.

- For any site (\mathcal{C}, J) , the condition for an object of the form $a_J(S)$ to be subterminal can be expressed as the requirement that the equalizer of the kernel pair of its unique arrow to the terminal object of $\mathbf{Sh}(\mathcal{C}, J)$ (which can be built from $a_J(S \rightarrow 1_{[\mathcal{C}^{\text{op}}, \mathbf{Set}]}) : a_J(S) \rightarrow a_J(1_{[\mathcal{C}^{\text{op}}, \mathbf{Set}]}) \cong 1_{\mathbf{Sh}(\mathcal{C}, J)}$ through geometric constructions) should be an isomorphism.
- For any site (\mathcal{C}, J) , the condition for an object of the form $a_J(S)$ to be indecomposable can be expressed as the requirement that any covering by pairwise disjoint subobjects (which are necessarily of the form $a_J(i) : a_J(S') \rightarrow a_J(S)$ for a canonical inclusion $i : S' \rightarrow S$) should contain an isomorphism.
- For any site (\mathcal{C}, J) , the condition for an object of the form $a_J(S)$ to be an atom can be expressed as the requirement that for any subobject $a_J(i) : a_J(S') \rightarrow a_J(S)$ of $a_J(S)$, either $a_J(i)$ is an isomorphism (that is, i is c_J -dense) or $a_J(S')$ is isomorphic to zero (that is, the canonical morphism $\emptyset \rightarrow S'$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is c_J -dense).
- For any site (\mathcal{C}, J) , the condition for an object of the form $a_J(S)$ to be well-supported can be expressed as the requirement that the morphism $a_J(S \rightarrow 1_{[\mathcal{C}^{\text{op}}, \mathbf{Set}]}) : a_J(S) \rightarrow a_J(1_{[\mathcal{C}^{\text{op}}, \mathbf{Set}]}) \cong 1_{\mathbf{Sh}(\mathcal{C}, J)}$ be epic, which is in turn equivalent to the condition that the monic part of the cover-mono factorization of the unique morphism $S \rightarrow 1_{[\mathcal{C}^{\text{op}}, \mathbf{Set}]}$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ be c_J -dense.

(b) The property of a topos to be equivalent to a presheaf topos satisfies condition (i) in the statement of Theorem 2.1 for any site, and condition (ii) for any subcanonical site. Indeed, a topos is equivalent to a presheaf topos if and only if it has a separating set of irreducible objects; condition (i) is satisfied since for any site (\mathcal{C}, J) and any irreducible object A of the topos $\mathbf{Sh}(\mathcal{C}, J)$, the fact that A is covered by objects of the form $l(c)$ implies that A is a retract of one of these objects and hence that it is, up to isomorphism, of the form $a_J(S)$ (where S is a covering sieve in \mathcal{C} , say on an object d), while if (\mathcal{C}, J) is subcanonical condition (ii) is satisfied since in order to prove that such an object $a_J(S)$ is irreducible it suffices to check that any covering

sieve on it generated by arrows having as domain objects of the form $l(c)$ contains the identity, and if (\mathcal{C}, J) is subcanonical then the condition that such an arrow $u : l(c) \rightarrow a_J(S)$ should be split epic can be formulated in terms of the existence of an arrow $s : \mathcal{C}(-, c) \rightarrow S$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ such that $a_J(s) \cong u$ and of an arrow $h : S \rightarrow \mathcal{C}(-, c)$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ such that the composite $s \circ h$ is sent by a_J to an isomorphism (notice that the condition of an arrow in a presheaf topos to be sent by a given associated sheaf functor to an isomorphism can always be reformulated in terms of the condition of certain monomorphisms to be dense, namely the equalizer of the kernel pair of the morphism and the image of the morphism).

- (c) The property of a topos to be coherent satisfies the hypotheses of Theorem 2.1 for any site $(\mathcal{C}, T_{\mathcal{C}})$, where $T_{\mathcal{C}}$ is the trivial topology on \mathcal{C} (that is, the Grothendieck topology whose only covering sieves are the maximal ones); see section 7 below.

In the following sections we shall analyze the invariants mentioned above in more detail, obtaining explicit site characterizations for them, as predicted by Theorem 2.1.

3. Localic toposes

In this section we shall address the problem of finding bijective site characterizations for the property of a topos to be localic.

Recall that a Grothendieck topos \mathcal{E} is said to be *localic* if it has a separating set of subterminal objects. We seek criteria for a topos $\mathbf{Sh}(\mathcal{C}, J)$ of sheaves on a site (\mathcal{C}, J) to be localic.

We start by observing that the objects of the form $l(c) := a_J(\mathcal{C}(-, c))$ are a separating set for $\mathbf{Sh}(\mathcal{C}, J)$; so this topos is localic if and only if for every $c \in \mathcal{C}$ the family of arrows from subterminal objects to $a_J(\mathcal{C}(-, c))$ is jointly epimorphic. Notice that the property of being a subterminal objects descends along epimorphisms, that is, if $f : a \rightarrow b$ is an epimorphism in a topos \mathcal{E} and a is a subterminal then b is a subterminal; indeed, if a is a subterminal then f is a monomorphism and hence an isomorphism.

Now, any arrow from a subterminal to $l(c)$ is monic (by definition of subterminal object) and hence, up to isomorphism, of the form $a_J(S \rightrightarrows \mathcal{C}(-, c))$ for a (J -closed) sieve S on c . On the other hand, for any sieve S on an object c , $a_J(S)$ is a subterminal object of $\mathbf{Sh}(\mathcal{C}, J)$ if and only if the equalizer of the kernel pair of its unique arrow to the terminal object of $\mathbf{Sh}(\mathcal{C}, J)$ should be an isomorphism. Now, this equalizer can be represented as $a_J(E_S)$, where $E_S \rightrightarrows S \rightrightarrows S$ is the equalizer of the two natural projections $S \times S \rightarrow S$, so it is an isomorphism if and only if the subobject $E_S \rightrightarrows S \rightrightarrows S$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is c_J -dense, i.e. for every arrows $f, g : d \rightarrow c$ in S the sieve $E_{f,g} := \{k : \text{dom}(k) \rightarrow d \mid f \circ k = g \circ k\}$ is J -covering.

Therefore, we obtain the following criterion for $\mathbf{Sh}(\mathcal{C}, J)$ to be localic.

¹This argument replaces the one given in the previous version of this paper, which worked only for subcanonical sites (\mathcal{C}, J) . The statement of Theorem 3.1 and the proofs of its two corollaries have also been modified accordingly.

3.1. THEOREM. *Let (\mathcal{C}, J) be a site. Then, with the above notation, the topos $\mathbf{Sh}(\mathcal{C}, J)$ is localic if and only if for every $c \in \mathcal{C}$ there exists a family \mathcal{F}_c of (J -closed) sieves on c in \mathcal{C} such that the sieve $\bigcup_{S \in \mathcal{F}_c} S$ is J -covering and any S in \mathcal{F}_c satisfies the property that for every arrows $f, g : d \rightarrow c$ in S the sieve $E_{f,g} := \{k : \text{dom}(k) \rightarrow d \mid f \circ k = g \circ k\}$ is J -covering.*

It is interesting to apply the theorem to presheaf toposes and to toposes of sheaves on a geometric site.

3.2. COROLLARY. *Let \mathcal{C} be a small category. Then the topos $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is localic if and only if \mathcal{C} is a preorder.*

PROOF. We can apply Theorem 3.1 by regarding $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ as the topos $\mathbf{Sh}(\mathcal{C}, J)$ where J is the trivial topology on \mathcal{C} . The theorem yields that $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is localic if and only if for any object c of \mathcal{C} , any two arrows $f, g : d \rightarrow c$ in \mathcal{C} are equal, in other words if and only if \mathcal{C} is a preorder. ■

3.3. COROLLARY. *Let $(\mathcal{C}, J_{\mathcal{C}})$ be a geometric site. Then the topos $\mathbf{Sh}(\mathcal{C}, J_{\mathcal{C}})$ is localic if and only if for every object $c \in \mathcal{C}$ there exists a covering family of arrows $\{f_i : \text{dom}(f_i) \rightarrow c \mid i \in I\}$ such that for every $i \in I$ and every object $d \in \mathcal{C}$ and every arrows $g, h : d \rightarrow \text{dom}(f_i)$, $f_i \circ g = f_i \circ h$.*

PROOF. It suffices to notice, using Theorem 3.1, that, by definition of geometric topology, the J -closed sieves are precisely those generated by a single arrow, and a family \mathcal{F}_c of sieves of the form (f_i) , where f_i is an arrow in \mathcal{C} with codomain c , satisfies the hypotheses of the theorem if and only if for every $i \in I$ and any arrows $m, n : d \rightarrow \text{dom}(f_i)$ which factor through f_i are equal (note that the J -covering sieves are generated by covering, and in particular epimorphic, families of arrows in \mathcal{C}). ■

4. Atomic toposes

In this section we shall investigate atomic toposes from the point of view of their site characterizations.

Given a topos \mathcal{E} , we recall that an object a of \mathcal{E} is said to be an *atom* of \mathcal{E} if the only subobjects of a in \mathcal{E} are the identity subobject and the zero one, and they are distinct from each other.

The following lemma, expressing the fact that atoms descend along epimorphisms, will be useful to us.

4.1. LEMMA. *Let $f : a \rightarrow b$ be an epimorphism in a topos \mathcal{E} . If a is an atom then b is an atom.*

PROOF. Let $m : b' \rightarrow b$ be a monomorphism in \mathcal{E} . Consider the pullback in \mathcal{E} of m along the arrow f .

$$\begin{array}{ccc} f^*(b') & \xrightarrow{g} & b' \\ f^*(m) \downarrow & & \downarrow m \\ a & \xrightarrow{f} & b \end{array}$$

The arrow $f^*(m) : f^*(b') \rightarrow a$ is a monomorphism (being the pullback of a monomorphism), and hence either $f^*(m) = 1_a$ or $f^*(m)$ is equal to the zero subobject of a . On the other hand, the arrow g is an epimorphism, it being the pullback of an epimorphism. So, if $f^*(m) = 1_a$ then by the uniqueness (up to isomorphism) of the epi-mono factorizations of arrows in a topos it follows that m is an isomorphism, while if $f^*(b') \cong 0$ then we have an epimorphism $g : 0 \rightarrow b'$ and hence $b' \cong 0$. ■

We shall also need the following proposition.

4.2. PROPOSITION. *Let (\mathcal{C}, J) be a site and S be a J -closed sieve on an object c of \mathcal{C} . Then $a_J(S)$ is an atom of $\mathbf{Sh}(\mathcal{C}, J)$ if and only if $\emptyset \notin J(\text{dom}(f))$ for some $f \in S$, and for every subsieve $S' \subseteq S$, either for every $f \in S' f^*(S') \in J(\text{dom}(f))$ or for every $g \in S' \emptyset \in J(\text{dom}(g))$.*

In particular, $l(c)$ is an atom if and only if for every sieve S on c , either for every arrow $f : \text{dom}(f) \rightarrow c$ in \mathcal{C}^2 , $f^(S) \in J(\text{dom}(f))$ or for every arrow $g \in S$, $\emptyset \in J(\text{dom}(g))$.*

PROOF. We start observing that for any local operator j on a topos \mathcal{E} , with associated sheaf functor $a_j : \mathcal{E} \rightarrow \mathbf{sh}_j(\mathcal{E})$, an object a of \mathcal{E} satisfies $a_j(a) \cong 0_{\mathbf{sh}_j(\mathcal{E})}$ if and only if, denoted by $a \rightarrow a' \rightarrow 1$ the epi-mono factorization of the unique arrow $a \rightarrow 1$ in \mathcal{E} , $a_j(a') \cong 0_{\mathbf{sh}_j(\mathcal{E})}$; this immediately follows from the fact that for any epimorphism in a topos its domain is isomorphic to zero if and only if its codomain is isomorphic to zero, combined with the observation that the associated sheaf functor preserves epimorphisms. Note in passing that this remark can be used to obtain explicit characterizations of the presheaves which are sent to the zero sheaf by a given associated sheaf functor.

Given a site (\mathcal{C}, J) , we would like to understand when a certain sieve S on an object $c \in \mathcal{C}$ has the property that $a_J(S) \cong 0$. We consider the epi-mono factorization $S \rightarrow I_S \rightarrow 1$ of the unique arrow $S \rightarrow \mathcal{C}(-, c) \rightarrow 1$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$. Since 1 is a J -sheaf the associated sheaf functor applied to a subterminal object U coincides with its J -closure $c_J(U)$. Therefore $a_J(S) \cong 0$ if and only if $c_J(I_S) \cong 0$. Now, I_S is clearly given by the formula $I_S = \{\text{dom}(f) \mid f \in S\}$, from which it follows, recalling that the zero subterminal in $\mathbf{Sh}(\mathcal{C}, J)$ corresponds to the ideal $\{c \in \mathcal{C} \mid \emptyset \in J(c)\}$, that $a_J(S) \cong 0$ if and only if for every $f \in S$, $\emptyset \in J(\text{dom}(f))$.

Now we want to investigate under what conditions $a_J(S)$ is an atom of $\mathbf{Sh}(\mathcal{C}, J)$. We have already characterized the conditions that make it non-zero. We observe that every subobject in $\mathbf{Sh}(\mathcal{C}, J)$ of $l(c)$ is of the form $a_J(i)$ for some inclusion of subsieves $i : S' \subseteq S$, and we have that $a_J(i)$ is an isomorphism if and only if i is c_J -dense (as a subobject in

²The condition “ $f : \text{dom}(f) \rightarrow c$ in \mathcal{C} ” incorrectly read “ $f \in S$ ” in the previous version of this paper.

$[\mathcal{C}^{\text{op}}, \mathbf{Set}]$), equivalently if $c_J(S)' \cong c_J(S)$ (or, alternatively, $S \subseteq c_J(S')$). The thesis thus follows immediately from the explicit description of the closure operator c_J . ■

4.3. REMARK. Notice that if $a_J(S)$ is an atom of the topos $\mathbf{Sh}(\mathcal{C}, J)$ and $f \in S$ is such that $\emptyset \notin J(\text{dom}(f))$ then $a_J((f)) \not\cong 0$ and hence $a_J((f)) \cong a_J(S)$. So, from the proof of the proposition we see that, to check that $a_J(S)$ is an atom it is equivalent to verify that for any arrow g such that $g^*((f)) \in J(\text{dom}(g))$ ³, either the sieve generated by it is sent by a_J to zero (equivalently, $\emptyset \in J(\text{dom}(g))$) or $f^*((g)) \in J(\text{dom}(f))$.

Now we would like to understand in concrete terms what it means for a topos $\mathbf{Sh}(\mathcal{C}, J)$ to have a separating set of atoms. First, we observe that this condition is equivalent to saying that for every $c \in \mathcal{C}$, $l(c)$ can be covered by an epimorphic family of arrows whose domains are atoms of $\mathbf{Sh}(\mathcal{C}, J)$. By Lemma 4.1, we can suppose without loss of generality (by possibly replacing any arrow in the given epimorphic family by its image), that all the arrows in the family are monic. Therefore, they are all of the form $a_J(S \rightarrow \mathcal{C}(-, c))$ for some sieve S on c . By Remark 4.3, we can suppose that S to be the J -closure of a sieve generated by a single arrow. Therefore, using Proposition 4.2, we obtain the following characterization theorem.

4.4. THEOREM. *Let (\mathcal{C}, J) be a site. Then the topos $\mathbf{Sh}(\mathcal{C}, J)$ is atomic if and only if for every $c \in \mathcal{C}$ there exists a J -covering sieve on c generated by arrows f with the property that $\emptyset \notin J(\text{dom}(f))$ and for every arrow g such that $g^*((f)) \in J(\text{dom}(g))$, either $\emptyset \in J(\text{dom}(g))$ or $f^*((g)) \in J(\text{dom}(f))$.*

5. Locally connected toposes

In this section we address the problem of establishing site characterizations for locally connected toposes.

Recall that a Grothendieck topos is locally connected if the inverse image functor of the unique geometric morphism from the topos to \mathbf{Set} has a left adjoint. Locally connected toposes can be equivalently characterized as the Grothendieck toposes which have a separating set of indecomposable objects (cf. [Caramello, 2012b]). Recall that a object of a Grothendieck topos is said to be indecomposable if it does not admit any non-trivial (set-indexed) coproduct decompositions.

Let us start with a lemma, which expresses the fact that indecomposable objects ‘descend’ along epimorphisms.

5.1. LEMMA. *Let $f : a \rightarrow b$ be an epimorphism in a Grothendieck topos \mathcal{E} . If a is indecomposable then b is indecomposable.*

³The condition “for any arrow g such that $g^*((f)) \in J(\text{dom}(g))$ ” replaces the incorrect condition “for any arrow g which factors through f ” stated in the previous version of this paper. Idem for the statement of Theorem 4.4.

PROOF. This follows immediately from the fact that if $f : a \rightarrow b$ is an epimorphism then the pullback functor $\mathcal{E}/b \rightarrow \mathcal{E}/a$ is logical and conservative (cf. [Mac Lane and Moerdijk, 1992]), in light of the fact that coproducts in a topos can be characterized as epimorphic families of pairwise disjoint subobjects. ■

Let us now turn to the problem of characterizing the indecomposable objects in a general topos $\mathbf{Sh}(\mathcal{C}, J)$.

5.2. PROPOSITION. *Let (\mathcal{C}, J) be a site. Then an object of the form $l(S)$, where S is a J -closed sieve on an object c of \mathcal{C} , is indecomposable if and only if the sieve S satisfies the property that for any family $\{S_i \mid i \in I\}$ of subsieves $S_i \subseteq S$ such that for any distinct i, i' and any $f \in S_i \cap S_{i'}$, $\emptyset \in J(\text{dom}(f))$, if the union S' of the S_i is c_J -dense in S (i.e., for any arrow $f \in S$, $f^*(\bigcup_{i \in I} S_i) \in J(\text{dom}(f))$) then some S_i is c_J -dense in S (i.e., for any arrow f in S , $f^*(S_i) \in J(\text{dom}(f))$).*

PROOF. It suffices to recall that coproducts in a topos can be characterized as epimorphic families of pairwise disjoint subobjects, and observe that, up to isomorphism, any subobject of $l(S)$ in $\mathbf{Sh}(\mathcal{C}, J)$ is, up to isomorphism, of the form $a_J(i) : a_J(T) \rightarrow a_J(S)$ where i is the canonical inclusion of a subsieve T of S into S . ■

We shall call a sieve S satisfying the property in the statement of the proposition a J -indecomposable sieve.

Using this Proposition, we can easily get a site characterization for a topos $\mathbf{Sh}(\mathcal{C}, J)$ to be locally connected.

5.3. THEOREM. *Let (\mathcal{C}, J) be a site. Then the topos $\mathbf{Sh}(\mathcal{C}, J)$ is locally connected if and only if for every $c \in \mathcal{C}$ there exists a family $\{S_i \mid i \in I\}$ of J -closed J -indecomposable sieves on c such that the union $\bigcup_{i \in I} S_i$ is J -covering.*

PROOF. The topos $\mathbf{Sh}(\mathcal{C}, J)$ is locally connected if and only if it has a separating set of indecomposable objects, equivalently for every $c \in \mathcal{C}$, the family of arrows from indecomposable objects to $l(c)$ is epimorphic. By Lemma 5.1, we can suppose without loss of generality that the arrows belonging to this family are monic, and hence are, up to isomorphism, of the form $a_J(i) : a_J(S) \rightarrow l(c) \cong a_J(\mathcal{C}(-, c))$ where i is the canonical inclusion $S \rightarrow \mathcal{C}(-, c)$ of a J -closed sieve S into $\mathcal{C}(-, c)$. Clearly a family of arrows of this form is epimorphic in $\mathbf{Sh}(\mathcal{C}, J)$ if and only if the union of the corresponding sieves is J -covering on c , from which our thesis follows. ■

6. Toposes which are equivalent to a presheaf topos

In this section we consider the invariant property of a topos to be equivalent to a presheaf topos in relation to the problem of obtaining site characterizations for it.

Let us start with a technical lemma.

6.1. LEMMA. *Let (\mathcal{C}, J) be a site and S be a J -closed sieve on an object c of \mathcal{C} . Then for any sieve R on $l(S)$ in $\mathbf{Sh}(\mathcal{C}, J)$, R is covering if and only if S is equal to the J -closure of the sieve $\{f \in S \mid l^S(f) \in R\}$ (where $l^S(f)$ denotes the factorization of $l(f)$ through the canonical monomorphism $a_J(S) \rightarrow l(c)$). In particular, a sieve R is epimorphic on $l(c)$ if and only if the sieve $\tilde{R}_S := \{f \in S \mid l(f) \in R\}$ is J -covering.*

PROOF. Given a sieve R on $l(S)$, let us consider its pullback in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ along the unit η_c of the reflection corresponding to the subtopos $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ at the object $\mathcal{C}(-, c)$. For any $f \in R$, we can cover $\eta_c^*(l(\text{dom}(f)))$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ with an epimorphic family whose domains are representable functors. By the fullness of the Yoneda Lemma, the arrow obtained by composing any such arrow $\mathcal{C}(-, d) \rightarrow \text{dom}(f)$ first with $\eta_c^*(l^S(f))$ and then with the canonical monomorphism $S \rightarrow \mathcal{C}(-, c)$ is of the form $\mathcal{C}(-, h)$ for some arrow $h : d \rightarrow c$ in \mathcal{C} . Notice that the arrows of the form $\mathcal{C}(-, h)$ corresponding to a given arrow $f \in R$ are jointly epimorphic on $\eta_c^*(l(\text{dom}(f)))$ and hence their images under the associated sheaf functor a_J yield a jointly epimorphic family T_f on $l(\text{dom}(f))$.

$$\begin{array}{ccccc}
 \eta_c^*(l(\text{dom}(f))) & \xrightarrow{\eta_c^*(l^S(f))} & S & \longrightarrow & \mathcal{C}(-, c) \\
 \downarrow & & \downarrow & & \downarrow \eta_c \\
 l(\text{dom}(f)) & \xrightarrow{l^S(f)} & l(S) & \longrightarrow & l(c)
 \end{array}$$

Now, clearly, R is epimorphic if and only if the sieve R^m on $l(S)$ obtained by ‘multi-composing’ R with the sieves T_f (for $f \in R$) is epimorphic. Notice that R^m is generated by the factorizations through the canonical monomorphism $a_J(S) \rightarrow l(c)$ of arrows of the form $l(h)$ where h is an arrow in \mathcal{C} with codomain c . Define the sieve A on c as

$$A := \{k \in S \mid l^S(k) \in R\}.$$

Clearly, the sieve A_l of arrows of the form $l^S(k)$ for $k \in A$ is contained in R and contains R^m , from which it follows that it is jointly epimorphic if and only if R (equivalently, R^m) is. From this our thesis clearly follows, since A_l is jointly epimorphic if and only if the image of the canonical monomorphism $A \rightarrow S$ under a_J is an epimorphism (equivalently, an isomorphism), that is if and only if $A \rightarrow S$ is c_J -dense, where c_J is the closure operator associated to the Grothendieck topology J . ■

We notice that the particular case of Lemma 6.1 when the sieve S is maximal was observed, but not proved, at p. 911 of [Johnstone, 2002].

Another useful remark concerning the relationship between sieves and their images under l functors is provided by the following proposition.

6.2. PROPOSITION. *Let (\mathcal{C}, J) be a site and S be a sieve on an object c of \mathcal{C} . Then the sieve*

$$\overline{S} = \{f : \text{dom}(f) \rightarrow c \mid l(f) \text{ factors through } l(g) \text{ for some } g \in S\}$$

is contained in the J -closure of S .

PROOF. Suppose that $l(f)$ factors through $l(g)$ for some $g \in S$. The canonical monomorphism $p : f^*(R) \rightarrow \mathcal{C}(-, \text{dom}(f))$ can be identified with the pullback of the canonical monomorphism $q : S \rightarrow \mathcal{C}(-, c)$ along $\mathcal{C}(-, f)$. Hence, if $l(f)$ factors through $l(g)$, $a_J(p)$ is isomorphic to the pullback of $a_J(q)$ along any factorization of $l(f)$ through $l(g)$; therefore, if $g \in S$ this latter pullback is an isomorphism and hence p is c_J -dense, in other words $f^*(R)$ is J -covering, i.e. f belongs to the J -closure of S . ■

We say that a sieve S on an object c is l -closed if $\overline{S} = S$. Note that the l -closed sieves are precisely those of the form \tilde{T} for some sieve T on c .

Notice that if J is subcanonical then every sieve is l -closed, since the functor l is full and faithful.

We can use the lemma to characterize the irreducible objects of a topos $\mathbf{Sh}(\mathcal{C}, J)$. Recall that an object of a Grothendieck topos is said to be irreducible if every covering sieve on it is maximal.

Suppose that P is an irreducible object of $\mathbf{Sh}(\mathcal{C}, J)$. We can cover P with a family of arrows whose domains are of the form $l(c)$ for $c \in \mathcal{C}$; there is thus an arrow of the family, say $e : l(c) \rightarrow P$, which is split epic, that is such that there is a monic arrow $m : P \rightarrow l(c)$ such that $e \circ m = 1_P$. Now, m being mono, P is, up to isomorphism, of the form $l(S)$ where S is a sieve on c . Now, consider the family of monomorphisms $\{l((f)) \rightarrow l(S) \mid f \in S\}$. This family covers $l(S)$ and hence, by the irreducibility of P , $l(S) \cong l((f))$ for some $f \in S$. Notice that the canonical monomorphism $(f) \rightarrow \mathcal{C}(-, c)$ is the monic part of the epi-mono factorization of the arrow $\mathcal{C}(-, f) : \mathcal{C}(-, \text{dom}(f)) \rightarrow \mathcal{C}(-, c)$ in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ and hence $m : l((f)) \rightarrow l(c)$ is the monic part of the epi-mono factorization of $l(f) : l(\text{dom}(f)) \rightarrow l(c)$ in $\mathbf{Sh}(\mathcal{C}, J)$. Let us denote by $e' : l(\text{dom}(f)) \rightarrow l((f))$ the epic part of this factorization (notice that this arrow is in fact split epic by the irreducibility of $l((f))$).

Now, given a sieve R on $l((f))$, R is maximal if and only if $f \in \tilde{R}$. Therefore the condition that for any sieve R on $l((f))$, R is epimorphic if and only if it is maximal can be equivalently expressed as the condition that \tilde{R} is dense in the c_J -closure of (f) if and only if it is maximal.

In order to obtain a criterion for irreducibility which does not involve constructions within the topos $\mathbf{Sh}(\mathcal{C}, J)$, we would like to replace the quantification over the sieves R in the topos $\mathbf{Sh}(\mathcal{C}, J)$ with a quantification over sieves in the category \mathcal{C} . To this end, we investigate whether if $l((f))$ is irreducible then it is the case that for every subsieve S of (f) - not just those of the form \tilde{R} for a sieve R on $l((f))$ - S is c_J -dense in the c_J -closure of (f) if and only if it is equal to $((f))$, and if not, whether it is possible to characterize intrinsically a class of subsieves which enjoy this property. Notice that S is c_J -dense in the c_J -closure of (f) if and only if the sieve S_p on $l((f))$ generated by the arrows of the form $l^{(f)}(g)$ for $g \in S$ is covering on $l((f))$, while, by Proposition 6.2, S_p is maximal if and only if $\overline{S} = (f)$. Therefore the required property is satisfied by all the l -closed sieves S ; that is for any such sieve S , S is c_J -dense in the c_J -closure of (f) (equivalently, $f^*(S) \in J(\text{dom}(f))$) if and only if it is equal to (f) (equivalently, $f \in S$).

Therefore we can conclude the following result.

6.3. PROPOSITION. *Let (\mathcal{C}, J) be a site and f be an arrow of \mathcal{C} . Then the object $l((f))$ is irreducible in $\mathbf{Sh}(\mathcal{C}, J)$ if and only if for every l -closed sieve $S \subseteq (f)$ on c , $f^*(S) \in J(\text{dom}(f))$ if and only if $f \in S$. The requirement of S to be l -closed can be omitted if the topology J is subcanonical.*

We now proceed to obtain, by using Proposition 6.3, an intrinsic site characterization of the toposes which are equivalent to presheaf toposes.

6.4. THEOREM. *Let (\mathcal{C}, J) be a site. Then the topos $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of irreducible objects (equivalently, is equivalent to a presheaf topos) if and only if for every object c of \mathcal{C} there exists a family $\{(k_i, w_i) \mid i \in I\}$ of pairs of composable arrows such that the sieve generated by the family $\{w_i \circ k_i \mid i \in I\}$ is J -covering and for every l -closed sieve $S \subseteq (k_i)$ on $\text{cod}(k_i)$, $k_i^*(S) \in J(\text{dom}(k_i))$ implies $k_i \in S$. The requirements of the sieves to be l -closed can be omitted if the topology J is subcanonical.*

PROOF. The topos $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of irreducible objects if and only if every object of the form $l(c)$ for an object c of \mathcal{C} is covered by a sieve generated by arrows whose domains are irreducible objects. Given any such arrow $u : P \rightarrow l(c)$ in the family (where P is an irreducible object), let us first show that there is a split epic arrow to P of the form $e : l(d) \rightarrow P$, with splitting $m : P \rightarrow l(d)$, which, composed with u , gives an arrow of the form $l(w)$. Consider the pullback $u' : P' \rightarrow \mathcal{C}(-, c)$ of u in $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ along the unit $\eta_c : \mathcal{C}(-, c) \rightarrow l(c)$; P' , regarded as an object of $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$, can be covered by an epimorphic family whose domains are representable functors. By the fullness of the Yoneda Lemma, any arrow obtained by composing such an arrow $\mathcal{C}(-, d) \rightarrow P'$ with u' is of the form $\mathcal{C}(-, w)$ for some arrow $w : d \rightarrow c$ in \mathcal{C} . Notice that the arrows of the form $\mathcal{C}(-, w)$ are jointly epimorphic on P' and hence their images under the associated sheaf functor a_J yield a jointly epimorphic family on P , which, by the irreducibility of P , must contain a split epic arrow, with monic splitting $m : P \rightarrow l(d)$. By the argument preceding Proposition 6.3, the monomorphism $m : P \rightarrow l(d)$ can be identified with the monic part of the epi-mono factorization $l(\text{dom}(k)) \twoheadrightarrow l((k)) = P \rightarrow l(d) = l(\text{cod}(k))$ of an arrow of the form $l(k)$ where k is an arrow $\text{dom}(k) \rightarrow d$ in \mathcal{C} , and $P \cong l((k))$. Let us denote by z the epic part $l(\text{dom}(k)) \twoheadrightarrow l((k))$ of this factorization. Then the arrow $u \circ z$ is equal to $l(w \circ k)$, since $u = u \circ 1_P = u \circ e \circ m \circ z = l(w) \circ m \circ z = l(w) \circ l(k) = l(w \circ k)$. Therefore, since the arrow z is epic, the family of the arrows of the form $l(w \circ k)$, where w and k vary as u does in the original epimorphic family, is epimorphic on $l(c)$, equivalently the sieve generated by the family of arrows $\{w \circ k\}$ is J -covering on c . Thus we can conclude that $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of irreducible objects if and only if for every object c of \mathcal{C} there exists a family $\{(k_i, w_i) \mid i \in I\}$ of pairs of composable arrows such that the sieve generated by the family $\{w_i \circ k_i \mid i \in I\}$ is J -covering and $l((k_i))$ is irreducible for each $i \in I$. Our thesis now follows by invoking Proposition 6.3. ■

6.5. REMARK. If \mathcal{C} is Cauchy-complete and J is subcanonical then the criterion of Theorem 6.4 significantly simplifies, as shown in [Caramello, 2012b], since all the irreducible objects in $\mathbf{Sh}(\mathcal{C}, J)$ are of the form $l(c)$ for some object $c \in \mathcal{C}$.

7. Other invariants

In this section we consider other two invariants on Grothendieck toposes, with the purpose of illustrating additional examples of invariants for which natural site characterizations can be achieved.

The first invariant that we shall investigate is the property of having a separating set of well-supported objects. Recall that an object A of a topos \mathcal{E} is said to be well-supported if the unique arrow $A \rightarrow 1$ is an epimorphism. Notice that the property of being well-supported descends along any arrow, that is for any arrow $f : A \rightarrow B$ in \mathcal{E} , if A is well-supported then B is well-supported.

Let (\mathcal{C}, J) be a site. Clearly, the topos $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of well-supported objects if and only if every object of the form $l(c)$ (for $c \in \mathcal{C}$) can be covered by a family of arrows whose domains are well-supported objects. By definition of epimorphic family, if the covering family on $l(c)$ is empty then $l(c)$ is isomorphic to zero, while if the family is non-empty then $l(c)$ is well-supported; from this remark we conclude (classically) that $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of well-supported objects if and only if for every $c \in \mathcal{C}$, either $l(c) \cong 0$ or $l(c)$ is well-supported. Now, $l(c)$ is well-supported if and only if every object of \mathcal{C} is covered by a J -covering sieve generated by arrows whose domains are objects which admit an arrow to c , while $l(c) \cong 0$ if and only if $\emptyset \in J(c)$. Therefore, we have the following result.

7.1. THEOREM. *Let (\mathcal{C}, J) be a site. Then the topos $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of well-supported objects if and only if for every $c \in \mathcal{C}$, either $\emptyset \in J(c)$ or for every $d \in \mathcal{C}$ there exists a sieve $S \in J(d)$ such that for every $f \in S$ there exists an arrow $\text{dom}(f) \rightarrow c$ in \mathcal{C} .*

We can straightforwardly apply this result to presheaf toposes and to toposes of sheaves on a geometric site.

7.2. COROLLARY. *Let \mathcal{C} be a small category. Then the topos $[\mathcal{C}^{op}, \mathbf{Set}]$ has a separating set of well-supported objects if and only if for any objects $c, d \in \mathcal{C}$ there exists an arrow $c \rightarrow d$ in \mathcal{C} .*

7.3. COROLLARY. *Let (\mathcal{C}, J) be a geometric site. Then the topos $\mathbf{Sh}(\mathcal{C}, J)$ has a separating set of well-supported objects if and only if for every $c \in \mathcal{C}$, either $c \cong 0_{\mathcal{C}}$ or the unique arrow $c \rightarrow 1_{\mathcal{C}}$ in \mathcal{C} is a cover.*

Finally, we consider a fundamental topos-theoretic invariant, namely the property of a topos to be coherent, from the point of view of site characterizations. The property of coherence for a topos is more problematic in relationship to bijective site characterization of the kind we seek than the other invariants considered above in the paper. In [Carmello, 2012b] we observed that a topos is coherent if and only if it has a separating set of compact objects which is closed under finite limits. The problem with this characterization is that we cannot suppose the separating set to be closed under quotients and therefore we cannot apply the usual technique of making the relevant property descend along epimorphisms.

On the other hand, the weaker invariant property of having a separating set of compact objects clearly admits a site characterization of the required form, since the property of an object of a topos to be compact descends along epimorphisms. Still, it is possible to achieve bijective site characterizations for the property of a topos to be coherent which hold for large classes of sites (cf. for example [Beke, 2004] for the case of presheaf toposes). For instance, by using the fact that in a presheaf topos all the representable functors are irreducible objects and that any retract of a coherent object in a coherent topos is coherent, one can immediately deduce that if $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is coherent then all the representable functors $y(c)$ are coherent objects; in particular, any finite product $y(c_1) \times \cdots \times y(c_n)$ of them is compact and the equalizer of any pair of arrows between such objects is compact (note that both conditions can be expressed as genuine properties of the category \mathcal{C} ; for example, the latter can be expressed by saying that for any arrows $f, g : c \rightarrow d$ in \mathcal{C} , the sieve consisting of all the arrows h with codomain c such that $f \circ h = g \circ h$ is generated by a finite family of arrows). Notice in passing that this kind of characterizations can be profitably applied in presence of any Morita-equivalence involving a presheaf topos according to the philosophy ‘toposes as bridges’ of [Caramello, 2010]; for example, they allow us to see that the syntactic property of a theory of presheaf type to be coherent has semantic consequences at the level of the ‘geometry’ of its category of finitely presentable models (for instance, the characterization provided by Theorem 2.1 [Beke, 2004] implies that for any theory of presheaf type \mathbb{T} , if \mathbb{T} is coherent then its category of finitely presentable models has fc finite colimits, in the sense of [Beke, 2004]).

This last example is just meant to give the reader an idea of the great amount of concrete mathematical results in distinct fields that can be obtained, in a ‘uniform’ and essentially automatic way, by using site characterizations for geometric invariants of toposes such as the ones that we have established in the present paper in connection with the philosophy ‘toposes as bridges’; other notable applications of the same general methodology can be found in [Caramello, 2012c], [Caramello, 2012b] and [Caramello, 2010].

Acknowledgements: I am very grateful to the anonymous referee for his comments and helpful suggestions, which have led to a more comprehensive and systematic presentation of the results in the paper.

References

- [Artin, Grothendieck and Verdier, 1972] M. Artin, A. Grothendieck and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas*, Séminaire de Géométrie Algébrique du Bois-Marie, année 1963-64; second edition published as Lecture Notes in Math., vols 269, 270 and 305 (Springer-Verlag, 1972).
- [Barr and Diaconescu, 1980] M. Barr and R. Diaconescu, Atomic toposes, *J. Pure Appl. Alg.* 17 (1980), 1-24.

- [Barr and Paré, 1980] M. Barr and R. Paré, Molecular toposes, *J. Pure Appl. Alg.* 17 (1980), 127-152.
- [Beke, 2004] T. Beke, Theories of presheaf type, *J. Symbolic Logic*, 69, 3 (2004), 923-934.
- [Caramello, 2010] O. Caramello, The unification of Mathematics via Topos Theory, *arXiv:math.CT/1006.3930*.
- [Caramello, 2011] O. Caramello, Atomic toposes and countable categoricity, *Applied Categorical Structures* 20 (4), 379-391 (2012).
- [Caramello, 2012a] O. Caramello, Syntactic characterizations of properties of classifying toposes, *Theory and Applications of Categories*, 26 (6), 176-193 (2012).
- [Caramello, 2012b] O. Caramello, Fraïssé's construction from a topos-theoretic perspective, *arXiv:math.CT/0805.2778v3*.
- [Johnstone, 2002] P. T. Johnstone, *Sketches of an Elephant: a topos theory compendium. Vols. 1-2*, vols. 43-44 of *Oxford Logic Guides* (Oxford University Press, 2002).
- [Mac Lane and Moerdijk, 1992] S. Mac Lane and I. Moerdijk, *Sheaves in geometry and logic: a first introduction to topos theory* (Springer-Verlag, 1992).

*Department of Pure Mathematics and Mathematical Statistics,
University of Cambridge,
Centre for Mathematical Sciences, Wilberforce Road,
Cambridge, CB3 0WB,
United Kingdom
Email: O.Caramello@dpms.cam.ac.uk*

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at [ftp://ftp.tac.mta.ca/pub/tac/html/volumes/26/25/26-25](ftp://ftp.tac.mta.ca/pub/tac/html/volumes/26/25/26-25.dvi). {dvi,ps,pdf}

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is \TeX , and use of $\text{\LaTeX}2\epsilon$ is strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

\TeX NICAL EDITOR Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT \TeX EDITOR Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS

Clemens Berger, Université de Nice-Sophia Antipolis, cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva: valeria.depaiiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne : kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh, Tom.Leinster@ed.ac.uk

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Edinburgh: Alex.Simpson@ed.ac.uk

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca