

SYNTACTIC CHARACTERIZATIONS OF PROPERTIES OF CLASSIFYING TOPOSES

OLIVIA CARMELLO

ABSTRACT. We give characterizations, for various fragments of geometric logic, of the class of theories classified by a locally connected (respectively connected and locally connected, atomic, compact, presheaf) topos, and exploit the existence of multiple sites of definition for a given topos to establish various results on quotients of theories of presheaf type.

1. Introduction

Given the fact that Grothendieck toposes are ‘the same thing as’ Morita-equivalence classes of geometric theories, it is naturally of interest to investigate how classical topos-theoretic properties of toposes translate into logical properties of the theories they classify.

Characterizations of the class of geometric theories classified by a Boolean (respectively De Morgan) topos have been provided in [Caramello, 2009a]. In section 3 of this paper, we provide syntactic characterizations, for various fragments of geometric logic, of the class of theories classified by a locally connected (respectively connected and locally connected, atomic, compact, presheaf) topos. Also, we establish criteria for a geometric theory over a given signature to be cartesian (respectively regular, coherent).

In section 4, given a quotient \mathbb{T}' of a theory of presheaf type \mathbb{T} corresponding to a Grothen-

-dieck topology J on the opposite of the category of finitely presentable \mathbb{T} -models via the duality of [Caramello, 2009c], we discuss how ‘geometrical’ properties of J translate into syntactic properties of \mathbb{T}' . In this context, we also show that, given a theory of presheaf type \mathbb{T} , the finitely presentable \mathbb{T} -models coincide with the finitely presented \mathbb{T} -models, in such a way that the opposite of the category of finitely presentable models of \mathbb{T} is equivalent to a full subcategory of the syntactic category of \mathbb{T} . This generalizes the well-known result for finitary algebraic (or more generally, cartesian) theories.

Our technique to establish these latter results is to transfer topos-theoretic invariants (that is, properties of objects of toposes which are stable under topos-theoretic equivalence) from one site of definition of a given topos to another, according to the principles introduced in [Caramello, 2010]. This paper thus represents a natural companion to [Caramello, 2009c] and [Caramello, 2009b], where subtoposes and universal models are

Received by the editors 2011-06-20 and, in revised form, 2012-02-29.

Transmitted by Jiri Rosicky. Published on 2012-03-06.

2000 Mathematics Subject Classification: 03G30, 18C10, 18B25.

Key words and phrases: Grothendieck topos, site characterizations, geometric logic, presheaf topos.

© Olivia Caramello, 2012. Permission to copy for private use granted.

investigated in particular in the context of theories of presheaf type.

The terminology used in the paper is borrowed from [Johnstone, 2002a] and [Johnstone, 2002b], if not otherwise stated. Our notion of site is that of a small Grothendieck site; anyway, all the results on sites established in the paper can be (trivially) extended to essentially small sites of the form (\mathcal{C}, J) where \mathcal{C} is an essentially small category.

2. Some geometric invariants

Let us start with some general facts about dense subcategories. We recall from [Johnstone, 2002b] the following definition.

2.1. DEFINITION. *Let (\mathcal{C}, J) be a site. We say a subcategory \mathcal{D} of \mathcal{C} is J -dense if*

- (a) *every object c of \mathcal{C} has a covering sieve $R \in J(c)$ generated by morphisms whose domains are in \mathcal{D} ; and*
- (b) *for any morphism $f : c \rightarrow d$ in \mathcal{C} with $d \in \mathcal{D}$, there is a covering sieve $R \in J(c)$ generated by morphisms $g : b \rightarrow c$ for which the composite $f \circ g$ is in \mathcal{D} .*

Let us denote by $\mathbf{Groth}(\mathcal{C})$ the Heyting algebra of Grothendieck topologies on a category \mathcal{C} (cfr. [Caramello, 2009c]). Given a subcategory \mathcal{C}' of \mathcal{C} , we denote by $\mathbf{Groth}_{\mathcal{C}'}(\mathcal{C})$ the subset of $\mathbf{Groth}(\mathcal{C})$ formed by the Grothendieck topologies J on \mathcal{C} such that \mathcal{C}' is J -dense.

There is an obvious notion of intersection of subcategories; specifically, given a collection $\{\mathcal{C}_i \mid i \in I\}$ of subcategories of a category \mathcal{C} , we can define their intersection \mathcal{C}' by putting $ob(\mathcal{C}') = \bigcap_{i \in I} ob(\mathcal{C}_i)$ and $arr(\mathcal{C}') = \bigcap_{i \in I} arr(\mathcal{C}_i)$, that is, given an arrow $f : a \rightarrow b$ in \mathcal{C} , f belongs to $arr(\mathcal{C}')$ if and only if it belongs to $arr(\mathcal{C}_i)$ for every $i \in I$. It is immediate to see that \mathcal{C}' is a subcategory of \mathcal{C} .

We note that, for any Grothendieck topology J on \mathcal{C} , any finite intersection of subcategories of \mathcal{C} which are J -dense is again J -dense; indeed, this easily follows from the fact that a finite intersection of J -covering sieves is again J -covering.

The following result provides a couple of useful facts about dense subcategories.

2.2. PROPOSITION. *Let \mathcal{C} be a category. Then*

- (i) *$\mathbf{Groth}_{\mathcal{C}'}(\mathcal{C})$ is closed in $\mathbf{Groth}(\mathcal{C})$ under arbitrary (non-empty) intersections (that is, meets in $\mathbf{Groth}(\mathcal{C})$) and under taking larger topologies, and hence it is an Heyting algebra inheriting the Heyting operations from $\mathbf{Groth}(\mathcal{C})$ whose maximal element is the maximal Grothendieck topology on \mathcal{C} and minimal element is the intersection of all the topologies in $\mathbf{Groth}_{\mathcal{C}'}(\mathcal{C})$;*
- (ii) *Let \mathcal{C}' and \mathcal{C}'' subcategories of \mathcal{C} such that \mathcal{C}' is a subcategory of \mathcal{C}'' . Then \mathcal{C}' is J -dense if and only if \mathcal{C}' is $J|_{\mathcal{C}''}$ -dense (as a subcategory of \mathcal{C}'') and \mathcal{C}'' is J -dense.*

PROOF. (i) The first assertion is obvious while the second easily follows from the fact that arbitrary (non-empty) unions of J -covering sieves are J -covering.

(ii) Suppose that \mathcal{C}' is J -dense. Then, by part (i) of the proposition, \mathcal{C}'' is J -dense, and it is immediate to see that \mathcal{C}' is $J|_{\mathcal{C}''}$ -dense (as a subcategory of \mathcal{C}''). The converse follows from the fact that ‘composition’ of J -covering sieves (in the sense of Definition 2.3 [Caramello, 2009c]) is J -covering. ■

We note that a small full subcategory of a Grothendieck topos \mathcal{E} is $J_{\mathcal{E}}$ -dense, where $J_{\mathcal{E}}$ is the canonical Grothendieck topology on \mathcal{E} , if and only if it is a separating set for \mathcal{E} .

Let us also recall the following notions.

2.3. DEFINITION. *Let \mathcal{E} be a Grothendieck topos and A an object of \mathcal{E} . Then*

- (a) *A is said to be an atom if the only subobjects of A in \mathcal{E} are the identity on A and the zero subobject, and they are distinct from each other;*
- (b) *A is said to be indecomposable if does not admit any non-trivial coproduct decompositions;*
- (c) *A is said to be irreducible if it is $J_{\mathcal{E}}$ -irreducible, where $J_{\mathcal{E}}$ is the canonical topology on \mathcal{E} ; in other words, if any sieve in \mathcal{E} containing a small epimorphic family contains the identity on A ;*
- (d) *A is said to be compact if every small covering family $\{A_i \rightarrow A \mid i \in I\}$ contains a finite covering subfamily;*
- (e) *A is said to be coherent if it is compact and, whenever we are given a morphism $f : B \rightarrow A$ with B compact, the domain of the kernel-pair of f is compact;*
- (f) *A is said to be supercompact if every small covering family $\{A_i \rightarrow A \mid i \in I\}$ contains a cover;*
- (g) *A is said to be regular if it is supercompact and, whenever we are given a morphism $f : B \rightarrow A$ with B supercompact, the domain of the kernel-pair of f is supercompact.*

Recall that an object in a locally connected topos is indecomposable if and only if it is connected (cfr. the discussion after the proof of Lemma C3.3.6 [Johnstone, 2002b]).

2.4. REMARK. It readily follows from the definitions that every coherent (respectively regular) object is compact (respectively supercompact), every atom is an indecomposable object, every irreducible object is supercompact, every supercompact object is indecomposable.

Let us recall from [Johnstone, 2002b] the following terminology.

A site (\mathcal{C}, J) is said to be locally connected if every J -covering sieve is connected, that is, for any $R \in J(c)$, R is connected as a full subcategory of \mathcal{C}/c .

A site (\mathcal{C}, J) is said to be atomic if \mathcal{C} satisfies the right Ore condition and J is the atomic topology on \mathcal{C} .

Given a site (\mathcal{C}, J) ,

- (a) we say an object c of \mathcal{C} is J -irreducible if the only J -covering sieve on c is the maximal sieve $M(c)$;
- (b) we say J is rigid if, for every $c \in \mathcal{C}$, the family of all morphisms from J -irreducible objects to c generates a J -covering sieve.

A site (\mathcal{C}, J) is said to be rigid if J is rigid as a Grothendieck topology on \mathcal{C} .

We will say that a site (\mathcal{C}, J) is coherent (respectively regular) if \mathcal{C} is cartesian and J is a finite-type Grothendieck topology on \mathcal{C} (respectively a Grothendieck topology J such that every J -covering sieve is generated by a single arrow).

Given a site (\mathcal{C}, J) , we denote by $l_J^{\mathcal{C}} : \mathcal{C} \rightarrow \mathbf{Sh}(\mathcal{C}, J)$ the composite of the Yoneda embedding $\mathcal{C} \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ with the associated sheaf functor $a_J : [\mathcal{C}^{\text{op}}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}, J)$. We have the following result.

2.5. PROPOSITION. *Let (\mathcal{C}, J) be a site. Then*

- (i) *if (\mathcal{C}, J) is locally connected then for each $c \in \mathcal{C}$, $l_J^{\mathcal{C}}(c)$ is an indecomposable (equivalently, connected) object in $\mathbf{Sh}(\mathcal{C}, J)$;*
- (ii) *if (\mathcal{C}, J) is atomic then for each $c \in \mathcal{C}$, $l_J^{\mathcal{C}}(c)$ is an atom in $\mathbf{Sh}(\mathcal{C}, J)$;*
- (iii) *if (\mathcal{C}, J) is rigid then for each $c \in \mathcal{C}$ such that c is J -irreducible, $l_J^{\mathcal{C}}(c)$ is an indecomposable projective (equivalently, an irreducible object) of $\mathbf{Sh}(\mathcal{C}, J)$;*
- (iv) *if (\mathcal{C}, J) is coherent (respectively regular) then for each $c \in \mathcal{C}$, $l_J^{\mathcal{C}}(c)$ is a coherent (respectively regular) object of $\mathbf{Sh}(\mathcal{C}, J)$.*

PROOF. (i) and (ii) were proved in [Caramello, 2011].

To prove (iii), we note that if (\mathcal{C}, J) is rigid then the Comparison Lemma yields $\mathbf{Sh}(\mathcal{C}, J) \simeq [\mathcal{D}^{\text{op}}, \mathbf{Set}]$ where \mathcal{D} is the full subcategory of \mathcal{C} on the J -irreducible objects (cfr. the discussion after Definition C2.2.18 [Johnstone, 2002b]) and, under this equivalence, for each $c \in \mathcal{D}$, $l_J^{\mathcal{C}}(c)$ corresponds to the representable $y(c) : \mathcal{D}^{\text{op}} \rightarrow \mathbf{Set}$. Now, it is well-known that all the representables on \mathcal{D} are indecomposable projective objects in $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$, from which our thesis follows.

Part (iv) was proved in [Johnstone, 2002b] (cfr. Theorem D3.3.7 and Remark D3.3.10). ■

We note that if \mathcal{E} is equivalent to a presheaf topos then any object of \mathcal{E} is irreducible if and only if it is indecomposable and projective. Indeed, by the argument in the proof of Lemma C2.2.20 [Johnstone, 2002b], any indecomposable projective object in \mathcal{E} is irreducible. Conversely, if \mathcal{G} is the full subcategory of \mathcal{E} on the irreducible objects then, by Lemma C2.2.20, \mathcal{G} is essentially small and $(\mathcal{G}, J_{\mathcal{E}|_{\mathcal{G}}})$ is a rigid site; moreover, by the Comparison Lemma $\mathbf{Sh}(\mathcal{E}, J_{\mathcal{E}|_{\mathcal{G}}}) \simeq \mathbf{Sh}(\mathcal{G}, J_{\mathcal{E}|_{\mathcal{G}}})$ and hence Proposition 2.5(iii) implies our thesis.

Let us recall from [Johnstone, 2002b] (C2.3.2(c)) that, for any essentially small site (\mathcal{C}, J) , a sieve R on $l_J^{\mathcal{C}}(c)$ in $\mathbf{Sh}(\mathcal{C}, J)$ is epimorphic iff the sieve $\{f : d \rightarrow c \mid l_J^{\mathcal{C}}(f) \in R\}$

is J -covering in \mathcal{C} . This fact enables us to express properties of objects of the form $l_J^{\mathcal{C}}(c)$ like compactness, supercompactness or irreducibility in terms of properties of J -covering sieves on c , as follows (point (i) of the following proposition was proved in [Johnstone, 2002b] as Lemma D3.3.4).

2.6. PROPOSITION. *Let (\mathcal{C}, J) be a site. Then, with the notation above, we have:*

- (i) $l_J^{\mathcal{C}}(c)$ is compact if and only if every J -covering sieve on c contains a finite family of arrows which generates a J -covering sieve;
- (ii) $l_J^{\mathcal{C}}(c)$ is supercompact if and only if every J -covering sieve on c contains a single arrow which generates a J -covering sieve;
- (iii) $l_J^{\mathcal{C}}(c)$ is irreducible if and only if every J -covering sieve on c is maximal, that is, c is J -irreducible.

It turns out that one can rephrase many interesting properties of Grothendieck toposes in terms of the existence of separating sets for them with particular properties. For example, it is well-known (cfr. Lemma C2.2.20 [Johnstone, 2002b]) that a Grothendieck topos is equivalent to a presheaf topos if and only if it has a separating set of indecomposable projective objects; moreover, we have the following characterizations.

2.7. THEOREM. *Let \mathcal{E} be a Grothendieck topos. Then*

- (i) \mathcal{E} is locally connected if and only if it has a separating set of indecomposable objects;
- (ii) \mathcal{E} is connected and locally connected if and only if it has a separating set of indecomposable objects containing the terminal object of \mathcal{E} ;
- (iii) \mathcal{E} is atomic if and only if it has a separating set of atoms;
- (iv) \mathcal{E} is coherent (respectively regular) if and only if it has a separating set of coherent (respectively regular) objects which is closed in \mathcal{E} under finite limits.

PROOF. (i) Suppose that \mathcal{E} is locally connected; then, by Theorem C3.3.10 [Johnstone, 2002b], \mathcal{E} is of the form $\mathbf{Sh}(\mathcal{C}, J)$ for a locally connected small site (\mathcal{C}, J) . Then the objects of the form $l_J^{\mathcal{C}}(c)$ for $c \in \mathcal{C}$ are indecomposable (by Proposition 2.5(i)) and hence they form a separating set for \mathcal{E} . Conversely, suppose that \mathcal{E} has a separating set of indecomposable objects; then, by arguing as in the proof of Theorem C3.3.10 [Johnstone, 2002b], we obtain that the full subcategory \mathcal{I} of \mathcal{E} on the indecomposable objects, equipped with the Grothendieck topology $J_{\mathcal{E}}|_{\mathcal{I}}$ on \mathcal{I} induced by the canonical coverage on \mathcal{E} is a locally connected site; but, by the Comparison Lemma, \mathcal{E} is equivalent to $\mathbf{Sh}(\mathcal{I}, J_{\mathcal{E}}|_{\mathcal{I}})$, so that the thesis follows from Theorem C3.3.10 [Johnstone, 2002b].

(ii) This follows analogously to part (i), by using the ‘connected and locally connected’ version of Theorem C3.3.10 [Johnstone, 2002b].

(iii) This was already proved in [Carmello, 2011] (Proposition 1.3(i)).

(iv) One direction follows from Theorem D3.3.7 [Johnstone, 2002b] and Remarks D3.3.9 and D3.3.10 [Johnstone, 2002b]. Conversely, suppose that \mathcal{E} has a separating set \mathcal{G} of coherent (respectively regular) objects which is closed in \mathcal{E} under finite limits. Then \mathcal{G} is a cartesian category and, by the Comparison Lemma, \mathcal{E} is equivalent to $\mathbf{Sh}(\mathcal{G}, J_{\mathcal{E}}|_{\mathcal{G}})$; now, by Proposition 2.6(i) (respectively Proposition 2.6(ii)), the site $(\mathcal{G}, J_{\mathcal{E}}|_{\mathcal{G}})$ is coherent (respectively regular), and hence, by Theorem D3.3.1 [Johnstone, 2002b], \mathcal{E} is a coherent (respectively regular) topos, as required. ■

2.8. **REMARK.** Notice that it follows from the theorem and Remark 2.4 that any presheaf topos is locally connected, any regular topos is locally connected and any atomic topos is locally connected. Moreover, it is clear that if \mathcal{E} is a Boolean topos then for any object A of \mathcal{E} , A is an atom if and only if it is non-zero and indecomposable, from which it follows that any Boolean locally connected topos is atomic.

2.9. **REMARK.** We note that the theorem implies that, given a site (\mathcal{C}, J) , if all the objects of the form $l_J^{\mathcal{C}}(c)$ for $c \in \mathcal{C}$ are indecomposable objects (respectively atoms) of $\mathbf{Sh}(\mathcal{C}, J)$ then the topos $\mathbf{Sh}(\mathcal{C}, J)$ is locally connected (respectively atomic). Also, provided that \mathcal{C} is cartesian, if all the objects of the form $l_J^{\mathcal{C}}(c)$ for $c \in \mathcal{C}$ are regular (respectively coherent) in $\mathbf{Sh}(\mathcal{C}, J)$ then the topos $\mathbf{Sh}(\mathcal{C}, J)$ is regular (respectively coherent). An application of this remark in the context of quotients of theories of presheaf type will be provided in section 4 below.

3. Syntactic criteria

Let \mathbb{T} be a cartesian (respectively regular, coherent, geometric) theory over a signature Σ . We denote by $\mathcal{C}_{\mathbb{T}}^{\text{cart}}$ (respectively $\mathcal{C}_{\mathbb{T}}^{\text{reg}}$, $\mathcal{C}_{\mathbb{T}}^{\text{coh}}$, $\mathcal{C}_{\mathbb{T}}^{\text{geom}}$) the cartesian (respectively regular, coherent, geometric) syntactic category of \mathbb{T} and by $J_{\mathbb{T}}^{\text{reg}}$ the regular (respectively coherent, geometric) topology on $\mathcal{C}_{\mathbb{T}}^{\text{reg}}$ (respectively $\mathcal{C}_{\mathbb{T}}^{\text{coh}}$, $\mathcal{C}_{\mathbb{T}}^{\text{geom}}$). Recall from [Johnstone, 2002b] that if \mathbb{T} is cartesian (respectively regular, coherent, geometric) then $[\mathcal{C}_{\mathbb{T}}^{\text{cart op}}, \mathbf{Set}]$ (respectively $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{reg}}, J_{\mathbb{T}}^{\text{reg}})$, $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{coh}}, J_{\mathbb{T}}^{\text{coh}})$, $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$) is a classifying topos for \mathbb{T} . Let us denote by $y_{\mathbb{T}}^{\text{cart}} : \mathcal{C}_{\mathbb{T}}^{\text{cart}} \rightarrow [\mathcal{C}_{\mathbb{T}}^{\text{cart op}}, \mathbf{Set}]$ (respectively $y_{\mathbb{T}}^{\text{reg}} : \mathcal{C}_{\mathbb{T}}^{\text{reg}} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{reg}}, J_{\mathbb{T}}^{\text{reg}})$, $y_{\mathbb{T}}^{\text{coh}} : \mathcal{C}_{\mathbb{T}}^{\text{coh}} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{coh}}, J_{\mathbb{T}}^{\text{coh}})$, $y_{\mathbb{T}}^{\text{geom}} : \mathcal{C}_{\mathbb{T}}^{\text{geom}} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$) the Yoneda embeddings.

Let us introduce the following notions. Below, by a \mathbb{T} -provably functional geometric formula from $\{\vec{x}.\phi\}$ to $\{\vec{y}.\psi\}$ we mean a geometric formula $\theta(\vec{x}, \vec{y})$ such that the sequents $(\phi \vdash_{\vec{x}} (\exists \vec{y})\theta)$, $(\theta \vdash_{\vec{x}, \vec{y}} \phi \wedge \psi)$ and $(\theta \wedge \theta[\vec{z}/\vec{y}] \vdash_{\vec{x}, \vec{y}, \vec{z}} \vec{y} = \vec{z})$ are provable in \mathbb{T} .

3.1. **DEFINITION.** *Let \mathbb{T} be a geometric theory over a signature Σ and $\phi(\vec{x})$ a geometric formula-in-context over Σ . Then*

- (a) *we say that $\phi(\vec{x})$ is \mathbb{T} -complete if the sequent $(\phi \vdash_{\vec{x}} \perp)$ is not provable in \mathbb{T} , and for every geometric formula ϕ in the same context either $(\chi \vdash_{\vec{x}} \perp)$ or $(\chi \wedge \phi \vdash_{\vec{x}} \perp)$ is provable in \mathbb{T} ;*
- (b) *we say that $\phi(\vec{x})$ is \mathbb{T} -indecomposable if for any family $\{\psi_i(\vec{x}) \mid i \in I\}$ of geometric formulae in the same context such that for each i , ψ_i \mathbb{T} -provably implies ϕ and for*

any distinct $i, j \in I$, $\psi_i \wedge \psi_j \vdash_{\vec{x}} \perp$ is provable in \mathbb{T} , we have that $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ provable in \mathbb{T} implies $\phi \vdash_{\vec{x}} \psi_i$ provable in \mathbb{T} for some $i \in I$;

- (c) we say that $\phi(\vec{x})$ is \mathbb{T} -irreducible if for any family $\{\theta_i \mid i \in I\}$ of \mathbb{T} -provably functional geometric formulae $\{\vec{x}_i, \vec{x}.\theta_i\}$ from $\{\vec{x}_i.\phi_i\}$ to $\{\vec{x}.\phi\}$ such that $\phi \vdash_{\vec{x}} \bigvee_{i \in I} (\exists \vec{x}_i)\theta_i$ is provable in \mathbb{T} , there exist $i \in I$ and a \mathbb{T} -provably functional geometric formula $\{\vec{x}, \vec{x}_i.\theta'\}$ from $\{\vec{x}.\phi\}$ to $\{\vec{x}_i.\phi_i\}$ such that $\phi \vdash_{\vec{x}} (\exists \vec{x}_i)(\theta' \wedge \theta_i)$ is provable in \mathbb{T} ;
- (d) we say that $\phi(\vec{x})$ is \mathbb{T} -compact if for any family $\{\psi_i(\vec{x}) \mid i \in I\}$ of geometric formulae in the same context, $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ provable in \mathbb{T} implies $\phi \vdash_{\vec{x}} \bigvee_{i \in I'} \psi_i$ provable in \mathbb{T} for some finite subset I' of I ;
- (e) we say that $\phi(\vec{x})$ is \mathbb{T} -supercompact if for any family $\{\psi_i(\vec{x}) \mid i \in I\}$ of geometric formulae in the same context, $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ provable in \mathbb{T} implies $\phi \vdash_{\vec{x}} \psi_i$ provable in \mathbb{T} for some $i \in I$.

3.2. LEMMA. *Let \mathbb{T} be a geometric theory over a signature Σ and $\phi(\vec{x})$ a geometric formula-in-context over Σ . Then*

- (i) $\phi(\vec{x})$ is \mathbb{T} -complete if and only if $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is an atom of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$;
- (ii) $\phi(\vec{x})$ is \mathbb{T} -indecomposable if and only if $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is an indecomposable object of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$;
- (iii) $\phi(\vec{x})$ is \mathbb{T} -irreducible if and only if $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is an irreducible object of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$;
- (iv) $\phi(\vec{x})$ is \mathbb{T} -compact if and only if $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is a compact object of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$;
- (v) $\phi(\vec{x})$ is \mathbb{T} -supercompact if and only if $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is a supercompact object of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$.

PROOF. Parts (iii), (iv) and (v) follow immediately from Proposition 2.6, by using Lemma D1.4.4(iv) [Johnstone, 2002b] and the fact that cover-mono factorizations of arrow exist in $\mathcal{C}_{\mathbb{T}}^{\text{geom}}$.

Parts (i) and (ii) follow from the fact that in a topos coproducts are the same thing as disjoint unions, since $y_{\mathbb{T}}^{\text{geom}}(\mathcal{C}_{\mathbb{T}}^{\text{geom}})$ is closed in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ under taking subobjects and $y_{\mathbb{T}}^{\text{geom}} : \mathcal{C}_{\mathbb{T}}^{\text{geom}} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ is a full and faithful geometric functor. \blacksquare

Recall from [Johnstone, 2002b] that a Grothendieck topos \mathcal{E} is compact if and only if the terminal object of \mathcal{E} is compact; hence, a geometric theory \mathbb{T} over a signature Σ is classified by a compact topos if and only if $y_{\mathbb{T}}^{\text{geom}}(\{\llbracket \cdot \rrbracket . \top\})$ is compact in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$, if and only if for any family $\{\psi_i \mid i \in I\}$ of geometric sentences, $\phi \vdash \bigvee_{i \in I} \psi_i$ provable in \mathbb{T} implies $\phi \vdash \bigvee_{i \in I'} \psi_i$ provable in \mathbb{T} for some finite subset I' of I .

We note that any cartesian (respectively regular, coherent) theory \mathbb{T} over a signature Σ can be regarded as a geometric theory, and hence we have an equivalence of classifying toposes

$$[\mathcal{C}_{\mathbb{T}}^{\text{cart}^{\text{op}}}, \mathbf{Set}] \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$$

(respectively

$$\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{reg}}, J_{\mathbb{T}}^{\text{reg}}) \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}}),$$

$$\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{coh}}, J_{\mathbb{T}}^{\text{coh}}) \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})).$$

Moreover, it is immediate to see that for any cartesian (respectively regular, coherent) formula $\phi(\vec{x})$ over Σ , $y_{\mathbb{T}}^{\text{cart}}(\{\vec{x}.\phi\})$ (respectively $y_{\mathbb{T}}^{\text{reg}}(\{\vec{x}.\phi\})$, $y_{\mathbb{T}}^{\text{coh}}(\{\vec{x}.\phi\})$) corresponds under the equivalence to $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$.

This remark, combined with Propositions 2.5 and 2.6, leads to the following results. Below, given a geometric theory \mathbb{T} over a signature Σ , by saying that \mathbb{T} is equivalent to a cartesian (respectively regular, coherent) theory we mean that \mathbb{T} can be axiomatized by cartesian (respectively regular, coherent) sequents over Σ .

3.3. THEOREM. *Let \mathbb{T} be a geometric theory over a signature Σ . Then \mathbb{T} is equivalent to a cartesian theory if and only if for any cartesian formula $\{\vec{x}.\phi\}$ over Σ , for any family $\{\theta_i \mid i \in I\}$ of \mathbb{T} -provably functional geometric formulae $\{\vec{x}_i, \vec{x}.\theta_i\}$ from $\{\vec{x}_i.\phi_i\}$ to $\{\vec{x}.\phi\}$ such that $\phi \vdash_{\vec{x}} \bigvee_{i \in I} (\exists \vec{x}_i)\theta_i$ is provable in \mathbb{T} , there exist $i \in I$ and a \mathbb{T} -provably functional geometric formula $\{\vec{x}, \vec{x}_i.\theta'\}$ from $\{\vec{x}.\phi\}$ to $\{\vec{x}_i.\phi_i\}$ such that $\phi \vdash_{\vec{x}} (\exists \vec{x}_i)(\theta' \wedge \theta_i)$ is provable in \mathbb{T} .*

PROOF. Let us suppose that \mathbb{T} is cartesian. Then the property of $\{\vec{x}.\phi\}$ in the statement of the proposition is equivalent, by Lemma 3.2(iii), to saying that $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is irreducible in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$. But this condition corresponds, under the equivalence

$$[\mathcal{C}_{\mathbb{T}}^{\text{cart}^{\text{op}}}, \mathbf{Set}] \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}}),$$

to saying that $y_{\mathbb{T}}^{\text{cart}}(\{\vec{x}.\phi\})$ is irreducible (equivalently, indecomposable projective) in the topos $[\mathcal{C}_{\mathbb{T}}^{\text{cart}^{\text{op}}}, \mathbf{Set}]$, and this true (cfr. Proposition 2.5(iii)).

Conversely, if \mathbb{T} is geometric and for any cartesian formula $\phi(\vec{x})$ over Σ , $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is irreducible in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ then, denoted by \mathcal{G} the full subcategory of $\mathcal{C}_{\mathbb{T}}^{\text{geom}}$ on the cartesian formulae, equivalently the cartesian syntactic category $\mathcal{C}_{\mathbb{T}'}^{\text{cart}}$ of the cartesianization \mathbb{T}' of \mathbb{T} (that is, the theory axiomatized by all the cartesian sequents over Σ which are provable in \mathbb{T}), we have that \mathcal{G} is $J_{\mathbb{T}}^{\text{geom}}$ -dense (by Lemma D1.3.8 [Johnstone, 2002b])

and $J_{\mathbb{T}}^{\text{geom}}|_{\mathcal{G}}$ is the trivial Grothendieck topology. Thus the Comparison Lemma yields an equivalence

$$\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}}) \simeq [\mathcal{C}_{\mathbb{T}'}^{\text{cartop}}, \mathbf{Set}].$$

Clearly, this equivalence sends the universal model of \mathbb{T} in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ to the universal model of \mathbb{T}' in $[\mathcal{C}_{\mathbb{T}'}^{\text{cartop}}, \mathbf{Set}]$, and hence, universal models being conservative, \mathbb{T} and \mathbb{T}' prove exactly the same geometric sequents over Σ , that is, they are equivalent, as required. ■

The ‘only if’ direction in the following result extends Lemma D3.3.11 [Johnstone, 2002b].

3.4. THEOREM. *Let \mathbb{T} be a geometric theory over a signature Σ . Then \mathbb{T} is equivalent to a regular theory if and only if for any regular formula $\{\vec{x}.\phi\}$ over Σ , for any family $\{\psi_i(\vec{x}) \mid i \in I\}$ of geometric formulae in the same context, $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ provable in \mathbb{T} implies $\phi \vdash_{\vec{x}} \psi_i$ provable in \mathbb{T} for some $i \in I$.*

PROOF. This follows similarly to Theorem 3.3 by using Lemma 3.2(v) and Proposition 2.5(iv). ■

3.5. THEOREM. *Let \mathbb{T} be a geometric theory over a signature Σ . Then \mathbb{T} is equivalent to a coherent theory if and only if for any coherent formula $\{\vec{x}.\phi\}$ over Σ , for any family $\{\psi_i(\vec{x}) \mid i \in I\}$ of geometric formulae in the same context, $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ provable in \mathbb{T} implies $\phi \vdash_{\vec{x}} \bigvee_{i \in I'} \psi_i$ provable in \mathbb{T} for some finite subset I' of I .*

PROOF. This follows similarly to Theorem 3.3 by using Lemma 3.2(iv) and Proposition 2.5(iv). ■

3.6. LOCALLY CONNECTED THEORIES The following result gives a syntactic characterization of the class of geometric theories classified by a locally connected (respectively connected and locally connected) topos.

3.7. THEOREM. *Let \mathbb{T} be a geometric theory over a signature Σ . Then \mathbb{T} is classified by a locally connected topos (respectively connected and locally connected topos) if and only if for any geometric formula $\phi(\vec{x})$ over Σ there exists a (unique) family $\{\psi_i(\vec{x}) \mid i \in I\}$ of \mathbb{T} -indecomposable geometric formulae in the same context (respectively, containing $\{[], \top\}$) such that*

- (i) for each i , ψ_i \mathbb{T} -provably implies ϕ ,
- (ii) for any distinct $i, j \in I$, $\psi_i \wedge \psi_j \vdash_{\vec{x}} \perp$ is provable in \mathbb{T} , and
- (iii) $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ is provable in \mathbb{T} .

PROOF. Let us suppose that the classifying topos $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ of \mathbb{T} is locally connected. Then, by Lemma D3.3.6 [Johnstone, 2002b], given a geometric formula $\phi(\vec{x})$ over Σ , the object $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ of the topos $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ is uniquely expressible as a coproduct of indecomposable objects. Now, since $y_{\mathbb{T}}^{\text{geom}}(\mathcal{C}_{\mathbb{T}}^{\text{geom}})$ is closed in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ under taking subobjects (the topos $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ being the ∞ -pretopos completion of $\mathcal{C}_{\mathbb{T}}^{\text{geom}}$), we can suppose that all the subobjects of $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$, and in particular the indecomposable objects arising in our coproduct, are of the form $y_{\mathbb{T}}^{\text{geom}}(c)$ for some $c \in \mathcal{C}_{\mathbb{T}}^{\text{geom}}$. The condition of the criterion then follows from Lemma 3.2(ii) and the fact that the functor $y_{\mathbb{T}}^{\text{geom}}$ is geometric and full and faithful.

Conversely, if the condition of the criterion is satisfied then we have, by Lemma 3.2(ii) and Proposition 2.2(ii), that the objects of the form $y_{\mathbb{T}}^{\text{geom}}(\{\vec{y}.\psi\})$ for a \mathbb{T} -indecomposable formula $\psi(\vec{y})$ form a separating set for $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ made of indecomposable objects; then $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ is locally connected by Theorem 2.7(i). ■

The following result is the coherent analogue of this theorem.

3.8. THEOREM. *Let \mathbb{T} be a coherent theory over a signature Σ . Then \mathbb{T} is classified by a locally connected topos (respectively connected and locally connected topos) if and only if for any coherent formula $\phi(\vec{x})$ over Σ there exists a (unique) finite family $\{\psi_i(\vec{x}) \mid i \in I\}$ of \mathbb{T} -indecomposable geometric formulae in the same context (resp. containing $\{\perp, \top\}$) such that*

- (i) for each i , ψ_i \mathbb{T} -provably implies ϕ ,
- (ii) for any distinct $i, j \in I$, $\psi_i \wedge \psi_j \vdash_{\vec{x}} \perp$ is provable in \mathbb{T} , and
- (iii) $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ is provable in \mathbb{T} .

PROOF. The proof proceeds analogously to the proof of Theorem 3.7, by using Theorem 3.5. ■

3.9. REMARK. From the proof of the theorems it is clear that, by using the notion of dense subcategory, one can obtain alternative (although equivalent) versions of the criteria. For example, a weaker (in the ‘if’ direction) version of the criterion of Theorem 3.7 reads as follows: a geometric theory \mathbb{T} is classified by a locally connected topos (respectively connected and locally connected topos) if and only if there exists a collection \mathcal{F} (respectively, a collection \mathcal{F} containing $\{\perp, \top\}$) of \mathbb{T} -indecomposable geometric formulae-in-context over Σ such that for any geometric formula $\{\vec{y}.\psi\}$ over Σ , there exist objects $\{\vec{x}_i.\phi_i\}$ in \mathcal{F} as i varies in I and \mathbb{T} -provably functional geometric formulae $\{\vec{x}_i, \vec{y}.\theta_i\}$ from $\{\vec{x}_i.\phi_i\}$ to $\{\vec{y}.\psi\}$ such that $\psi \vdash_{\vec{y}} \bigvee_{i \in I} (\exists \vec{x}_i)\theta_i$ is provable in \mathbb{T} . Naturally, the ‘coherent’ version of this criterion also holds.

We note that, by Theorem 3.5, if \mathbb{T} is coherent and $\phi(\vec{x})$ is a coherent formula over Σ then $\phi(\vec{x})$ is \mathbb{T} -indecomposable if and only if for any finite family $\{\psi_i(\vec{x}) \mid i \in I\}$ of geometric formulae in the same context such that for each i , ψ_i \mathbb{T} -provably implies ϕ and

for any distinct $i, j \in I$, $\psi_i \wedge \psi_j \vdash_{\vec{x}} \perp$ is provable in \mathbb{T} , we have that $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \psi_i$ provable in \mathbb{T} implies $\phi \vdash_{\vec{x}} \psi_i$ provable in \mathbb{T} for some $i \in I$.

Regarding regular theories, their classifying toposes are always connected and locally connected; this was already observed in [Johnstone, 2002b], and also follows from our Remark 2.4.

We have already noticed that any atomic topos \mathcal{E} is locally connected; in fact, as it is observed in [Johnstone, 2002b], the atoms of \mathcal{E} are exactly the connected (equivalently, indecomposable) objects of \mathcal{E} . Hence, in view of Theorem 2.7, by replacing ‘ \mathbb{T} -indecomposable’ with ‘ \mathbb{T} -complete’ everywhere in the criteria above, one obtains syntactic criteria for a geometric (respectively coherent) theory to be classified by an atomic topos; also, one can obtain alternative versions of these criteria in the same spirit as in Remark 3.9.

Concerning atomic toposes, let us notice that if $\phi(\vec{x})$ is a \mathbb{T} -complete formula then $\phi(\vec{x})$ is \mathbb{T} -provably equivalent to a regular formula; indeed, this follows immediately from the fact that any geometric formula is provably equivalent to a disjunction of regular formulae (Lemma D1.3.8 [Johnstone, 2002b]).

We can give the following criterion for a regular theory to be classified by an atomic topos.

3.10. PROPOSITION. *Let \mathbb{T} be a regular theory over a signature Σ . Then \mathbb{T} is classified by an atomic topos if and only if every regular formula over Σ is either \mathbb{T} -provably equivalent to \perp or \mathbb{T} -complete.*

PROOF. Let $\phi(\vec{x})$ be a regular formula over Σ . If \mathbb{T} is classified by an atomic topos then, by the discussion above, ϕ is expressible as a disjunction of \mathbb{T} -complete regular formulae; but this implies, by Theorem 3.4, that either ϕ is \mathbb{T} -provably equivalent to \perp or it is \mathbb{T} -provably equivalent to one of these formulae and hence \mathbb{T} -complete.

Conversely, if every regular formula over Σ is either \mathbb{T} -provably equivalent to \perp or \mathbb{T} -complete then, by Lemma 3.2 and the fact that the set of objects of the form $y_{\mathbb{T}}^{\text{reg}}(\{\vec{x}.\phi\})$ for a regular formula $\phi(\vec{x})$ over Σ form a separating set for the classifying topos $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{reg}}, \mathcal{J}_{\mathbb{T}}^{\text{reg}})$ of \mathbb{T} , we have that $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{reg}}, \mathcal{J}_{\mathbb{T}}^{\text{reg}})$ of \mathbb{T} has a separating set of atoms and hence, by Theorem 2.7(iii), it is atomic, as required. ■

In passing, we note an interesting property of theories classified by atomic toposes.

3.11. THEOREM. *Let \mathbb{T} be a regular (respectively coherent) theory over a signature Σ which is classified by an atomic (equivalently, Boolean) topos. Then every geometric formula over Σ is \mathbb{T} -provably equivalent to a regular (respectively coherent) formula over Σ .*

PROOF. Let $\phi(\vec{x})$ be a geometric formula over Σ . Then, the classifying topos of \mathbb{T} being Boolean, $\top \vdash_{\vec{x}} \phi(\vec{x}) \vee \neg^{\mathbb{T}} \phi(\vec{x})$ is provable in \mathbb{T} (where $\neg^{\mathbb{T}} \phi(\vec{x})$ denotes the pseudocomplementation of $\phi(\vec{x})$ in $\text{Sub}_{\mathcal{C}_{\mathbb{T}}^{\text{geom}}}(\{\vec{x}.\top\})$ as in [Carmello, 2009b]). But, by Theorem 3.4, $\top(\vec{x})$ is \mathbb{T} -supercompact (respectively \mathbb{T} -compact) and, since by Lemma D1.3.8 [Johnstone, 2002b] $\phi(\vec{x})$ is (\mathbb{T}) -provably equivalent to a disjunction of regular formulae, it thus

follows that $\phi(\vec{x})$ is \mathbb{T} -provably equivalent to a single regular formula (respectively a finite disjunction of regular formulae), as required. ■

3.12. THEORIES OF PRESHEAF TYPE In this section we give a characterization of the class of geometric (respectively coherent, regular) theories classified by a presheaf topos. We recall that a theory classified by a presheaf topos is said to be of presheaf type.

Below, for a subcanonical site (\mathcal{C}, J) , we denote by $y : \mathcal{C} \rightarrow \mathbf{Sh}(\mathcal{C}, J)$ the factorization through $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ of the Yoneda embedding.

3.13. THEOREM. *Let (\mathcal{C}, J) be a subcanonical site such that $y(\mathcal{C})$ is closed in $\mathbf{Sh}(\mathcal{C}, J)$ under retracts. Then $\mathbf{Sh}(\mathcal{C}, J)$ is equivalent to a presheaf topos if and only if J is rigid.*

PROOF. The ‘if’ direction follows at once from the Comparison Lemma. Let us then prove the ‘only if’ direction. If $\mathcal{E} = \mathbf{Sh}(\mathcal{C}, J)$ is equivalent to a presheaf topos then, by Lemma C2.2.20 [Johnstone, 2002b], \mathcal{E} has a separating set of indecomposable projective objects. Now, suppose A is an indecomposable projective in \mathcal{E} . Then, as it is observed in the proof of Lemma C2.2.20 [Johnstone, 2002b], given any epimorphic family $\{f_i : B_i \rightarrow A \mid i \in I\}$, at least one f_i must be a split epimorphism; in particular A is $J_{\mathcal{E}}$ -irreducible, where $J_{\mathcal{E}}$ is the canonical coverage on \mathcal{E} . Hence, by taking as epimorphic family the collection of all the arrows in \mathcal{E} from objects of the form $y(c)$ to A , we obtain that A is a retract in \mathcal{E} of an object of the form $y(c)$. Thus, by our hypotheses, A is itself, up to isomorphism, of the form $y(c)$ for some $c \in \mathcal{C}$. Let us denote by \mathcal{C}' the full subcategory of \mathcal{C} on the objects c such that $y(c)$ is indecomposable and projective in \mathcal{E} ; then the objects in $y(\mathcal{C}')$ form a separating set for \mathcal{E} . Thus, for any object B of \mathcal{E} the family of all the arrows in \mathcal{E} from objects of the form $y(c)$ for $c \in \mathcal{C}'$ to B generates a $J_{\mathcal{E}}$ -covering sieve. But, J being subcanonical, $J = J_{\mathcal{E}}|_{\mathcal{C}}$ (by Proposition C2.2.16 [Johnstone, 2002b]) and hence for any object $c \in \mathcal{C}$ the collection of all arrows in \mathcal{C} from objects of \mathcal{C}' to c is J -covering; so, by Proposition 2.6(iii), J is rigid. ■

3.14. REMARK. We note that, under the hypotheses of the theorem, if \mathcal{C} is Cauchy-complete (in particular if \mathcal{C} is cartesian) then $y(\mathcal{C})$ is closed in $\mathcal{E} = \mathbf{Sh}(\mathcal{C}, J)$ under retracts. Indeed, let $i : A \rightarrow y(c)$, $r : y(c) \rightarrow A$ be a retract of A in \mathcal{E} , that is, $r \circ i = 1_A$. Then $i \circ r : y(c) \rightarrow y(c)$ is idempotent. Now, since y is full and faithful, $i \circ r = y(e)$ for some idempotent $e : c \rightarrow c$ in \mathcal{C} . Since \mathcal{C} is Cauchy complete then e splits as $s \circ t$ where $t \circ s = 1$. Then $y(s)$ and $y(t)$ form a retract of A and hence, by the uniqueness up to isomorphism of the splitting of an idempotent in a category, it follows that r is isomorphic to $y(t)$ and i is isomorphic to $y(s)$, and in particular A is isomorphic to $y(\text{dom}(s))$.

By Remark 3.14, the regular (respectively coherent, geometric) syntactic sites for regular (respectively coherent, geometric) theories all satisfy the hypotheses of Theorem 3.13. Thus we obtain the following results.

3.15. COROLLARY. *Let \mathbb{T} be a geometric theory over a signature Σ . Then \mathbb{T} is of presheaf type if and only if there exists a collection \mathcal{F} of geometric formulae-in-context over Σ satisfying the following properties:*

- (i) for any geometric formula $\{\vec{y}.\psi\}$ over Σ , there exist objects $\{\vec{x}_i.\phi_i\}$ in \mathcal{F} as i varies in I and \mathbb{T} -provably functional geometric formulae $\{\vec{x}_i, \vec{y}.\theta_i\}$ from $\{\vec{x}_i.\phi_i\}$ to $\{\vec{y}.\psi\}$ such that $\psi \vdash_{\vec{y}} \bigvee_{i \in I} (\exists \vec{x}_i)\theta_i$ is provable in \mathbb{T} ;
- (ii) for any formula $\{\vec{x}.\phi\}$ in \mathcal{F} , for any family $\{\theta_i \mid i \in I\}$ of \mathbb{T} -provably functional geometric formulae $\{\vec{x}_i, \vec{x}.\theta_i\}$ from $\{\vec{x}_i.\phi_i\}$ to $\{\vec{x}.\phi\}$ such that $\phi \vdash_{\vec{x}} \bigvee_{i \in I} (\exists \vec{x}_i)\theta_i$ is provable in \mathbb{T} , there exist $i \in I$ and a \mathbb{T} -provably functional geometric formula $\{\vec{x}, \vec{x}_i.\theta'\}$ from $\{\vec{x}.\phi\}$ to $\{\vec{x}_i.\phi_i\}$ such that $\phi \vdash_{\vec{x}} (\exists \vec{x}_i)(\theta' \wedge \theta_i)$ is provable in \mathbb{T} .

Note that condition (ii) in the corollary says precisely that $\{\vec{x}.\phi\}$ is \mathbb{T} -irreducible; in particular, $\{\vec{x}.\phi\}$ is \mathbb{T} -supercompact, that is, for any family $\{\{\vec{x}.\phi_i\} \mid i \in I\}$ of geometric formulae in the same context which \mathbb{T} -provably imply $\{\vec{x}.\phi\}$ and such that $\phi \vdash_{\vec{x}} \bigvee_{i \in I} \phi_i$ is provable in \mathbb{T} , there exists $i \in I$ such that ϕ_i and ϕ are \mathbb{T} -provably equivalent.

The following results are the coherent and regular analogues of this corollary.

3.16. COROLLARY. *Let \mathbb{T} be a coherent theory over a signature Σ . Then \mathbb{T} is of presheaf type if and only if there exists a collection \mathcal{F} of coherent formulae-in-context over Σ satisfying the following properties:*

- (i) for any coherent formula $\{\vec{y}.\psi\}$ over Σ , there exists a finite number of objects $\{\vec{x}_i.\phi_i\}$ in \mathcal{F} as i varies in I and \mathbb{T} -provably functional coherent formulae $\{\vec{x}_i, \vec{y}.\theta_i\}$ from $\{\vec{x}_i.\phi_i\}$ to $\{\vec{y}.\psi\}$ such that $\psi \vdash_{\vec{y}} \bigvee_{i \in I} (\exists \vec{x}_i)\theta_i$ is provable in \mathbb{T} ;
- (ii) for any formula $\{\vec{x}.\phi\}$ in \mathcal{F} , for any finite family $\{\theta_i \mid i \in I\}$ of \mathbb{T} -provably functional coherent formulae $\{\vec{x}_i, \vec{x}.\theta_i\}$ from $\{\vec{x}_i.\phi_i\}$ to $\{\vec{x}.\phi\}$ such that $\phi \vdash_{\vec{x}} \bigvee_{i \in I} (\exists \vec{x}_i)\theta_i$ is provable in \mathbb{T} , there exist $i \in I$ and a \mathbb{T} -provably functional coherent formula $\{\vec{x}, \vec{x}_i.\theta'\}$ from $\{\vec{x}.\phi\}$ to $\{\vec{x}_i.\phi_i\}$ such that $\phi \vdash_{\vec{x}} (\exists \vec{x}_i)(\theta' \wedge \theta_i)$ is provable in \mathbb{T} .

3.17. COROLLARY. *Let \mathbb{T} be a regular theory over a signature Σ . Then \mathbb{T} is of presheaf type if and only if there exists a collection \mathcal{F} of regular formulae-in-context over Σ satisfying the following properties:*

- (i) for any regular formula $\{\vec{y}.\psi\}$ over Σ , there exists an object $\{\vec{x}.\phi\}$ in \mathcal{F} and a \mathbb{T} -provably functional formula $\{\vec{x}, \vec{y}.\theta\}$ from $\{\vec{x}.\phi\}$ to $\{\vec{y}.\psi\}$ such that $\psi \vdash_{\vec{y}} (\exists \vec{x})\theta$ is provable in \mathbb{T} ;
- (ii) for any formula $\{\vec{y}.\psi\}$ in \mathcal{F} , for any \mathbb{T} -provably functional regular formulae $\{\vec{x}, \vec{y}.\theta\}$ from $\{\vec{x}.\phi\}$ to $\{\vec{y}.\psi\}$ such that $\psi \vdash_{\vec{y}} (\exists \vec{x})\theta$ is provable in \mathbb{T} , there exist a \mathbb{T} -provably functional regular formula $\{\vec{y}, \vec{x}.\theta'\}$ from $\{\vec{y}.\psi\}$ to $\{\vec{x}.\phi\}$ such that $\psi \vdash_{\vec{y}} (\exists \vec{x})(\theta' \wedge \theta)$ is provable in \mathbb{T} .

4. Syntactic properties of quotients of theories of presheaf type

Let \mathbb{T} be a theory of presheaf type over a signature Σ . Recall that a \mathbb{T} -model M in \mathbf{Set} is said to be finitely presentable if the representable functor $\mathrm{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(M, -) : \mathbb{T}\text{-mod}(\mathbf{Set}) \rightarrow \mathbf{Set}$ preserves filtered colimits. The full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$ on the finitely presentable models will be denoted by $\mathrm{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$.

A \mathbb{T} -model M in \mathbf{Set} is said to be finitely presented if there is a geometric formula $\{\vec{x}.\phi\}$ over Σ and a string of elements $(\xi_1, \dots, \xi_n) \in MA_1 \times \dots \times MA_n$ (where A_1, \dots, A_n are the sorts of the variables in \vec{x}), called the generators of M , such that for any \mathbb{T} -model N in \mathbf{Set} and string of elements $(b_1, \dots, b_n) \in MA_1 \times \dots \times MA_n$ such that $(b_1, \dots, b_n) \in [[\phi]]_N$, there exists a unique arrow $f : M \rightarrow N$ in $\mathbb{T}\text{-mod}(\mathbf{Set})$ such that $(fA_1 \times \dots \times fA_n)((\xi_1, \dots, \xi_n)) = (b_1, \dots, b_n)$.

We recall from [Caramello, 2009b] that, given a theory of presheaf type \mathbb{T} , a canonical Morita-equivalence for \mathbb{T} (in the sense of [Caramello, 2009b]) induces an equivalence of classifying toposes

$$[\mathrm{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}] \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\mathrm{geom}}, J_{\mathbb{T}}^{\mathrm{geom}})$$

sending the universal model $M_{\mathbb{T}}$ of \mathbb{T} in the topos $[\mathrm{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ to the universal model $U_{\mathbb{T}}$ of \mathbb{T} in $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\mathrm{geom}}, J_{\mathbb{T}}^{\mathrm{geom}})$. Recall that $M_{\mathbb{T}}$ is the \mathbb{T} -model in $[\mathrm{f.p.}\mathbb{T}\text{-Mod}(\mathbf{Set}), \mathbf{Set}]$ which assigns to a sort A the functor $M_{\mathbb{T}}A$ given by $(M_{\mathbb{T}}A)(M) = MA$, to a function symbol $f : A_1 \cdots A_n \rightarrow B$ the morphism $M_{\mathbb{T}}A_1 \times \cdots \times M_{\mathbb{T}}A_n \rightarrow M_{\mathbb{T}}B$ given by $(M_{\mathbb{T}}f)(M) = Mf$ and to a relation symbol $R \rhd A_1 \cdots A_n$ the subobject $M_{\mathbb{T}}R \rhd M_{\mathbb{T}}A_1 \times \cdots \times M_{\mathbb{T}}A_n$ given by $(M_{\mathbb{T}}R)(M) = MR$ (for any $M \in \mathrm{f.p.}\mathbb{T}\text{-Mod}(\mathbf{Set})$), while $U_{\mathbb{T}}$ is the image, under the Yoneda embedding $y : \mathcal{C}_{\mathbb{T}} \hookrightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$, of the ‘universal’ model $M_{\mathbb{T}}$ of \mathbb{T} in $\mathcal{C}_{\mathbb{T}}$, which assigns to a sort A the object $\{x^A.\top\}$ where x^A is a variable of sort A , to a function symbol $f : A_1 \cdots A_n \rightarrow B$ the morphism

$$\{x_1, \dots, x_n.\top\} \xrightarrow{[f(x_1, \dots, x_n)=y]} \{y.\top\}$$

and to a relation symbol $R \rhd A_1 \cdots A_n$ the subobject

$$\{x_1, \dots, x_n.R(x_1, \dots, x_n)\} \xrightarrow{[R(x_1, \dots, x_n)]} \{x_1, \dots, x_n.\top\}$$

as in Lemma D1.4.4(iv) [Johnstone, 2002b].

As a consequence, if $M \in \mathrm{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ is a \mathbb{T} -model presented by a formula $\phi(\vec{x})$ over Σ then, denoted by $y : \mathrm{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\mathrm{op}} \rightarrow [\mathrm{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ the Yoneda embedding, $y(M)$ is equal to $[[\vec{x}.\phi]]_{M_{\mathbb{T}}}$ and hence corresponds, under the equivalence

$$[\mathrm{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}] \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\mathrm{geom}}, J_{\mathbb{T}}^{\mathrm{geom}}),$$

to the functor $y_{\mathbb{T}}^{\mathrm{geom}}(\{\vec{x}.\phi\}) = [[\vec{x}.\phi]]_{U_{\mathbb{T}}}$.

We recall that in [Caramello, 2009c] a bijection is established between the subtoposes of the classifying topos of a geometric theory \mathbb{T} and the quotients of \mathbb{T} (i.e., geometric extensions of \mathbb{T} over its signature), considered up to the obvious notion of syntactic

equivalence (i.e. the equivalence which identifies two quotients of \mathbb{T} precisely when they prove the same geometric sequents over the signature of \mathbb{T}); the subtopos corresponding via this duality to a given quotient of \mathbb{T} can be identified with its classifying topos (cf. Theorem 3.6 [Caramello, 2009c]).

Now, if \mathbb{T}' is a quotient of \mathbb{T} then the subtopos of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$ corresponding to it via this duality transfers via the equivalence

$$[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}] \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$$

to a subtopos

$$\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J) \hookrightarrow [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$$

of $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$; the topology J will be called the associated \mathbb{T} -topology of \mathbb{T}' . This gives rise to an equivalence

$$\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J) \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}^{\text{geom}}, J_{\mathbb{T}'}^{\text{geom}})$$

of classifying toposes of \mathbb{T}' which sends $l_J^{\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}}(M)$ to the functor $y_{\mathbb{T}'}^{\text{geom}}(\{\vec{x}.\phi\}) = [[\vec{x}.\phi]]_{U_{\mathbb{T}'}}$.

The following result provides a link between ‘geometrical’ properties of J and syntactic properties of \mathbb{T}' .

4.1. THEOREM. *Let \mathbb{T} be a theory of presheaf type over a signature Σ , \mathbb{T}' be a quotient of \mathbb{T} with associated \mathbb{T} -topology J on $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ and $\phi(\vec{x})$ be a geometric formula over Σ which presents a \mathbb{T} -model M . Then*

- (i) $\phi(\vec{x})$ is \mathbb{T} -irreducible; in particular, $\phi(\vec{x})$ is \mathbb{T} -provably equivalent to a regular formula;
- (ii) if the site $(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J)$ is locally connected (for example when $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ satisfies the right Ore condition and every J -covering sieve is non-empty) then $\phi(\vec{x})$ is \mathbb{T}' -indecomposable;
- (iii) if $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ satisfies the right Ore condition and J is the atomic topology on $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ then $\phi(\vec{x})$ is \mathbb{T}' -complete;
- (iv) if every J -covering sieve on M contains a J -covering sieve generated by a finite family of morphisms (respectively by a single morphism) then $\phi(\vec{x})$ is \mathbb{T}' -compact (respectively \mathbb{T}' -supercompact).

PROOF. (i) By Lemma 3.2(iii), $\phi(\vec{x})$ is \mathbb{T} -irreducible if and only if $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\})$ is an irreducible object of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}})$. But, by the discussion above, this is equivalent to saying that $y(M)$ is irreducible in $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$, and this is true by Proposition 2.5(iii). The fact that $\phi(\vec{x})$ is \mathbb{T} -provably equivalent to a regular formula then follows from Lemma D1.3.8(ii) [Johnstone, 2002b].

(ii) and (iii) By Lemma 3.2(ii) (respectively Lemma 3.2(i)), $\phi(\vec{x})$ is \mathbb{T}' -indecomposable (respectively \mathbb{T}' -complete) if and only if $y_{\mathbb{T}'}^{\text{geom}}(\{\vec{x}.\phi\})$ is an indecomposable object (respectively an atom) of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}^{\text{geom}}, J_{\mathbb{T}'}^{\text{geom}})$; but this is equivalent to saying that $l_J^{\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}}(M)$ is an indecomposable object (respectively an atom) of $\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J)$, and this is true by Proposition 2.5(i) (respectively Proposition 2.5(ii)).

(iv) By Lemma 3.2(iii), $\phi(\vec{x})$ is \mathbb{T}' -compact (respectively \mathbb{T}' -supercompact) if and only if $y_{\mathbb{T}'}^{\text{geom}}(\{\vec{x}.\phi\})$ is a compact (respectively supercompact) object of $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}^{\text{geom}}, J_{\mathbb{T}'}^{\text{geom}})$; but this is equivalent to saying that $l_J^{\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}}(M)$ is a compact (respectively supercompact) object of $\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J)$, and this is true by Proposition 2.6(i) (respectively Proposition 2.6(ii)). \blacksquare

4.2. REMARK. The theorem can be profitably applied in the context of cartesian theories. Indeed, if \mathbb{T} is a cartesian theory then every cartesian formula over Σ presents a \mathbb{T} -model so that we have an equivalence between $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ and the cartesian syntactic category of \mathbb{T} . Thus the theorem provides syntactic properties of cartesian formulae in particular quotients of \mathbb{T} ; for example, for any such formula $\phi(\vec{x})$, part (iii) of the lemma yields that $\phi(\vec{x})$ is \mathbb{T}' -complete where \mathbb{T}' is the Booleanization of \mathbb{T} (as defined in [Caramello, 2009a]).

As an application of the notion of irreducible object in a topos, we can prove the following result.

4.3. THEOREM. *Let \mathbb{T} be a theory of presheaf type over a signature Σ . Then*

(i) *Any finitely presentable \mathbb{T} -model in \mathbf{Set} is presented by a \mathbb{T} -irreducible geometric formula $\phi(\vec{x})$ over Σ ;*

(ii) *Conversely, any \mathbb{T} -irreducible geometric formula $\phi(\vec{x})$ over Σ presents a \mathbb{T} -model.*

In fact, the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ is equivalent to the full subcategory of $\mathcal{C}_{\mathbb{T}}^{\text{geom}}$ on the \mathbb{T} -irreducible formulae.

PROOF. We have already observed that we have an equivalence

$$\tau : \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}}) \simeq [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$$

of classifying toposes for \mathbb{T} . Now, if $\mathcal{C}_{\mathbb{T}}^{\text{irr}}$ is the full subcategory of $\mathcal{C}_{\mathbb{T}}^{\text{geom}}$ on the \mathbb{T} -irreducible formulae then, by Theorem 3.13, we have that $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}^{\text{geom}}, J_{\mathbb{T}}^{\text{geom}}) \simeq [(\mathcal{C}_{\mathbb{T}}^{\text{irr}})^{\text{op}}, \mathbf{Set}]$ via the Comparison Lemma (cf. the proof of Theorem 2.5). Now, if $\tilde{\mathcal{C}}_{\mathbb{T}}^{\text{irr}} \hookrightarrow \mathcal{C}_{\mathbb{T}}^{\text{geom}}$ is the Cauchy-completion of $\mathcal{C}_{\mathbb{T}}^{\text{irr}}$ then $[(\mathcal{C}_{\mathbb{T}}^{\text{irr}})^{\text{op}}, \mathbf{Set}] \simeq [(\tilde{\mathcal{C}}_{\mathbb{T}}^{\text{irr}})^{\text{op}}, \mathbf{Set}]$ and the resulting equivalence

$$[(\tilde{\mathcal{C}}_{\mathbb{T}}^{\text{irr}})^{\text{op}}, \mathbf{Set}] \simeq [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$$

restricts to an equivalence

$$l : \tilde{\mathcal{C}}_{\mathbb{T}}^{\text{irr}} \simeq \text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$$

between the subcategories of indecomposable projective objects. Now, given $\{\vec{x}.\phi\} \in \tilde{\mathcal{C}}_{\mathbb{T}}^{\text{irr}}$, τ sends the functor $y_{\mathbb{T}}^{\text{geom}}(\{\vec{x}.\phi\}) = [[\vec{x}.\phi]]_{U_{\mathbb{T}}}$ to $y(l(\{\vec{x}.\phi\})) = [[\vec{x}.\phi]]_{M_{\mathbb{T}}}$, from which it follows that the model $l(\{\vec{x}.\phi\})$ is finitely presented by $\{\vec{x}.\phi\}$. Then, by Theorem 4.1(i), $\phi(\vec{x})$ is \mathbb{T} -irreducible. So we conclude that $\tilde{\mathcal{C}}_{\mathbb{T}}^{\text{irr}}$ is equal to $\mathcal{C}_{\mathbb{T}}^{\text{irr}}$, that is, $\mathcal{C}_{\mathbb{T}}^{\text{irr}}$ is Cauchy complete, and hence l gives an equivalence

$$\tilde{\mathcal{C}}_{\mathbb{T}}^{\text{irr}\text{op}} \simeq \text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}).$$

It is easy to verify that this equivalence coincide with the dualizing functor d of Theorem 3.6 [Caramello, 2009b]. ■

4.4. **REMARK.** As an application of Theorem 4.3 and Remark 2.9, suppose that J is the associated \mathbb{T} -topology of a quotient \mathbb{T}' of \mathbb{T} . Then, if for any \mathbb{T} -irreducible formula (equivalently, formula presenting a \mathbb{T} -model) $\phi(\vec{x})$, $\phi(\vec{x})$ is \mathbb{T}' -indecomposable (respectively \mathbb{T}' -complete) then the classifying topos $\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J)$ of \mathbb{T}' is locally connected (respectively atomic); indeed, as observed above, $\phi(\vec{x})$ is \mathbb{T}' -indecomposable (respectively \mathbb{T}' -atomic) if and only if $y_{\mathbb{T}'}^{\text{geom}}(\{\vec{x}.\phi\})$ is an indecomposable object (respectively an atom) of the topos $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}^{\text{geom}}, J_{\mathbb{T}'}^{\text{geom}})$, if and only if $l_J^{\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}}(M)$ is an indecomposable object (respectively an atom) of the topos $\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J)$.

Acknowledgements: I am grateful to Peter Johnstone for useful discussions on the subject matter of this paper, and to the anonymous referee for his suggestions, which have helped improve the readability of this paper.

References

- [Artin, Grothendieck and Verdier, 1972] M. Artin, A. Grothendieck and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas*, Lecture Notes in Math., vols 269, 270 and 305 (Springer-Verlag, 1972).
- [Caramello, 2011] O. Caramello, Atomic toposes and countable categoricity, *Applied Categorical Structures*, doi:10.1007/s10485-010-9244-x.
- [Caramello, 2009a] O. Caramello, De Morgan classifying toposes, *Advances in Mathematics* 222 (2009), 2117-2144.
- [Caramello, 2009b] O. Caramello, Universal models and definability, *Math. Proc. Camb. Phil. Soc.* (2012), 152, 279302, 2011, doi:10.1017/S0305004111000624.
- [Caramello, 2009c] O. Caramello, Lattices of theories, *arXiv:math.CT/0905.0299*.
- [Caramello, 2010] O. Caramello, The unification of Mathematics via Topos Theory, *arXiv:math.CT/1006.3930v1*.
- [Johnstone, 2002a] P. T. Johnstone, *Sketches of an Elephant: a topos theory compendium. Vol.1*, vol. 43 of *Oxford Logic Guides* (Oxford University Press, 2002).

[Johnstone, 2002b] P. T. Johnstone, *Sketches of an Elephant: a topos theory compendium. Vol.2*, vol. 44 of *Oxford Logic Guides* (Oxford University Press, 2002).

[Mac Lane and Moerdijk, 1992] S. Mac Lane and I. Moerdijk, *Sheaves in geometry and logic: a first introduction to topos theory* (Springer-Verlag, 1992).

[Makkai and Reyes, 1977] M. Makkai and G. E. Reyes, *First order categorical logic*, Lecture Notes in Math. vol. 611 (Springer-Verlag, 1977).

*Department of Pure Mathematics and Mathematical Statistics,
University of Cambridge,
Centre for Mathematical Sciences, Wilberforce Road,
Cambridge, CB3 0WB,
United Kingdom*

Email: `O.Caramello@dpms.cam.ac.uk`

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/26/6/26-06.{dvi,ps,pdf}>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\text{e}$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

$\text{T}_{\text{E}}\text{X}$ NICAL EDITOR Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT $\text{T}_{\text{E}}\text{X}$ EDITOR Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS

Clemens Berger, Université de Nice-Sophia Antipolis, cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva: valeria.depaiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne : kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

P. T. Johnstone, University of Cambridge: ptj@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Glasgow, Tom.Leinster@glasgow.ac.uk

Jean-Louis Loday, Université de Strasbourg: loday@math.u-strasbg.fr

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Edinburgh: Alex.Simpson@ed.ac.uk

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca