

THE CORE OF ADJOINT FUNCTORS

ROSS STREET

ABSTRACT. There is a lot of redundancy in the usual definition of adjoint functors. We define and prove the core of what is required. First we do this in the hom-enriched context. Then we do it in the cocompletion of a bicategory with respect to Kleisli objects, which we then apply to internal categories. Finally, we describe a doctrinal setting.

1. Introduction

Kan [7] introduced the notion of adjoint functors. By defining the unit and counit natural transformations, he paved the way for the notion to be internalized to any 2-category. This was done by Kelly [8] whose interest at the time was particularly in the 2-category of \mathcal{V} -categories for a monoidal category \mathcal{V} (in the sense of Eilenberg-Kelly [4]).

During my Topology lectures at Macquarie University in the 1970s, the students and I realized, in proving that a function f between posets was order preserving when there was a function u in the reverse direction such that $f(x) \leq a$ if and only if $x \leq u(a)$, did not require u to be order preserving. I realized then that knowing functors in the two directions only on objects and the usual hom adjointness isomorphism implied the effect of the functors on homs was uniquely determined. Writing this down properly led to the present paper.

Section 2 merely reviews adjunctions between enriched categories. Section 3 introduces the notion of core of an enriched adjunction: it only involves the object assignments of the two functors and a hom isomorphism with no naturality requirement. The main result characterizes when such a core is an adjunction.

The material becomes increasingly for mature audiences; that is, for those with knowledge of bicategories. Sections 4 and 5 present results about adjunctions in the Kleisli object cocompletion of a bicategory in the sense of [13]. In particular, this is applied in Section 6 to adjunctions for categories internal to a finitely complete category. By a different choice of bicategory, where enriched categories can be seen as monads (see [2]), we could rediscover the work of Section 3; however, we leave this to the interested reader. In Section 7 we describe a general setting involving a pseudomonad (doctrine) on a bicategory.

The author gratefully acknowledges the support of an Australian Research Council Discovery Grant DP1094883.

Received by the editors 2012-01-01 and, in revised form, 2012-05-29.

Published on 2012-06-11 in the volume of articles from CT2011.

2000 Mathematics Subject Classification: 18A40; 18D10; 18D05.

Key words and phrases: adjoint functor; enriched category; bicategory; Kleisli cocompletion.

© Ross Street, 2012. Permission to copy for private use granted.

2. Adjunctions

For \mathcal{V} -categories \mathcal{A} and \mathcal{X} , an *adjunction* consists of

1. \mathcal{V} -functors $U : \mathcal{A} \longrightarrow \mathcal{X}$ and $F : \mathcal{X} \longrightarrow \mathcal{A}$;
2. a \mathcal{V} -natural family of isomorphisms $\pi : \mathcal{A}(FX, A) \cong \mathcal{X}(X, UA)$ in \mathcal{V} indexed by $A \in \mathcal{A}$, $X \in \mathcal{X}$.

We write $\pi : F \dashv U : \mathcal{A} \longrightarrow \mathcal{X}$.

The following result is well known; for example see Section 1.11 of [10].

2.1. PROPOSITION. *Suppose $U : \mathcal{A} \longrightarrow \mathcal{X}$ is a \mathcal{V} -functor, $F : \text{ob}\mathcal{X} \longrightarrow \text{ob}\mathcal{A}$ is a function, and, for each $X \in \mathcal{X}$, $\pi : \mathcal{A}(FX, A) \cong \mathcal{X}(X, UA)$ is a family of isomorphisms \mathcal{V} -natural in $A \in \mathcal{A}$. Then there exists a unique adjunction $\pi : F \dashv U : \mathcal{A} \longrightarrow \mathcal{X}$ for which $F : \text{ob}\mathcal{X} \longrightarrow \text{ob}\mathcal{A}$ is the effect of the \mathcal{V} -functor $F : \mathcal{X} \longrightarrow \mathcal{A}$ on objects.*

3. Cores

3.1. DEFINITION. *For \mathcal{V} -categories \mathcal{A} and \mathcal{X} , an *adjunction core* consists of*

1. *functions $U : \text{ob}\mathcal{A} \longrightarrow \text{ob}\mathcal{X}$ and $F : \text{ob}\mathcal{X} \longrightarrow \text{ob}\mathcal{A}$;*
2. *a family of isomorphisms $\pi : \mathcal{A}(FX, A) \cong \mathcal{X}(X, UA)$ in \mathcal{V} indexed by $A \in \mathcal{A}$, $X \in \mathcal{X}$.*

Given such a core, we make the following definitions:

- (a) $\beta_X : X \longrightarrow UFX$ is the composite

$$I \xrightarrow{j} \mathcal{A}(FX, FX) \xrightarrow{\pi} \mathcal{X}(X, UFX);$$

- (b) $\alpha_A : FUA \longrightarrow A$ is the composite

$$I \xrightarrow{j} \mathcal{X}(UA, UA) \xrightarrow{\pi^{-1}} \mathcal{A}(FUA, X);$$

- (c) $U_{AB} : \mathcal{A}(A, B) \longrightarrow \mathcal{X}(UA, UB)$ is the composite

$$\mathcal{A}(A, B) \xrightarrow{\mathcal{A}(\alpha_A, 1)} \mathcal{A}(FUA, B) \xrightarrow{\pi} \mathcal{X}(UA, UB);$$

- (d) $F_{XY} : \mathcal{X}(X, Y) \longrightarrow \mathcal{A}(FX, FY)$ is the composite

$$\mathcal{X}(X, Y) \xrightarrow{\mathcal{X}(1, \beta_Y)} \mathcal{X}(X, UFY) \xrightarrow{\pi^{-1}} \mathcal{A}(FX, FY).$$

Clearly each adjunction $\pi : F \dashv U : \mathcal{A} \longrightarrow \mathcal{X}$ includes an adjunction core as part of its data. Then it follows directly from the Yoneda lemma and the definitions (a) and (b) that the effect of U and F on homs are as in (c) and (d).

3.2. THEOREM. *An adjunction core extends to an adjunction if and only if one of the diagrams (3.1) or (3.2) below commutes. The adjunction is unique when it exists.*

$$\begin{array}{ccc} \mathcal{A}(A, B) \otimes \mathcal{A}(FX, A) & \xrightarrow{U_{A,B} \otimes \pi} & \mathcal{X}(UA, UB) \otimes \mathcal{X}(X, UA) \\ \text{comp} \downarrow & & \downarrow \text{comp} \\ \mathcal{A}(FX, B) & \xrightarrow{\pi} & \mathcal{X}(X, UB) \end{array} \quad (3.1)$$

$$\begin{array}{ccc} \mathcal{X}(Y, UA) \otimes \mathcal{X}(X, Y) & \xrightarrow{\pi^{-1} \otimes F_{X,Y}} & \mathcal{A}(FY, A) \otimes \mathcal{A}(FX, FY) \\ \text{comp} \downarrow & & \downarrow \text{comp} \\ \mathcal{X}(X, UA) & \xrightarrow{\pi^{-1}} & \mathcal{A}(FX, A) \end{array} \quad (3.2)$$

PROOF. We deal first with the version involving diagram (3.1). For an adjunction, (3.1) expresses the \mathcal{V} -naturality of π in $A \in \mathcal{A}$. Conversely, given an adjunction core satisfying (3.1), we paste to the left of (3.1) with $X = UC$, the diagram

$$\begin{array}{ccc} \mathcal{A}(A, B) \otimes \mathcal{A}(C, A) & \xrightarrow{1 \otimes \mathcal{A}(\alpha_C, 1)} & \mathcal{A}(A, B) \otimes \mathcal{A}(FUC, A) \\ \text{comp} \downarrow & & \downarrow \text{comp} \\ \mathcal{A}(C, B) & \xrightarrow{\mathcal{A}(\alpha_C, 1)} & \mathcal{A}(FUC, B) \end{array} \quad (3.3)$$

which commutes by naturality of composition. This leads to the following commutative square.

$$\begin{array}{ccc} \mathcal{A}(A, B) \otimes \mathcal{A}(C, A) & \xrightarrow{U \otimes U} & \mathcal{X}(UA, UB) \otimes \mathcal{X}(UC, UA) \\ \text{comp} \downarrow & & \downarrow \text{comp} \\ \mathcal{A}(C, B) & \xrightarrow{U} & \mathcal{X}(UC, UB) \end{array} \quad (3.4)$$

We also have the equality

$$\left(I \xrightarrow{j} \mathcal{A}(A, A) \xrightarrow{U} \mathcal{X}(UA, UB) \right) = \left(I \xrightarrow{j} \mathcal{X}(UA, UB) \right) \quad (3.5)$$

straight from the definitions (b) and (c). Together (3.4) and (3.5) tell us that U is a \mathcal{V} -functor. Now the general diagram (3.1) expresses the \mathcal{V} -naturality of π in A . By Proposition 2.1, we have an adjunction determined uniquely by the core.

Writing \mathcal{V}^{rev} for \mathcal{V} with the reversed monoidal structure $A \otimes^{\text{rev}} B = B \otimes A$, and applying the first part of this proof to the \mathcal{V}^{rev} -enriched adjunction $\pi^{-1} : U^{\text{op}} \dashv F^{\text{op}} : \mathcal{X}^{\text{op}} \rightarrow \mathcal{A}^{\text{op}}$, which is the same as an adjunction $\pi : F \dashv U : \mathcal{A} \rightarrow \mathcal{X}$, we see that it is equivalent to an adjunction core satisfying (3.2). ■

3.3. COROLLARY. *If \mathcal{V} is a poset then adjunction cores are adjunctions.*

PROOF. All diagrams, including (3.1), commute in such a \mathcal{V} . ■

An object X of a category is often called *preterminal* when, for all objects K , there is at most one morphism $K \rightarrow X$. In the category \mathbf{Set} , both the initial and terminal objects are preterminal. A poset is a category in which every object is preterminal. A \mathcal{V} -category \mathcal{A} is *locally preterminal* when each $\mathrm{hom} \mathcal{A}(A, B)$ is preterminal in \mathcal{V} . Using this terminology, we can state a more general corollary.

3.4. COROLLARY. *If in Definition 3.1 either \mathcal{A} or \mathcal{X} is locally preterminal then the adjunction core is an adjunction.*

4. Adjunctions between monads

This section will discuss adjunctions in a particular bicategory $\mathrm{KL}(\mathcal{K})$ of monads in a bicategory \mathcal{K} . The results will apply to adjunctions between categories internal to a category \mathcal{C} with pullbacks.

As well as defining bicategories Bénabou [1] defined, for each pair of bicategories \mathcal{A} and \mathcal{K} , a bicategory $\mathrm{Bicat}(\mathcal{A}, \mathcal{K})$ whose objects are morphisms $\mathcal{A} \rightarrow \mathcal{K}$ of bicategories (also called lax functors), whose morphisms are transformations (also called lax natural transformations), and whose 2-cells are modifications. In particular, $\mathrm{Bicat}(\mathbf{1}, \mathcal{K})$ is one bicategory whose objects are monads in \mathcal{K} ; it was called $\mathrm{Mnd}(\mathcal{K})$ in [17] for the case of a 2-category \mathcal{K} , where it was used to discuss Eilenberg–Moore objects in \mathcal{K} . We shall also use the notation $\mathrm{Mnd}(\mathcal{K})$ when \mathcal{K} is a bicategory.

We write $\mathcal{K}^{\mathrm{op}}$ for the dual of \mathcal{K} obtained by reversing morphisms (not 2-cells). Monads in $\mathcal{K}^{\mathrm{op}}$ are the same as monads in \mathcal{K} . So we also have the bicategory

$$\mathrm{Mnd}^{\mathrm{op}}(\mathcal{K}) = \mathrm{Bicat}(\mathbf{1}, \mathcal{K}^{\mathrm{op}})^{\mathrm{op}}$$

whose objects are monads in \mathcal{K} . This was used in [17] to discuss Kleisli objects in \mathcal{K} .

Two more bicategories $\mathrm{EM}(\mathcal{K})$ and $\mathrm{KL}(\mathcal{K})$, with objects monads in \mathcal{K} , were defined in [13]. The first freely adjoins Eilenberg–Moore objects and the second freely adjoins Kleisli objects to \mathcal{K} . In fact, $\mathrm{EM}(\mathcal{K})$ has the same objects and morphisms as $\mathrm{Mnd}(\mathcal{K})$ but different 2-cells while $\mathrm{KL}(\mathcal{K})$ has the same objects and morphisms as $\mathrm{Mnd}^{\mathrm{op}}(\mathcal{K})$ but different 2-cells.

A *monad* in a bicategory \mathcal{K} is an object A equipped with a morphism $s : A \rightarrow A$ and 2-cells $\eta : 1_A \rightarrow s$ and $\mu : ss \rightarrow s$ such that

$$\begin{array}{ccc}
 & s(ss) & \\
 \cong \nearrow & & \searrow s\mu \\
 (ss)s & & ss \\
 \mu s \downarrow & & \downarrow \mu \\
 ss & \xrightarrow{\mu} & w
 \end{array} \tag{4.1}$$

and the composites

$$s1 \xrightarrow{s\eta} ss \xrightarrow{\mu} s \quad \text{and} \quad 1s \xrightarrow{\eta s} ss \xrightarrow{\mu} s \tag{4.2}$$

should be the canonical isomorphisms. We shall use the same symbols η and μ for the *unit* and *multiplication* of all monads; so we simply write (A, s) for the monad.

For monads (A, s) and (A', s') in \mathcal{K} , a *monad opmorphism* $(f, \phi) : (A, s) \rightarrow (A', s')$ consists of a morphism $f : A \rightarrow A'$ and a 2-cell $\phi : fs \rightarrow s'f$ in \mathcal{K} such that

$$\begin{aligned} \left((fs) s \xrightarrow{\phi s} (s'f) s \xrightarrow{\cong} s'(fs) \xrightarrow{s'\phi} s'(s'f) \xrightarrow{\cong} s'(s'f) \xrightarrow{\mu f} s'f \right) \\ = \left((fs) s \xrightarrow{\cong} f(ss) \xrightarrow{f\mu} fs \xrightarrow{\phi} s'f \right) \end{aligned} \quad (4.3)$$

and

$$\left(f1 \xrightarrow{f\eta} fs \xrightarrow{\phi} s'f \right) = \left(f1 \xrightarrow{\cong} 1f \xrightarrow{\eta f} s'f \right). \quad (4.4)$$

The composite of monad opmorphisms $(f, \phi) : (A, s) \rightarrow (A', s')$ and $(f', \phi') : (A', s') \rightarrow (A'', s'')$ is defined to be $(f'f, \phi' \star \phi) : (A, s) \rightarrow (A'', s'')$ where $\phi' \star \phi$ is the composite

$$(f'f) s \xrightarrow{\cong} f'(fs) \xrightarrow{f'\phi} f'(s'f) \xrightarrow{\cong} (f's') f \xrightarrow{\phi' f} (s''f') f \xrightarrow{\cong} s''(f'f) . \quad (4.5)$$

The objects of both $\text{Mnd}^{\text{op}}(\mathcal{K})$ and $\text{KL}(\mathcal{K})$ are monads (A, s) in \mathcal{K} . The morphisms in both are the opmorphisms (f, ϕ) . The 2-cells $\sigma : (f, \phi) \rightarrow (g, \psi) : (A, s) \rightarrow (A', s')$ in $\text{Mnd}^{\text{op}}(\mathcal{K})$ are 2-cells $\sigma : f \rightarrow g$ in \mathcal{K} such that the following square commutes.

$$\begin{array}{ccc} fs & \xrightarrow{\phi} & s'f \\ \sigma s \downarrow & & \downarrow s'\sigma \\ gs & \xrightarrow{\psi} & s'g \end{array} \quad (4.6)$$

Vertical and horizontal composition in $\text{Mnd}^{\text{op}}(\mathcal{K})$ are performed in the obvious way so that the projection $\text{Und} : \text{Mnd}^{\text{op}}(\mathcal{K}) \rightarrow \mathcal{K}$, taking (A, s) to A , (f, ϕ) to f , and σ to σ , preserves them. The associativity and unit isomorphisms in $\text{Mnd}^{\text{op}}(\mathcal{K})$ are also such that Und preserves them, making Und a strict morphism of bicategories.

A 2-cell $\rho : (f, \phi) \rightarrow (g, \psi) : (A, s) \rightarrow (A', s')$ in $\text{KL}(\mathcal{K})$ is a 2-cell $\rho : f \rightarrow s'g$ in \mathcal{K} such that

$$\begin{aligned} \left(fs \xrightarrow{\phi} s'f \xrightarrow{s'\rho} s'(s'g) \xrightarrow{\cong} (s's') g \xrightarrow{\mu g} s'g \right) \\ = \left(fs \xrightarrow{\rho s} (s'g) s \xrightarrow{\cong} s'(gs) \xrightarrow{s'\psi} s'(s'g) \xrightarrow{\cong} (s's') g \xrightarrow{\mu g} s'g \right). \end{aligned} \quad (4.7)$$

The vertical composite of the 2-cells $\rho : (f, \phi) \rightarrow (g, \psi)$ and $\tau : (g, \psi) \rightarrow (h, \theta)$ is the 2-cell

$$f \xrightarrow{\rho} s'g \xrightarrow{s'\tau} s'(s'h) \xrightarrow{\cong} (s's') h \xrightarrow{\mu h} s'h . \quad (4.8)$$

The horizontal composite of 2-cells $\rho : (f, \phi) \rightarrow (g, \psi) : (A, s) \rightarrow (A', s')$ and $\rho' : (f', \phi') \rightarrow (g', \psi') : (A', s') \rightarrow (A'', s'')$ is the 2-cell

$$\begin{aligned} f'f \xrightarrow{f'\rho} f'(s'g) \xrightarrow{\cong} (f's') g \xrightarrow{\phi' g} (s''f') g \\ \xrightarrow{(s''\rho')g} (s''(s'g')) g \xrightarrow{\cong} (s''s'')(g'g) \xrightarrow{\mu(g'g)} s''(g'g) . \end{aligned} \quad (4.9)$$

Each 2-cell $\sigma : (f, \phi) \longrightarrow (g, \psi)$ in $\text{Mnd}^{\text{op}}(\mathcal{K})$ defines a 2-cell $\rho : (f, \phi) \longrightarrow (g, \psi)$ in $\text{KL}(\mathcal{K})$ via $\rho = \eta g \cdot \sigma$. The associativity and unit isomorphisms for $\text{KL}(\mathcal{K})$ are determined by the condition that we have a strict morphism of bicategories

$$K : \text{Mnd}^{\text{op}}(\mathcal{K}) \longrightarrow \text{KL}(\mathcal{K})$$

which is the identity on objects and morphisms and takes each 2-cell σ to $\eta g \cdot \sigma$.

Henceforth we shall invoke the coherence theorem (see [15] and [5]) that every bicategory is biequivalent to a 2-category to write as if we were working in a 2-category $\text{KL}(\mathcal{K})$. We also recommend reworking the proofs below using the string diagrams of [6] as adapted for bicategories in [19] and [20].

Now we are in a position to examine what is involved in an adjunction

$$(f, \phi) \dashv (u, v) : (A, s) \longrightarrow (X, t) \quad (4.10)$$

with counit $\alpha : (f, \phi) \cdot (u, v) \longrightarrow 1_{(A, s)}$ and unit $\beta : 1_{(X, t)} \longrightarrow (u, v) \cdot (f, \phi)$ in $\text{KL}(\mathcal{K})$.

We have morphisms $u : A \longrightarrow X$ and $f : X \longrightarrow A$ in \mathcal{K} . We have 2-cells $v : uf \longrightarrow tu$ and $\phi : ft \longrightarrow sf$ both satisfying (4.3) and (4.4) with the variables appropriately substituted.

We have a 2-cell $\alpha : fu \longrightarrow s$ satisfying

$$\left(fus \xrightarrow{\alpha s} ss \xrightarrow{\mu} s \right) = \left(fus \xrightarrow{fv} ftu \xrightarrow{\phi u} sfu \xrightarrow{s\alpha} ss \xrightarrow{\mu} s \right) \quad (4.11)$$

which is (4.7) for α .

We have a 2-cell $\beta : 1_X \longrightarrow tuf$ satisfying

$$\left(t \xrightarrow{t\beta} ttuf \xrightarrow{\mu u f} tuf \right) = \left(t \xrightarrow{\beta t} tuft \xrightarrow{tu\phi} tusf \xrightarrow{tvf} ttuf \xrightarrow{\mu u f} tuf \right) \quad (4.12)$$

which is (4.7) for β .

Using the rules for compositions in $\text{KL}(\mathcal{K})$, we see that the two triangle conditions for the counit and unit of an adjunction become, in this case, the identities

$$\left(f \xrightarrow{\eta f} sf \right) = \left(f \xrightarrow{f\beta} ftuf \xrightarrow{\phi u f} sfuf \xrightarrow{s\alpha f} ssf \xrightarrow{\mu f} sf \right) \quad (4.13)$$

and

$$\left(u \xrightarrow{\eta u} tu \right) = \left(u \xrightarrow{\beta u} tufu \xrightarrow{tu\alpha} tus \xrightarrow{tv} ttu \xrightarrow{\mu u} tu \right) . \quad (4.14)$$

It is common to call a morphism $f : X \longrightarrow A$ in a bicategory \mathcal{K} a *map* when it has a right adjoint. We write $f^* : A \longrightarrow X$ for a selected right adjoint, $\eta_f : 1_X \longrightarrow f^*f$ for the unit, and $\varepsilon_f : ff^* \longrightarrow 1_A$ for the counit.

4.1. THEOREM. *Suppose (4.10) is an adjunction in $\text{KL}(\mathcal{K})$ with counit α and unit β , and suppose $f : X \longrightarrow A$ is a map in \mathcal{K} . Then the composite 2-cell*

$$\pi : f^*s \xrightarrow{\beta f^*s} tuf f^*s \xrightarrow{tu\varepsilon_f s} tus \xrightarrow{tv} ttu \xrightarrow{\mu u} tu \quad (4.15)$$

in \mathcal{K} is invertible with inverse defined by the composite 2-cell

$$\pi^{-1} : tu \xrightarrow{\eta_f tu} f^*ftu \xrightarrow{f^*\phi u} f^*sfu \xrightarrow{f^*s\alpha} f^*ss \xrightarrow{f^*\mu} f^*s. \quad (4.16)$$

PROOF. Without yet knowing that π^{-1} as given by (4.16) is inverse to π , we calculate $\pi\pi^{-1}$:

$$\begin{aligned}
& \mu u \cdot tv \cdot tu\varepsilon_f s \cdot \beta f^* s \cdot f^* \mu \cdot f^* s \alpha \cdot f^* \phi u \cdot \eta_f t u \\
= & \mu u \cdot tv \cdot t\mu \cdot tus\alpha \cdot tu\phi u \cdot tu\varepsilon_f t u \cdot tuf\eta_f t u \cdot \beta t u \\
= & \mu u \cdot tv \cdot t\mu \cdot tus\alpha \cdot tu\phi u \cdot \beta t u \\
= & \mu u \cdot \mu t u \cdot t t v \cdot t v s \cdot tus\alpha \cdot tu\phi u \cdot \beta t u \\
= & \mu u \cdot tv \cdot t u \alpha \cdot \mu u f u \cdot t v f u \cdot tu\phi u \cdot \beta t u \\
= & \mu u \cdot tv \cdot t u \alpha \cdot \mu u f u \cdot t \beta u \\
= & \mu u \cdot tv \cdot \mu u s \cdot t t u \alpha \cdot t \beta u \\
= & \mu u \cdot t \mu u \cdot t t v \cdot t t u \alpha \cdot t \beta u \\
= & \mu u \cdot t \eta u \\
= & 1_{t u}.
\end{aligned}$$

The first, fourth, sixth and seventh equalities above follow purely from properties of composition in \mathcal{K} . The second equality uses the triangular equation appropriate to the unit and counit for f and its right adjoint. The third equality uses the opmorphism property of (u, v) and associativity of μ . The fifth equality uses (4.12). The eighth equality uses (4.14).

Now we calculate $\pi^{-1}\pi$:

$$\begin{aligned}
& f^* \mu \cdot f^* s \alpha \cdot f^* \phi u \cdot \eta_f t u \cdot \mu u \cdot tv \cdot tu\varepsilon_f s \cdot \beta f^* s \\
= & f^* \mu \cdot f^* s \mu \cdot f^* s s \alpha \cdot f^* s \phi u \cdot f^* \phi t u \cdot \eta_f t t u \cdot tv \cdot tu\varepsilon_f s \cdot \beta f^* s \\
= & f^* \mu \cdot f^* \mu s \cdot f^* s \alpha s \cdot f^* \phi u s \cdot \eta_f t u s \cdot t u \eta_f s \beta f^* s \\
= & f^* \mu \cdot f^* s \varepsilon_f s \cdot f^* \eta f f^* s \cdot \eta_f f^* s \\
= & 1_{f^* s},
\end{aligned}$$

using the associativity and unit conditions for the monads, the opmorphism property of (f, ϕ) , equation (4.11), and equation (4.14). ■

As expected by general principles of doctrinal adjunction [9], a monad opmorphism $(f, \phi) : (X, t) \rightarrow (A, s)$ for which f is a map in \mathcal{K} gives rise to a monad morphism $(f^*, \hat{\phi}) : (A, s) \rightarrow (X, t)$ where $\hat{\phi} : t f^* \rightarrow f^* s$ is the mate of ϕ under the adjunction $f \dashv f^*$ in the sense of [11].

4.2. REMARK. *The Kleisli construction for \mathcal{K} is a left biadjoint for the pseudofunctor $\mathcal{K} \rightarrow \text{KL}(\mathcal{K})$ taking A to $(A, 1_A)$. The value of the biadjoint at (A, t) is denoted by A_t . When the Kleisli construction exists, it takes the adjunction (4.10) in $\text{KL}(\mathcal{K})$ to an adjunction $f_\phi \dashv u_v : A_s \rightarrow X_t$ in \mathcal{K} .*

5. Cores between monads

5.1. DEFINITION. An adjunction core (u, g, π) between monads (A, s) and (X, t) in a bicategory \mathcal{K} consists of the following data in \mathcal{K} :

1. morphisms $u : A \rightarrow X$ and $g : A \rightarrow X$;
2. an invertible 2-cell $\pi : gs \rightarrow tu$.

Given such a core, we make the following definitions:

(a) $\bar{\beta} : g \rightarrow tu$ is the composite

$$g \xrightarrow{g\eta} gs \xrightarrow{\pi} tu;$$

(b) $\bar{\alpha} : u \rightarrow gs$ is the composite

$$u \xrightarrow{\eta u} tu \xrightarrow{\pi^{-1}} gs;$$

(c) $v : us \rightarrow tu$ is the composite

$$us \xrightarrow{\bar{\alpha}s} gss \xrightarrow{g\mu} gs \xrightarrow{\pi} tu;$$

(d) $\psi : tg \rightarrow gs$ is the composite

$$tg \xrightarrow{t\bar{\beta}} ttu \xrightarrow{\mu u} tu \xrightarrow{\pi^{-1}} gs.$$

5.2. PROPOSITION. An adjunction core between monads (A, s) and (X, t) is obtained from the data of Theorem 4.1 by putting $g = f^*$. Moreover, the $\bar{\beta}$ of (a) and the $\bar{\alpha}$ of (b) are the mates of the unit β and counit α , respectively, the composite in (c) recovers v , and the ψ of (d) is the mate $\hat{\phi}$ of ϕ .

PROOF. That we have an adjunction core follows from the invertibility of π according to Theorem 4.1. Next we look at the composite in (a):

$$\begin{aligned} & \pi \cdot f^* \eta \\ &= \mu u \cdot tv \cdot tu \varepsilon_f s \cdot \beta f^* s \cdot f^* \eta \\ &= \mu u \cdot tv \cdot t\eta \cdot tu \varepsilon_f \cdot \beta f^* \\ &= \mu u \cdot t\eta u \cdot tu \varepsilon_f \cdot \beta f^* \\ &= tu \varepsilon_f \cdot \beta f^* \end{aligned}$$

which is the mate $\bar{\beta}$ of β . That the composite in (b) gives $\bar{\alpha}$ is a similar calculation.

Next we calculate:

$$\begin{aligned} & \pi^{-1} s \cdot \eta us \\ &= f^* \mu s \cdot f^* s \alpha s \cdot f^* \phi us \cdot \eta_f tus \cdot \eta us \\ &= f^* \mu s \cdot f^* s \alpha s \cdot f^* \phi us \cdot f^* f \eta_f us \cdot \eta_f us \\ &= f^* \mu s \cdot f^* s \alpha s \cdot f^* \eta_f us \cdot \eta_f us \\ &= f^* \mu s \cdot f^* \eta_f ss \cdot f^* \alpha s \cdot \eta_f us \\ &= f^* \alpha s \cdot \eta_f us, \end{aligned}$$

so the composite in (c) is:

$$\begin{aligned}
 & \mu u \cdot tv \cdot tu\varepsilon_f s \cdot \beta f^* s a \cdot f^* \mu \cdot \pi^{-1} s \cdot \eta u s \\
 = & \mu u \cdot tv \cdot tu\varepsilon_f s \cdot \beta f^* s \cdot f^* \mu \cdot f^* \alpha s \cdot \eta_f u s \\
 = & \mu u \cdot tv \cdot tu\varepsilon_f s \cdot tu f f^* \mu \cdot \beta f^* s s \cdot f^* \alpha s \cdot \eta_f u s \\
 = & \mu u \cdot tv \cdot tu\varepsilon_f s \cdot tu f f^* \mu \cdot tu f f^* \alpha s \cdot \beta f^* f u s \cdot \eta_f u s \\
 = & \mu u \cdot tv \cdot t\mu \cdot tu\varepsilon_f s s \cdot tu f f^* \alpha s \cdot \beta f^* f u s \cdot \eta_f u s \\
 = & \mu u \cdot t\mu u \cdot ttv \cdot tvs \cdot tu\varepsilon_f s s \cdot tu f f^* \alpha s \cdot \beta f^* f u s \cdot \eta_f u s \\
 = & \mu u \cdot t\mu u \cdot ttv \cdot tvs \cdot tu\alpha s \cdot tu\varepsilon_f f u s \cdot \beta f^* f u s \cdot \eta_f u s \\
 = & \mu u \cdot t\mu u \cdot ttv \cdot tvs \cdot tu\alpha s \cdot tu\varepsilon_f f u s \cdot tu f \eta_f s \cdot \beta u s \\
 = & \mu u \cdot t\mu u \cdot ttv \cdot tvs \cdot tu\alpha s \cdot \beta u s \\
 = & \mu u \cdot tv \cdot \mu u s \cdot tvs \cdot tu\alpha s \cdot \beta u s \\
 = & \mu u \cdot tv \cdot \eta u s \\
 = & \mu u \cdot \eta t u \cdot v \\
 = & v,
 \end{aligned}$$

as required. The calculation for the composite in (d) is similar. \blacksquare

The Corollary of the following result should be compared with Theorem 3.2.

5.3. THEOREM. *For an adjunction core (u, g, π) between monads (A, s) and (X, t) in a bicategory \mathcal{K} , the following two commutativity conditions (5.1) and (5.2) are equivalent.*

$$\begin{array}{ccc}
 & tus & \\
 \pi s \nearrow & & \searrow tv \\
 gss & & ttu \\
 g\mu \downarrow & & \downarrow \mu u \\
 gs & \xrightarrow{\pi} & tu
 \end{array} \tag{5.1}$$

$$\begin{array}{ccc}
 & tgs & \\
 t\pi^{-1} \nearrow & & \searrow \psi s \\
 ttu & & gss \\
 \mu u \downarrow & & \downarrow g\mu \\
 tu & \xrightarrow{\pi^{-1}} & gs
 \end{array} \tag{5.2}$$

Moreover, under these conditions, using definitions (a), (b), (c) and (d),

- (i) $(u, v) : (A, s) \longrightarrow (X, t)$ is a monad opmorphism;
- (ii) $(g, \psi) : (A, s) \longrightarrow (X, t)$ is a monad morphism;

(iii) π is equal to the composite

$$gs \xrightarrow{\bar{\beta}s} tus \xrightarrow{tv} ttu \xrightarrow{\mu u} tu;$$

(iv) π^{-1} is equal to the composite

$$tu \xrightarrow{t\bar{\alpha}} tgs \xrightarrow{\psi s} gss \xrightarrow{g\mu} gs;$$

(v) the following identity holds

$$(us \xrightarrow{\bar{\alpha}s} gss \xrightarrow{g\mu} gs) = (us \xrightarrow{v} tu \xrightarrow{t\bar{\alpha}} tgs \xrightarrow{\psi s} gss \xrightarrow{g\mu} gs);$$

(vi) the following identity holds

$$(tg \xrightarrow{t\bar{\beta}} ttu \xrightarrow{\mu u} tu) = (tg \xrightarrow{\psi} gs \xrightarrow{\bar{\beta}s} tus \xrightarrow{tv} ttu \xrightarrow{\mu u} tu);$$

(vii) the following identity holds

$$(g \xrightarrow{g\eta} gs) = (g \xrightarrow{\bar{\beta}} tu \xrightarrow{t\bar{\alpha}} tgs \xrightarrow{\psi s} gss \xrightarrow{g\mu} gs);$$

(viii) the following identity holds

$$(u \xrightarrow{\eta u} tu) = (u \xrightarrow{\bar{\alpha}} gs \xrightarrow{\bar{\beta}s} tus \xrightarrow{tv} ttu \xrightarrow{\mu u} tu).$$

PROOF. Assuming (5.1) at the first step, we have the calculation:

$$\begin{aligned} & \pi \cdot g\mu \cdot \psi s \\ &= \mu u \cdot tv \cdot \pi s \cdot \psi s \\ &= \mu u \cdot tv \cdot \pi s \cdot \pi^{-1} s \cdot \mu us \cdot t\bar{\beta}s \\ &= \mu u \cdot tv \cdot \mu us \cdot t\bar{\beta}s \\ &= \mu u \cdot \mu tu \cdot ttv \cdot t\bar{\beta}s \\ &= \mu u \cdot t\mu u \cdot ttv \cdot t\bar{\beta}s \\ &= \mu u \cdot t\pi, \end{aligned}$$

proving (5.2). The converse is dual.

(i) Using (5.1) at the second step, we have the calculation:

$$\begin{aligned} & \mu u \cdot tv \cdot vs \\ &= \mu u \cdot tv \cdot \pi s \cdot g\mu s \cdot \bar{\alpha}ss \\ &= \pi \cdot g\mu \cdot g\mu s \cdot \bar{\alpha}ss \\ &= \pi \cdot g\mu \cdot gs\mu \cdot \bar{\alpha}ss \\ &= \pi \cdot g\mu \cdot \bar{\alpha}s \cdot u\mu \\ &= v \cdot u\mu. \end{aligned}$$

We also have:

$$\begin{aligned}
& v \cdot u\eta \\
&= \pi \cdot g\mu \cdot \bar{\alpha}s \cdot u\eta \\
&= \pi \cdot g\mu \cdot \pi^{-1}s \cdot \eta us \cdot u\eta \\
&= \pi \cdot g\mu \cdot \pi^{-1}s \cdot tu\eta \cdot \eta u \\
&= \pi \cdot g\mu \cdot g\eta \cdot \pi^{-1} \cdot \eta u \\
&= \pi \cdot \pi^{-1} \cdot \eta u \\
&= \eta u.
\end{aligned}$$

Hence (u, v) is a monad opmorphism.

(ii) This is dual to (i) using (5.2) instead of (5.1).

(iii) Using (5.1), we have:

$$\begin{aligned}
& \pi \\
&= \pi \cdot g\mu \cdot g\eta s \\
&= \mu u \cdot tv \cdot \pi s \cdot g\eta s \\
&= \mu u \cdot tv \cdot \bar{\beta}s.
\end{aligned}$$

(iv) This is dual to (iii) using (5.2) instead of (5.1).

(v) Using (iv), we immediately have:

$$\begin{aligned}
& g\mu \cdot \psi s \cdot t\bar{\alpha} \cdot v \\
&= \pi^{-1} \cdot v \\
&= \pi^{-1} \cdot \pi \cdot g\mu \cdot \bar{\alpha}s \\
&= g\mu \cdot \bar{\alpha}s.
\end{aligned}$$

(vi) This is dual to (v) using (iii) instead of (iv).

(vii) Using (iv), we immediately have:

$$\begin{aligned}
& g\mu \cdot \psi s \cdot t\bar{\alpha} \cdot \bar{\beta} \\
&= \pi^{-1} \cdot \bar{\beta} \\
&= \pi^{-1} \cdot \pi \cdot g\eta \\
&= g\eta.
\end{aligned}$$

(viii) This is dual to (vii) using (iii) instead of (iv). ■

5.4. COROLLARY. *An adjunction core of the form (u, f^*, π) between monads (A, s) and (X, t) in a bicategory \mathcal{K} extends to an adjunction (4.10) in $\text{KL}(\mathcal{K})$ if and only if one of the diagrams (5.1) or (5.2) commutes. The adjunction is unique when it exists.*

PROOF. Properties (i)–(viii) of Theorem 5.3, when re-expressed with the mates α, β and ϕ replacing $\bar{\alpha}, \bar{\beta}$ and ψ , give precisely what is required for an adjunction (4.10). The converse is Proposition 5.2. ■

5.5. **REMARK.** Recall from [13] that the free completion $\text{EM}(\mathcal{K})$ of \mathcal{K} with respect to the Eilenberg–Moore construction is obtained as

$$\text{EM}(\mathcal{K}) = \text{KL}(\mathcal{K}^{\text{op}})^{\text{op}} .$$

Monads in \mathcal{K}^{op} are the same as monads in \mathcal{K} . An adjunction opcore between monads (X, t) and (A, s) in \mathcal{K} is an adjunction core in \mathcal{K}^{op} . This consists of morphisms $f, v : X \rightarrow A$ and an invertible 2-cell $\pi : sv \cong ft$. Also recall from [13] that a wreath in \mathcal{K} is a monad in $\text{EM}(\mathcal{K})$. Many examples of wreaths were provided in that paper. A wreath on A consists of a monad $t : A \rightarrow A$ with unit $\eta : 1 \Rightarrow t$ and multiplication $\mu : tt \Rightarrow t$, an endomorphism $s : A \rightarrow A$ with 2-cells $\sigma : 1 \Rightarrow st$ and $\nu : ss \Rightarrow st$, and a 2-cell $\lambda : ts \Rightarrow st$ satisfying conditions. This gives an adjunction cocore $s, 1_A : A \rightarrow A$ between the monad (A, s) and the wreath product monad (A, st) where π is the identity of st . Since 1_A has itself as adjoint and condition (5.1) can be checked, we obtain an example of an adjunction between monads from every wreath. To be explicit, we have the adjunction

$$(s, s\mu.\nu t.s\lambda) \dashv (1_A, s\mu.\sigma t) : (A, st) \rightarrow (A, t)$$

associated with the Eilenberg–Moore construction for the monad (s, λ) on (A, t) in $\text{EM}(\mathcal{K})$.

6. Cores between internal categories

This section will apply our results to categories internal to a category \mathcal{C} which admits pullbacks. For this example, we take the bicategory \mathcal{K} of the previous sections to be the bicategory $\text{Span}(\mathcal{C})$ of spans in \mathcal{C} as constructed by Bénabou in [1].

The objects of the bicategory $\text{Span}(\mathcal{C})$ are those of \mathcal{C} . A morphism $S = (s_0, S, s_1) : U \rightarrow V$ is a so-called span

$$U \xleftarrow{s_0} S \xrightarrow{s_1} V$$

from U to V in \mathcal{C} . A 2-cell $r : (s_0, S, s_1) \rightarrow (t_0, T, t_1) : U \rightarrow V$ is a morphism $r : S \rightarrow T$ in \mathcal{C} such that $t_0 r = s_0$ and $t_1 r = s_1$. Vertical composition of 2-cells is simply that of \mathcal{C} . Horizontal composition uses pullback in \mathcal{C} ; more precisely,

$$(U \xrightarrow{(s_0, S, s_1)} V \xrightarrow{(t_0, T, t_1)} W) = (U \xrightarrow{(s_0 p, P, t_1 q)} W)$$

where

$$\begin{array}{ccc} P & \xrightarrow{q} & T \\ p \downarrow & & \downarrow t_0 \\ S & \xrightarrow{s_1} & V \end{array} \quad (6.1)$$

is a pullback square.

Each morphism $f : U \rightarrow V$ in \mathcal{C} determines a span $f_\star = (1_U, U, f) : U \rightarrow V$. We write $f^\star : V \rightarrow U$ for the span $(f, V, 1_V) : V \rightarrow U$. It is well known that we have an

adjunction $f_* \dashv f^*$ in $\text{Span}(\mathcal{C})$; in fact, it is shown in [3] that the maps in $\text{Span}(\mathcal{C})$ are all isomorphic to spans of the form $f_* : U \rightarrow V$ for some $f : U \rightarrow V$ in \mathcal{C} .

One of the reasons for interest in the free Kleisli object cocompletion $\text{KL}(\mathcal{K})$ in the paper [13] is that the 2-category $\text{Cat}(\mathcal{C})$ of categories in \mathcal{C} is equivalent to the sub-2-category of $\text{KL}(\text{Span}(\mathcal{C}))$ obtained by restricting to the morphisms whose underlying morphisms in $\text{Span}(\mathcal{C})$ are maps. We shall explain this in a bit more detail.

A *category* in \mathcal{C} is a monad (A, S) in $\text{Span}(\mathcal{C})$. The object A of \mathcal{C} is called the *object of objects*. The span $S = (s_0, S, s_1) : A \rightarrow A$ provides the *object of morphisms* S , the *source operation* s_1 and the *target operation* s_0 . The multiplication for the monad provides the *composition operation* and the unit for the monad provides the *identities operation*.

A *functor* between categories in \mathcal{C} is a monad opmorphism of the form $(f_*, \phi) : (X, T) \rightarrow (A, S)$ in $\text{Span}(\mathcal{C})$. The morphism $f : X \rightarrow A$ in \mathcal{C} is called the *effect on objects* of the functor and the morphism $\phi : T \rightarrow S$ in \mathcal{C} is called the *effect on morphisms* of the functor.

A *natural transformation* between functors in \mathcal{C} is precisely a 2-cell between them in $\text{KL}(\text{Span}(\mathcal{C}))$.

As an immediate consequence of Corollary 5.4 we have:

6.1. COROLLARY. *An adjunction core of the form (u_*, f^*, π) between categories (A, S) and (X, T) in a category \mathcal{C} extends to an adjunction $(f_*, \phi) \dashv (u_*, v)$ in $\text{Cat}(\mathcal{C})$ if and only if one of the diagrams (5.1) or (5.2) mutatis mutandis commutes. The adjunction is unique when it exists.*

6.2. REMARK. *If in Corollary 6.1 we denote the categories (A, S) and (X, T) in \mathcal{C} by \mathbb{A} and \mathbb{X} and the functors (f_*, ϕ) and (u_*, v) by $F : \mathbb{X} \rightarrow \mathbb{A}$ and $U : \mathbb{A} \rightarrow \mathbb{X}$ then the π is an isomorphism of spans from A to X between the object of objects of the comma category F/\mathbb{A} and the object of objects of the comma category \mathbb{X}/U . We leave it to the reader to interpret diagram (5.1) in this notation.*

7. A doctrinal setting

Let D be a pseudomonad (also called a doctrine in [14], [9], [21] and [18]) on a bicategory \mathcal{K} . It means that we have a pseudofunctor $D : \mathcal{K} \rightarrow \mathcal{K}$, a unit pseudonatural transformation denoted by $n : 1_{\mathcal{K}} \rightarrow D$, and a multiplication pseudonatural transformation denoted by $m : DD \rightarrow D$. For example, see [16] or [12] for the axioms.

A *lax D -algebra* (A, s) consists of an object A , a morphism $s : DA \rightarrow A$, and 2-cells $\mu : s \cdot Ds \Longrightarrow s \cdot m_A$ and $\eta : 1_A \Longrightarrow s \cdot n_A$, satisfying coherence conditions.

For lax D -algebras (A, s) and (A', s') in \mathcal{K} , a *lax opmorphism* $(f, \phi) : (A, s) \rightarrow (A', s')$ consists of a morphism $f : A \rightarrow A'$ and a 2-cell $\phi : fs \Longrightarrow s'Df$ in \mathcal{K} such that

$$\begin{array}{ccc}
& s'Df \cdot Ds \xrightarrow{s'D\phi} s'Ds' \cdot D^2f & \\
\phi Ds \nearrow & & \searrow \mu D^2f \\
fsDs & & s'm_{A'}D^2f \\
f\mu \searrow & & \nearrow s'm_f \\
& fsm_A \xrightarrow{\phi m_A} s'Df \cdot m_A &
\end{array} \tag{7.1}$$

$$\begin{array}{ccc}
f s n_A & \xrightarrow{\phi n_A} & s' D f \cdot n_A \\
\uparrow f \eta & & \downarrow s' n_f \\
f & \xrightarrow{\eta f} & s' n_{A'}
\end{array} \tag{7.2}$$

The composite of lax opmorphisms $(f, \phi) : (A, s) \longrightarrow (A', s')$ and $(f', \phi') : (A', s') \longrightarrow (A'', s'')$ is defined to be $(f'f, \phi' \star \phi) : (A, s) \longrightarrow (A'', s'')$ where $\phi' \star \phi$ is the composite

$$f'f s \xrightarrow{f'\phi} f's'Df \xrightarrow{\phi'Df} s''Df' \cdot Df \ . \tag{7.3}$$

There is a bicategory $\text{KL}(\mathcal{K}, D)$ whose objects are lax D -algebras and whose morphisms are lax opmorphisms. A 2-cell $\rho : (f, \phi) \Longrightarrow (g, \psi) : (A, s) \longrightarrow (A', s')$ in $\text{KL}(\mathcal{K}, D)$ (in non-reduced form, in the terminology of 13) is a 2-cell $\hat{\rho} : fs \Longrightarrow s'Dg$ in \mathcal{K} such that the diagrams (7.4) and (7.5) commute.

$$\begin{array}{ccc}
& s'Dg \cdot Ds \xrightarrow{s'D\psi} s'Ds'D^2g & \\
\hat{\rho}Ds \nearrow & & \searrow \mu D^2g \\
fsDs & & s'm_{A'}D^2g \\
f\mu \searrow & & \nearrow s'm_g \\
& fsm_A \xrightarrow{\hat{\rho}m_A} s'Dg \cdot m_A &
\end{array} \tag{7.4}$$

$$\begin{array}{ccc}
& s'Df \cdot Ds \xrightarrow{s'D\hat{\rho}} s'Ds'D^2g & \\
\phi Ds \nearrow & & \searrow \mu D^2g \\
fsDs & & s'm_{A'}D^2g \\
f\mu \searrow & & \nearrow s'm_g \\
& fsm_A \xrightarrow{\hat{\rho}m_A} s'Dg \cdot m_A &
\end{array} \tag{7.5}$$

The reduced form of the 2-cell $\text{KL}(\mathcal{K}, D)$ is a 2-cell $\rho : f \Longrightarrow s'Dg \cdot n_A$ in \mathcal{K} such that equality (7.6) holds.

$$\begin{aligned}
f s \xrightarrow{\phi} s' D f \xrightarrow{s' D \rho} s' D s' D^2 g D n_A \xrightarrow{\mu D^2 g D n_A} s' m_{A'} D^2 g D n_A \\
\xrightarrow{s' m_g^{-1} D n_A} s' D g \cdot m_A D n_A \xrightarrow{\cong} s' D g \\
= \\
f s \xrightarrow{\rho s} s' D g \cdot n_A s \xrightarrow{s' n_g s} s' n_{A'} g s \xrightarrow{s' n_{A'} \psi} s' n_{A'} s' D g \\
\xrightarrow{s' n_{s'}^{-1} D g} s' D s' n_{D A'} D g \xrightarrow{\mu n_{D A'} D g} s' m_{A'} n_{D A'} D g \xrightarrow{\cong} s' D g. \quad (7.6)
\end{aligned}$$

The bijection between reduced and non-reduced forms takes ρ to the $\hat{\rho}$ defined by either side of (7.6).

The vertical composite of the 2-cells $\rho : (f, \phi) \longrightarrow (g, \psi)$ and $\tau : (g, \psi) \longrightarrow (h, \theta)$ is the 2-cell

$$f' f s \xrightarrow{f' \hat{\rho}} f' s' D g \xrightarrow{s' \tau} s' (s' h) \xrightarrow{\cong} (s' s') h \xrightarrow{\mu h} s' h. \quad (7.7)$$

The horizontal composite of 2-cells $\rho : (f, \phi) \longrightarrow (g, \psi) : (A, s) \longrightarrow (A', s')$ and $\rho' : (f', \phi') \longrightarrow (g', \psi') : (A', s') \longrightarrow (A'', s'')$ has non-reduced form the 2-cell

$$\begin{aligned}
f \xrightarrow{\rho} s' D g \cdot n_A \xrightarrow{s' D \tau \cdot n_A} s' D s' D^2 h D n_A \cdot n_A \xrightarrow{\mu D^2 h D n_A \cdot n_A} s' m_{A'} D^2 h D n_A \cdot n_A \\
\xrightarrow{s' m_h^{-1} D n_A \cdot n_A} s' D h \cdot m_A D n_A \cdot n_A \xrightarrow{\cong} s' D h \cdot n_A. \quad (7.8)
\end{aligned}$$

This completes the definition of the bicategory $\text{KL}(\mathcal{K}, D)$ with the exception of giving the coherent associativity and unit isomorphisms. As forewarned, we have been writing as if \mathcal{K} were a 2-category in which case $\text{KL}(\mathcal{K}, D)$ would also be a 2-category. Putting in all the coherent isomorphisms as we did in the definition of $\text{KL}(\mathcal{K})$, we can readily give them for $\text{KL}(\mathcal{K}, D)$ as we did for the special case where D was the identity pseudomonad.

In a future paper we shall explain the universal properties of the construction taking \mathcal{K} to $\text{KL}(\mathcal{K})$.

7.1. DEFINITION. *An adjunction core (u, g, π) between lax D -algebras (A, s) and (X, t) in a bicategory \mathcal{K} consists of the following data in \mathcal{K} :*

1. *morphisms $u : A \longrightarrow X$ and $g : A \longrightarrow X$;*
2. *an invertible 2-cell $\pi : g s \longrightarrow t D u$.*

Given such a core, we make the following definitions:

(a) $\bar{\beta} : g \Longrightarrow t D u \cdot n_A$ is the composite

$$g \xrightarrow{g \eta} g s n_A \xrightarrow{\pi n_A} t D u \cdot n_A ;$$

(b) $\bar{\alpha} : u \Longrightarrow g s n_A$ is the composite

$$u \xrightarrow{\eta u} t n_X u \xrightarrow{\cong} t D u \cdot n_A \xrightarrow{\pi^{-1} n_A} g s n_A ;$$

(c) $v : us \Longrightarrow tDu$ is the composite

$$us \xrightarrow{\bar{\alpha}s} gsn_{AS} \xrightarrow{gsn_s} gsDs \cdot n_{DA} \xrightarrow{g\mu n_{DA}} gsm_{AN_{DA}} \xrightarrow{\cong} gs \xrightarrow{\pi} tDu \ ;$$

(d) $\psi : tDg \Longrightarrow gs$ is the composite

$$tDg \xrightarrow{tD\bar{\beta}} tDtD^2uDn_A \xrightarrow{\mu n_u^{-1}} tm_X Dn_X Du \xrightarrow{\cong} tDu \xrightarrow{\pi^{-1}} gs \ .$$

If $g = f^*$ for some map $f : X \longrightarrow A$ in \mathcal{K} then $\bar{\alpha}$ and $\bar{\beta}$ have mates $\alpha : fu \Longrightarrow sn_A$ and $\beta' : 1_X \Longrightarrow tDu \cdot n_A f$. We take $\beta : 1_X \Longrightarrow tDu Df \cdot n_X$ to be the composite of β' and $tn_f : tDu \cdot n_A f \cong tDu Df \cdot n_X$.

7.2. THEOREM. *An adjunction core of the form (u, f^*, π) between lax D -algebras (A, s) and (X, t) in a bicategory \mathcal{K} extends to an adjunction*

$$(f, \phi) \dashv (u, v) : (A, s) \longrightarrow (X, t) \quad (7.9)$$

with counit α and unit β in $\text{KL}(\mathcal{K}, D)$ if and only if one of the diagrams (7.10) or (7.11) commutes. The adjunction is unique when it exists.

$$\begin{array}{ccc} & tDuDs & \xrightarrow{tDv} tDtD^2u \\ \pi Ds \nearrow & & \searrow \mu D^2u \\ gsDs & & tm_X D^2u \\ g\mu \searrow & & \nearrow tm_u \\ & gsm_A \xrightarrow{\pi m_A} tDu \cdot m_A & \end{array} \quad (7.10)$$

$$\begin{array}{ccc} & tDgDs & \xrightarrow{\psi Ds} gsDs \\ tD\pi^{-1} \nearrow & & \searrow g\mu \\ tDtD^2u & & gsm_A \\ \mu D^2u \searrow & & \nearrow \pi^{-1} m_A \\ & tm_X D^2u \xrightarrow{tm_u^{-1}} tDu \cdot m_A & \end{array} \quad (7.11)$$

References

- [1] Jean Bénabou, *Introduction to bicategories*, Lecture Notes in Mathematics **47** (Springer-Verlag, 1967) 1–77.
- [2] Renato Betti, Aurelio Carboni, Ross Street and Robert Walters, *Variation through enrichment*, Journal of Pure and Applied Algebra **29** (1983) 109–127.

- [3] Aurelio Carboni, Stefano Kasangian and Ross Street, *Bicategories of spans and relations*, Journal of Pure and Applied Algebra **33** (1984) 259–267.
- [4] Samuel Eilenberg and G. Max Kelly, *Closed categories*, Proceedings of the Conference on Categorical Algebra (La Jolla, 1965), Springer (1966) 421–562.
- [5] Robert Gordon, A. John Power and Ross Street, *Coherence for tricategories*, Memoirs of the American Math. Society **117** no. 558 (1995) vi+81 pp.
- [6] André Joyal and Ross Street, *The geometry of tensor calculus*, Advances in Math. **88** (1991) 55–112.
- [7] Daniel M. Kan, *Adjoint functors*, Transactions of the American Mathematical Society **87** (1958) 294–329.
- [8] G. Max Kelly, *Adjunction for enriched categories*, Lecture Notes in Mathematics **106** (Springer-Verlag, 1969) 166–177.
- [9] G. Max Kelly, *Doctrinal adjunction*, Lecture Notes in Mathematics **420** (Springer-Verlag, 1974) 257–280.
- [10] G. Max Kelly, *Basic concepts of enriched category theory*, London Mathematical Society Lecture Note Series **64** (Cambridge University Press, Cambridge, 1982).
- [11] G. Max Kelly and Ross Street, *Review of the elements of 2-categories*, Lecture Notes in Mathematics **420** (Springer-Verlag, 1974) 75–103.
- [12] Stephen Lack, *A coherent approach to pseudomonads*, Advances in Math. **152** (2000) 179–202.
- [13] Stephen Lack and Ross Street, *The formal theory of monads II*, Journal of Pure and Applied Algebra **175** (2002) 243–265.
- [14] F. William Lawvere, *Ordinal sums and equational doctrines*, Lecture Notes in Mathematics **80** (Springer-Verlag, 1969) 141–155.
- [15] Saunders Mac Lane and Robert Paré, *Coherence for bicategories and indexed categories*, Journal of Pure and Applied Algebra **37** (1985) 59–80.
- [16] Francisco Marmolejo, *Distributive laws for pseudomonads*, Theory and Applications of Categories **5** (1999) 91–147.
- [17] Ross Street, *The formal theory of monads*, Journal of Pure and Applied Algebra **2** (1972) 149–168.
- [18] Ross Street, *Fibrations in bicategories*, Cahiers de topologie et géométrie différentielle **21** (1980) 111–160.

- [19] Ross Street, *Higher categories, strings, cubes and simplex equations*, Applied Categorical Structures **3** (1995) 29–77.
- [20] Ross Street, *Categorical structures*, Handbook of Algebra **1** (edited by M. Hazewinkel, Elsevier Science, 1996) 529–577.
- [21] Volker Zöberlein, *Doctrines on 2-categories*, Math. Zeitschrift **148** (1976) 267–279.

Department of Mathematics, Macquarie University, NSW 2109, Australia

Email: `ross.street@mq.edu.au`

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/27/4/27-04.{dvi,ps,pdf}>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is \TeX , and $\text{\LaTeX}2\epsilon$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

\TeX NICAL EDITOR Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT \TeX EDITOR Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS

Clemens Berger, Université de Nice-Sophia Antipolis, cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva: valeria.depaiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne : kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

P. T. Johnstone, University of Cambridge: ptj@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Glasgow, Tom.Leinster@glasgow.ac.uk

Jean-Louis Loday, Université de Strasbourg: loday@math.u-strasbg.fr

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Edinburgh: Alex.Simpson@ed.ac.uk

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca