

A C-SYSTEM DEFINED BY A UNIVERSE CATEGORY

VLADIMIR VOEVODSKY

ABSTRACT. This is the third paper in a series started in [9]. In it we construct a C-system $CC(\mathcal{C}, p)$ starting from a category \mathcal{C} together with a morphism $p : \tilde{U} \rightarrow U$, a choice of pull-back squares based on p for all morphisms to U and a choice of a final object of \mathcal{C} . Such a quadruple is called a universe category. We then define universe category functors and construct homomorphisms of C-systems $CC(\mathcal{C}, p)$ defined by universe category functors.

In the sections before the last section we give, for any C-system CC , three different constructions of pairs $((\mathcal{C}, p), H)$ where (\mathcal{C}, p) is a universe category and $H : CC \rightarrow CC(\mathcal{C}, p)$ is an isomorphism.

In the last section we construct for any (set) category C with a choice of a final object and fiber products a C-system and an equivalence between C and the precategory underlying CC .

1. Introduction

The concept of a C-system in its present form was introduced in [9]. The type of C-systems is constructively equivalent to the type of contextual categories defined by Cartmell in [4] and [3] but the definition of a C-system is slightly different from Cartmell's foundational definition.

In [8] we constructed for any pair (R, LM) where R is a monad on *Sets* and LM a left R -module with values in *Sets* a C-system $CC(R, LM)$. In the particular case of pairs (R, LM) corresponding to binding signatures (cf. [1], [5], [6, p.228]) the regular sub-quotients of $CC(R, LM)$ are the C-systems corresponding to dependent type theories of Martin-Löf genus.

In this paper we describe another construction that generates C-systems. This time the input data is a quadruple that consists of a category \mathcal{C} , a morphism $p : \tilde{U} \rightarrow U$ in this category, a choice of pull-back squares based on p for all morphisms to U and a choice

1. Work on this paper was supported by NSF grant 1100938 and Clay Mathematical Institute. 2. This material is based on research sponsored by the United States Air Force Research Laboratory under agreement number FA9550-15-1-0053. The US Government is authorized to reproduce and distribute reprints for Governmental purposes notwithstanding any copyright notation thereon. 3. The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing the official policies or endorsements, either expressed or implied, of the United States Air Force Research Laboratory, the US Government or Carnegie Mellon University.

Received by the editors 2015-02-02 and, in revised form, 2015-09-06.

Transmitted by Martin Hyland. Published on 2015-09-17.

2010 Mathematics Subject Classification: 03F50, 03B15, 03G25 .

Key words and phrases: contextual category, universe category, C-system.

© Vladimir Voevodsky, 2015. Permission to copy for private use granted.

of a final object in \mathcal{C} . Such a quadruple is called a universe category. For any universe category we construct a C-system that we denote by $CC(\mathcal{C}, p)$.

We then define the notion of a universe category functor and construct homomorphisms of C-systems of the form $CC(\mathcal{C}, p)$ corresponding to universe category functors. For universe category functors satisfying certain conditions these homomorphisms are isomorphisms. In particular, any equivalence $F : \mathcal{C} \rightarrow \mathcal{C}'$ together with an isomorphism $F(p) \cong p'$ (in the category of morphisms) defines a universe category functor whose associated homomorphism of C-systems is an isomorphism. This implies the C-systems that correspond to two different choices of final objects and pull-backs for the same \mathcal{C} and p are connected by a given isomorphism which justifies our simplified notation $CC(\mathcal{C}, p)$.

To the best of our knowledge it is the only known construction of a C-system (or, equivalently, a contextual category) from category level data that transforms equivalences into isomorphisms. Because of this fact we find it important to present both the construction of the C-system and the construction of the homomorphisms defined by universe functors in detail.

This construction is also important as the starting point in the construction of the model of the inference rules of Martin-Löf type theory that satisfies the univalence axiom. The underlying C-system of the univalent model is obtained by application of our Construction 2.12 to the universal Kan fibration of well-ordered simplicial sets (in a given Grothendieck universe).

Next we explore the question of how to construct, for a given C-system CC , a universe category (\mathcal{C}, p) together with an isomorphism $CC \rightarrow CC(\mathcal{C}, p)$. It is clear from Lemma 4.8 that if this problem has a solution then it has many solutions. We construct three such solutions each having certain advantages and disadvantages.

The set of universe categories in a given Grothendieck universe has a structure of a 2-category suggested by Definition 4.1. It seems likely that our main construction extends to a construction of a functor from this 2-category to the 1-category of C-systems. We leave the investigations of the properties of this 2-category and of this functor for the future.

To avoid the abuse of language inherent in the use of the Theorem-Proof style of presenting mathematics when dealing with constructions we use the pair of names Problem-Construction for the specification of the goal of a construction and the description of the particular solution.

In the case of a Theorem-Proof pair one usually refers (by name or number) to the statement when using both the statement and the proof. This is acceptable in the case of theorems because the future use of their proofs is such that only the fact that there is a proof but not the particulars of the proof matter.

In the case of a Problem-Construction pair the content of the construction often matters in the future use. Because of this we often have to refer to the construction and not to the problem and we assign in this paper numbers both to Problems and to the Constructions.

Following the approach used in [9] we write the composition of morphisms in categories

in the diagrammatic order, i.e., for $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ their composition is written as $f \circ g$. This makes it much easier to translate between diagrams and equations involving morphisms.

The methods of this paper are fully constructive and the style we write in is the “formalization ready” style where the proofs are spelled out in detail even when the assertion may appear obvious to the practitioners of a particular tradition in mathematics. This particular paper is written having in mind the possibility of formalization both in the Zermelo-Fraenkel set theory (without the axiom of choice) and its constructive versions and in any type theory including Church’s type theory or HOL.

Following the distinction that becomes essential in the univalent formalization (cf. [2]) we use the word “category” in the contexts where the corresponding object is used in a way that is functorial for equivalences of categories and the word “precategory” otherwise.

The main construction of this paper was introduced in [7]. I am grateful to the Centre for Quantum Mathematics and Computation (QMAC) and the Mathematical Institute of the University of Oxford for their hospitality during my work on the previous version of the paper and to the Department of Computer Science and Engineering of the University of Gothenburg and Chalmers University of Technology for its the hospitality during my work on the present version.

2. Construction of $CC(\mathcal{C}, p)$.

2.1. DEFINITION. *Let \mathcal{C} be a category. A universe structure on a morphism $p : \tilde{U} \rightarrow U$ in \mathcal{C} is a mapping that assigns to any morphism $f : X \rightarrow U$ in \mathcal{C} a pull-back square*

$$\begin{array}{ccc} (X; f) & \xrightarrow{Q(f)} & \tilde{U} \\ p_{X,f} \downarrow & & \downarrow p \\ X & \xrightarrow{f} & U \end{array}$$

A universe in \mathcal{C} is a morphism p together with a universe structure on it.

In what follows we will write $(X; f_1, \dots, f_n)$ for $(\dots((X; f_1); f_2) \dots; f_n)$.

2.2. EXAMPLE. Let G be a group. Consider the category BG with one object pt whose monoid of endomorphisms is G . Recall that any commutative square where all four arrows are isomorphisms is a pull-back square. Let $p : pt \rightarrow pt$ be the unit element of G . Then a universe structure on p can be defined by specifying, for every $g : pt \rightarrow pt$, of the horizontal morphism $Q(g)$ in the corresponding canonical square. There are no restrictions on the choice of $Q(g)$ since for any such choice one can take the vertical morphism to be $Q(g)g^{-1}$ obtaining a pull-back square. Therefore, the set of universe structures on p is G^G . The automorphisms of BG are given by $Aut(G)$ (with two automorphisms being isomorphic as functors if they differ by an inner automorphisms of G). Therefore, there are $(G^G)/Aut(G)$ isomorphism classes of categories with universes with the underlying

category BG and the underlying universe morphism being $Id : pt \rightarrow pt$. Note that in this case all auto-equivalences of the category are automorphisms and so simply saying that we will consider universes up to an equivalence of the underlying category does not change the answer. To have, as is suggested by category-theoretic intuition, no more than one universe structure on a morphism one needs to consider categories with universes up to equivalences of categories with universes and then one has the obligation to prove that the constructions that are supposed to produce objects such as C-systems map equivalences of categories with universes to isomorphisms. In the case of the main construction of this paper it is achieved in Lemma 4.8.

Suppose that we are given a morphism $F : X \rightarrow U$. Then for $f : W \rightarrow X$ and $g : W \rightarrow \tilde{U}$ such that $f \circ F = g \circ p$ we will denote by $f * g$ the unique morphism $W \rightarrow (X; F)$ such that

$$\begin{aligned} (f * g) \circ p_{X,F} &= f \\ (f * g) \circ Q(F) &= g \end{aligned}$$

For $X' \xrightarrow{f} X \xrightarrow{F} U$ we let $Q(f, F)$ denote the morphism

$$(p_{X',f \circ F} \circ f) * Q(f \circ F) : (X'; f \circ F) \rightarrow (X; F)$$

such that in particular

$$Q(f, F) \circ Q(F) = Q(f \circ F) \tag{1}$$

2.3. LEMMA. *The square*

$$\begin{array}{ccc} (X'; f \circ F) & \xrightarrow{Q(f,F)} & (X; F) \\ p_{X',f \circ F} \downarrow & & \downarrow p_{X,F} \\ X' & \xrightarrow{f} & X \end{array} \tag{2}$$

is a pull-back square.

PROOF. Consider the diagram

$$\begin{array}{ccccc} (X'; f \circ F) & \xrightarrow{Q(f,F)} & (X; F) & \xrightarrow{Q(F)} & \tilde{U} \\ p_{X',f \circ F} \downarrow & & \downarrow p_{X,F} & & \downarrow p \\ X' & \xrightarrow{f} & X & \xrightarrow{F} & U \end{array}$$

The composition of two squares of this diagram equals the square with the sides $p_{X',f \circ F}$, $f \circ F$, $Q(f \circ F)$ and p , which is a pull-back square. The right hand side square in this diagram is a pull-back square. This implies that the left hand side square is a pull-back square. ■

2.4. LEMMA. *If $f : X' \rightarrow X$ is an isomorphism then $Q(f, F)$ is an isomorphism.*

PROOF. It follows from Lemma 2.3 by general properties of pull-back squares. ■

2.5. LEMMA. *For $f' : X'' \rightarrow X'$, $f : X' \rightarrow X$ and $F : X \rightarrow U$ one has*

$$Q(f', f \circ F) \circ Q(f, F) = Q(f' \circ f, F)$$

PROOF. Both sides of the equality are morphisms to $(X; F)$, therefore it is sufficient to verify that

$$Q(f', f \circ F) \circ Q(f, F) \circ Q(F) = Q(f' \circ f, F) \circ Q(F)$$

and

$$Q(f', f \circ F) \circ Q(f, F) \circ p_{X,F} = Q(f' \circ f, F) \circ p_{X,F}$$

For the first one we have

$$Q(f', f \circ F) \circ Q(f, F) \circ Q(F) = Q(f', f \circ F) \circ Q(f \circ F) = Q(f' \circ f \circ F)$$

and

$$Q(f' \circ f, F) \circ Q(F) = Q(f' \circ f \circ F)$$

and for the second one we have

$$Q(f', f \circ F) \circ Q(f, F) \circ p_{X,F} = Q(f', f \circ F) \circ p_{X',f \circ F} \circ f = p_{X'',f' \circ f \circ F} \circ f' \circ f$$

and

$$Q(f' \circ f, F) \circ p_{X,F} = p_{X'',f' \circ f \circ F} \circ f' \circ f.$$

■

2.6. DEFINITION. *A universe category is a triple (\mathcal{C}, p, pt) where \mathcal{C} is a category, $p : \tilde{U} \rightarrow U$ is a morphism in \mathcal{C} with a universe structure on it and pt is a final object in \mathcal{C} .*

We will often denote a universe category by a pair (\mathcal{C}, p) .

Let (\mathcal{C}, p) be a universe category. Define by induction on n pairs $(Ob_n(\mathcal{C}, p), int_n)$ where $Ob_n = Ob_n(\mathcal{C}, p)$ are sets and $int_n : Ob_n \rightarrow Ob(\mathcal{C})$ are functions, as follows:

1. $Ob_0 = unit$ where $unit$ is the distinguished set with only one point tt and int_0 maps this point to pt .
2. $Ob_{n+1} = \coprod_{A \in Ob_n} Hom_{\mathcal{C}}(int_n(A), U)$ and $int_{n+1}(A, F) = (int_n(A); F)$.

In what follows we will write int instead of int_n since n can usually be inferred.

Define for each n the function $ft_{n+1} : Ob_{n+1} \rightarrow Ob_n$ by the formula $ft_{n+1}(A, F) = A$ and define ft_0 as the identity function of Ob_0 .

For each $B = (ft(B), F) \in Ob_{n+1}$ define $p_B : int(B) \rightarrow int(ft(B))$ as $p_{int(ft(B)), F}$. For $B \in Ob_0$ define p_B as $Id_{int(B)}$.

For each $A \in Ob_m$, $B = (ft(B), F) \in Ob_{n+1}$ and $f : int(A) \rightarrow int(ft(B))$ define $f^*(B) \in Ob_{m+1}$ as

$$f^*(B) = (A, f \circ F) \tag{3}$$

and $q(f, B) : int(f^*(B)) \rightarrow int(B)$ as

$$q(f, B) = Q(f, F) \tag{4}$$

Recall the concept of a C0-system defined in [9, Definition 2.1].

2.7. PROBLEM. For each universe category (\mathcal{C}, p, pt) to define a C0-system $CC0(\mathcal{C}, p)$.

2.8. CONSTRUCTION. We set

$$Ob(CC0(\mathcal{C}, p)) = \coprod_{n \geq 0} Ob_n(\mathcal{C}, p)$$

where $Ob_n = Ob_n(\mathcal{C}, p)$ are the sets introduced above. Let

$$int_{Ob} : Ob(CC0(\mathcal{C}, p)) \rightarrow \mathcal{C}$$

be the sum of the functions int_n . Let

$$Mor(CC0(\mathcal{C}, p)) = \coprod_{\Gamma, \Gamma' \in Ob(CC0(\mathcal{C}, p))} Hom_{\mathcal{C}}(int_{Ob}(\Gamma), int_{Ob}(\Gamma'))$$

Define the function

$$int_{Mor} : Mor(CC0(\mathcal{C}, p)) \rightarrow Mor(\mathcal{C})$$

by the formula

$$int_{Mor}(\Gamma, (\Gamma', a)) = a$$

We will often write simply int for int_{Ob} and int_{Mor} .

The identity morphisms and the composition of morphisms are defined as in \mathcal{C} . The proofs of the axioms of a category are straightforward.

The definition of the length function is obvious.

We define pt as the unique element $(0, tt)$ of $Ob(CC0(\mathcal{C}, p))$ of length zero.

The function $ft : Ob(CC0) \rightarrow Ob(CC0)$ is defined as the sum of functions ft_n defined above.

The p -morphisms $p_{(n,A)}$ are defined such that $int(p_{(n,A)}) = p_A$ where p_A where defined above.

Similarly one defines the morphisms $q(f, (n + 1, B))$ such that $int(q(f, (n + 1, B))) = q(f, B)$.

Lemma 2.5 shows that the structure that we have defined satisfies the axioms of a C0-system given in [9, Definition 2.1].

Let us also note the following formulas. For $\Delta = (n + 1, (B, F))$ and $\Gamma = (n, B)$ one has

$$p_{\Delta} = (\Delta, (\Gamma, p_{int(B,F)})) \tag{5}$$

For $\Gamma' = (m, A)$, $\Gamma = (n, B)$ and $f : \Gamma' \rightarrow \Gamma$ one has

$$f^*(n + 1, (B, F)) = (m + 1, (A, \text{int}(f) \circ F)) \tag{6}$$

$$q(f, (n + 1, (B, F))) = (f^*(\Delta), (\Delta, Q(\text{int}(f), F))) \tag{7}$$

2.9. LEMMA. *The functions int_{Ob} and int_{Mor} defined above form a fully faithful functor from the category underlying the C0-system $CC0(\mathcal{C}, p)$ to \mathcal{C} .*

PROOF. Easy from the construction. ■

2.10. REMARK. The image of int on objects consists of those objects for which the unique morphism to pt can be represented as a composition of morphisms of the form $p_{X,F}$. Note that int need not be an injection on the sets of objects. For example, if \mathcal{C} is the one point category with its unique structure of a universe category then $Ob(CC(\mathcal{C}, p))$ will be isomorphic to the set of natural numbers.

2.11. PROBLEM. *For each universe category (\mathcal{C}, p, pt) to define a C-system $CC(\mathcal{C}, p)$.*

2.12. CONSTRUCTION. We will define $CC(\mathcal{C}, p)$ as an extension of $CC0(\mathcal{C}, p)$ using [9, Proposition 2.4]. In particular $Ob(CC) = Ob(CC0)$, $Mor(CC) = Mor(CC0)$ and similarly for the length function, ft , p -morphisms and q -morphisms.

The canonical squares of $CC0(\mathcal{C}, p)$ are of the form

$$\begin{array}{ccc} f^*(\Gamma) & \xrightarrow{q(f,\Gamma)} & \Gamma \\ p_{f^*(\Gamma)} \downarrow & & \downarrow p_\Gamma \\ \Gamma' & \xrightarrow{f} & ft(\Gamma) \end{array} \tag{8}$$

For $\Gamma = (n + 1, (B, F))$ where $B \in Ob_n(\mathcal{C}, p)$ and $F : \text{int}(A) \rightarrow U$, $\Gamma' = (m, A)$ where $A \in Ob_m(\mathcal{C}, p)$, and $f = (\Gamma', (\Gamma, a))$ the image of this square under the functor int is of the form

$$\begin{array}{ccc} (\text{int}(A); a \circ F) & \xrightarrow{Q(a,F)} & (\text{int}(B); F) \\ p_{\text{int}(A), a \circ F} \downarrow & & \downarrow p_{\text{int}(B), F} \\ \text{int}(A) & \xrightarrow{F} & \text{int}(B) \end{array}$$

This is one of the squares of the form (2) and therefore by Lemma 2.3 it is a pull-back square. Since int is fully faithful by Lemma 2.9, the squares (8) are pull-back squares in the codomain of a fully faithful functor and therefore they are also pull-back squares in the domain of this functor, i.e., in $CC0(\mathcal{C}, p)$. In view of [9, Proposition 2.4] this implies that the C0-system $CC0(\mathcal{C}, p)$ has a unique structure of a C-system and we denote this C-system by $CC(\mathcal{C}, p)$.

2.13. REMARK. Recall that in [9] we suggested the notation $Ob_n(CC)$ for the set of objects of length n of a C-system CC . We will avoid using this notation here because the sets $Ob_n(\mathcal{C}, p)$ are not equal to the subsets of elements of length n in $CC(\mathcal{C}, p)$. Indeed, the elements of $\{\Gamma \in Ob(CC(\mathcal{C}, p)) \mid l(\Gamma) = n\}$ are not the elements of $Ob_n(\mathcal{C}, p)$ but pairs of the form (n, A) where $A \in Ob_n(\mathcal{C}, p)$.

2.14. EXAMPLE. An important example of a C-system of the form $CC(\mathcal{C}, p)$ is “the” C-system Fam of families of sets considered in [3] and [4]. The definition of Fam in [4, p.238] as well as the preceding it discussion in [4, p.232] is somewhat incomplete in that the notion of “a set” and moreover the notion of “a family of sets” are taken as being uniquely determined by some previous agreement that is never explicitly referred to.

To define Fam as a C-system of the form $CC(\mathcal{C}, p)$ let us choose two Grothendieck universes U and U_1 in our set theory such that U_1 is an element of U . One then defines the category $Sets(U)$ of sets as the category whose set of objects is U and such that for $X, Y \in U$ the set of morphisms from X to Y in $Sets(U)$ is the set of functions from X to Y in the ambient set theory (which automatically is an element of U). This category will contain U_1 as an object and also, because of the closure conditions that U satisfies, it will contain as an object the set \tilde{U}_1 of pairs (X, x) where $X \in U_1$ and $x \in X$. Since morphisms in $Sets$ are the same as functions in the ambient set theory we also get $p_{U_1} : \tilde{U}_1 \rightarrow U_1$ that takes (X, x) to X . Using the standard construction of pull-backs in sets we obtain a universe structure on p . Now we can define:

$$Fam(U, U_1) := CC(Sets(U), p_{U_1})$$

The explicit definition given in [4] avoids the use of the second universe (universe U in our notations) by constructing the same C-system “by hand”. In our approach we have to use U but the resulting C-system does not depend on U . Indeed, if our set theory assumes two Grothendieck universes U and U' such that both contain U_1 as an element then one can show that

$$CC(Sets(U), p_{U_1}) = CC(Sets(U'), p_{U_1}) \tag{9}$$

where the equality means in particular that the sets of objects of these two C-systems are equal as sets. Because of this one can denote this C-system as $Fam(U_1)$.

3. On homomorphisms of C-systems

We will need below the concept of a homomorphism of C-systems. Homomorphisms of C-systems were defined in [9, Remark 2.8]. Let us recall it here in a more detailed form.

3.1. DEFINITION. *Let CC_1, CC_2 be C-systems. A homomorphism F from CC_1 to CC_2 is a pair of functions $F_{Ob} : Ob(CC_1) \rightarrow Ob(CC_2)$, $F_{Mor} : Mor(CC_1) \rightarrow Mor(CC_2)$ such that:*

1. F commutes with the length functions, i.e., for all $X \in Ob(CC_1)$ one has

$$l(F_{Ob}(X)) = l(X)$$

2. F commutes with the ft function, i.e., for all $X \in Ob(CC_1)$ one has

$$ft(F_{Ob}(X)) = F_{Ob}(ft(X))$$

3. F is a functor, i.e., one has:

- (a) F_{Mor} and F_{Ob} commute with the domain and codomain functions,
- (b) for all $X \in Ob(CC_1)$ one has

$$F_{Mor}(Id_X) = Id_{F_{Mor}(X)}$$

(c) for all $f, g \in Mor(CC_1)$ of the form $f : X \rightarrow Y$, $g : Y \rightarrow Z$ one has

$$F_{Mor}(f \circ g) = F_{Mor}(f) \circ F_{Mor}(g)$$

4. F takes canonical projections to canonical projections, i.e., for all $X \in Ob(CC_1)$ one has

$$p_{F_{Ob}(X)} = F_{Mor}(p_X)$$

5. F takes q -morphisms to q -morphisms, i.e., for all $X, Y \in Ob(CC_1)$ such that $l(Y) > 0$ and all $f : X \rightarrow ft(Y)$ one has

$$F_{Mor}(q(f, Y)) = q(F_{Mor}(f), F_{Ob}(Y))$$

6. F takes s -morphisms to s -morphisms, i.e., for all $X, Y \in Ob(CC_1)$ such that $l(Y) > 0$ and $f : X \rightarrow Y$ one has

$$s_{F_{Mor}(f)} = F_{Mor}(s_f)$$

In what follows we will write F for both F_{Ob} and F_{Mor} since the choice of which one is meant is determined by the type of the argument. Note that the condition that F commutes with the domain function together with the q -morphism condition implies that for all $X, Y \in Ob(CC_1)$ such that $l(Y) > 0$ and all $f : X \rightarrow ft(Y)$ one has

$$F(f^*(Y)) = F(f)^*(F(Y)) \tag{10}$$

3.2. LEMMA. *Let $F : CC_1 \rightarrow CC_2$ and $G : CC_2 \rightarrow CC_3$ be homomorphisms of C -systems. Then the compositions of functions $F_{Ob} \circ G_{Ob}$ and $F_{Mor} \circ G_{Mor}$ is a homomorphism of C -systems.*

PROOF. The proof is relatively straightforward but long and we leave it for the formal version(s) of the paper. ■

3.3. REMARK. Since homomorphisms of C-systems are pairs of functions between sets satisfying certain conditions and the composition is given by composition of these functions, the associativity and unitality of this composition follows easily from the associativity and unitality of the composition of functions between sets. Therefore, if we restrict our attention to the C-systems whose sets Ob and Mor are elements of a chosen set (“universe”) U that contains natural numbers and is closed under the power-set operation, then such C-systems, their homomorphisms, compositions of these homomorphisms and the identity homomorphisms form a category of C-systems in U .

3.4. LEMMA. *Let CC_1, CC_2, F_{ob} and F_{Mor} be as above. Assume further that these data satisfy all of the conditions of the definition except, possibly, the s -morphisms condition. Then (F_{Ob}, F_{Mor}) satisfies the s -morphisms condition and forms a homomorphism of C-systems.*

PROOF. Let $f : X \rightarrow Y$ be as in the s -morphism condition. We need to show that $F(s_f) = s_{F(f)}$. Observe first that the right hand side is well defined since $l(F(Y)) = l(Y) > 0$. We have $F(s_f) : F(X) \rightarrow F((f \circ p_Y)^*(Y))$ and $s_{F(f)} : F(X) \rightarrow (F(f) \circ p_{F(Y)})^*(F(Y))$. One proves that codomains of both morphisms are equal using that F is a functor, the p -morphisms condition and (10).

Since the canonical squares of CC_2 are pull-back squares the object

$$(F(f) \circ p_{F(Y)})^*(F(Y))$$

is a fiber product with the projections $q(F(f) \circ p_{F(Y)}, F(Y))$ and $p_{F(f)^*(F(Y))}$. Therefore it is sufficient to check that one has

$$F(s_f) \circ q(F(f) \circ p_{F(Y)}, F(Y)) = s_{F(f)} \circ q(F(f) \circ p_{F(Y)}, F(Y)) \quad (11)$$

and

$$F(s_f) \circ p_{F(f)^*(F(Y))} = s_{F(f)} \circ p_{F(f)^*(F(Y))} \quad (12)$$

We have

$$F(s_f) \circ q(F(f) \circ p_{F(Y)}, F(Y)) = F(s_f) \circ q(F(f) \circ F(p_Y), F(Y)) =$$

$$F(s_f) \circ q(F(f \circ p_Y), F(Y)) = F(s_f) \circ F(q(f \circ p_Y, Y)) = F(s_f \circ q(f \circ p_Y, Y)) = F(f)$$

where the first equality holds by condition (4) of Definition 3.1, the second and the fourth equalities by condition (3), the third equality by condition (5) and the fifth equality by axiom [9, Definition 2.3(3)] of the operation s for CC_1 .

On the other hand

$$s_{F(f)} \circ q(F(f) \circ p_{F(Y)}, F(Y)) = F(f)$$

directly by the axiom [9, Definition 2.3(3)] of the operation s for CC_2 . This proves (11).

For the equation (12) we have

$$F(s_f) \circ p_{F(f)^*(F(Y))} = F(s_f) \circ p_{F(f^*(Y))} = F(s_f) \circ F(p_{f^*(Y)}) = F(s_f \circ p_{f^*(Y)}) =$$

$$F(Id_X) = Id_{F(X)}$$

where the first equation holds by (10), the second one by condition (4), the third one by condition (3), the fourth one by the axiom [9, Definition 2.3(2)] of the operation s for CC_1 and the fifth one by condition (3).

On the other hand

$$s_{F(f)} \circ p_{F(f)^*(F(Y))} = Id_{F(X)}$$

directly by the axiom [9, Definition 2.3(2)] of the operation s for CC_2 . This completes the proof of Lemma 3.4. ■

3.5. REMARK. As defined in [9], a C-system without operation s is called a C0-system. The pairs F_{Ob}, F_{Mor} that satisfy all of the conditions of Definition 3.1 other than, possibly, the s -morphism condition are homomorphisms of C0-systems. Therefore, if one defines a categories of C-systems and C0-systems based on a particular universe of sets as outlined in Remark 3.3 then Lemma 3.4 implies that the forgetting functor from the category of C-systems in U to C0-systems in U is a full embedding.

4. Functoriality of $CC(\mathcal{C}, p)$

4.1. DEFINITION. Let (\mathcal{C}, p, pt) and (\mathcal{C}', p', pt') be universe categories. A functor of universe categories from (\mathcal{C}, p, pt) to (\mathcal{C}', p', pt') is a triple $(\Phi, \phi, \tilde{\phi})$ where $\Phi : \mathcal{C} \rightarrow \mathcal{C}'$ is a functor and $\phi : \Phi(U) \rightarrow U', \tilde{\phi} : \Phi(\tilde{U}) \rightarrow \tilde{U}'$ are morphisms such that:

1. Φ takes the canonical pull-back squares based on p to pull-back squares,
2. Φ takes pt to a final object of \mathcal{C}' ,
3. the square

$$\begin{array}{ccc} \Phi(\tilde{U}) & \xrightarrow{\tilde{\phi}} & \tilde{U}' \\ \Phi(p) \downarrow & & \downarrow p' \\ \Phi(U) & \xrightarrow{\phi} & U' \end{array}$$

is a pull-back square.

Let

$$(\Phi, \phi, \tilde{\phi}) : (\mathcal{C}, p, pt) \rightarrow (\mathcal{C}', p', pt')$$

be a functor of universes categories. Let $Ob_n = Ob_n(\mathcal{C}, p)$ and $Ob'_n = Ob_n(\mathcal{C}', p')$. Let int and int' be the corresponding functions to \mathcal{C} and \mathcal{C}' .

Denote by ψ the isomorphism $\psi : pt' \rightarrow \Phi(pt)$. Define, by induction on n , pairs (H_n, ψ_n) where $H_n : Ob_n \rightarrow Ob'_n$ and ψ_n is a family of isomorphisms of the form

$$\psi_n(A) : int'(H_n(A)) \rightarrow \Phi(int(A))$$

given for all $A \in Ob_n$. We set:

1. for $n = 0$, H_0 is the unique map from a one point set to a one point set and $\psi_0(A) = \psi$,
2. for the successor of n we set

$$H_{n+1}(A, F) = (H_n(A), \psi_n(A) \circ \Phi(F) \circ \phi) \tag{13}$$

and define

$$\psi_{n+1}(A, F) : (int(H_n(A)); \psi_n(A) \circ \Phi(F) \circ \phi) \rightarrow \Phi(int(A, F))$$

as the unique morphism such that the left hand side square of the diagram

$$\begin{array}{ccccccc}
 int'(H_{n+1}(A, F)) & \xrightarrow{\psi_{n+1}(A, F)} & \Phi(int(A, F)) & \xrightarrow{\Phi(Q(F))} & \Phi(\tilde{U}) & \xrightarrow{\tilde{\phi}} & \tilde{U}' \\
 \downarrow p_{H_{n+1}(A, F)} & & \downarrow \Phi(p_{(A, F)}) & & \downarrow \Phi(p) & & \downarrow p' \\
 int'(H_n(A)) & \xrightarrow{\psi_n(A)} & \Phi(int(A)) & \xrightarrow{\Phi(F)} & \Phi(U) & \xrightarrow{\phi} & U'
 \end{array} \tag{14}$$

commutes, i.e.,

$$\psi(A, F) \circ \Phi(p_{(A, F)}) = p_{H(A, F)} \circ \psi(A) \tag{15}$$

and

$$\psi_{n+1}(A, F) \circ \Phi(Q(F)) \circ \tilde{\phi} = Q(\psi_n(A) \circ \Phi(F) \circ \phi) \tag{16}$$

Note that the existence and uniqueness of $\psi_{n+1}(A, F)$ follows from the fact that the right hand side squares of (14) are pull-back squares as a corollary of the definition of a universe category functor and the fact that the canonical square for the morphism $\psi_n(A) \circ \Phi(F) \circ \phi$ commutes.

Moreover since the outer square of (14) is a pull-back square, the left-most square commutes and the two right hand side squares are pull-back squares we conclude that the left hand side square is a pull-back square. In combination with the inductive assumption that $\psi_n(A)$ is an isomorphism this implies that $\psi_{n+1}(A, F)$ is an isomorphism.

In what follows we will write $\psi(A)$ instead of $\psi_n(A)$ since n can often be inferred.

4.2. LEMMA. *The functions H commute with the functions ft , i.e., for $A \in Ob_n$ one has*

$$ft(H(A)) = H(ft(A))$$

PROOF. Immediate from the construction. ■

Let $A \in Ob_m$, $A' \in Ob_{m'}$ and $a : int(A) \rightarrow int(A')$. Define a morphism

$$H(a) : int'(H(A)) \rightarrow int'(H(A'))$$

as

$$H(a) = \psi(A) \circ \Phi(a) \circ \psi(A')^{-1} \tag{17}$$

4.3. LEMMA. For Φ as above one has:

1. for $A \in Ob_n$ one has $H(Id_{int(A)}) = Id_{int(H(A))}$,
2. for $a' : int(A'') \rightarrow int(A')$ and $a : int(A') \rightarrow int(A)$ one has $H(a' \circ a) = H(a') \circ H(a)$.

PROOF. Immediate from the construction. ■

4.4. LEMMA. For $A \in Ob_n$ one has $H(p_A) = p_{H(A)}$.

PROOF. If $n = 0$ the statement is obvious. For $(A, F) \in Ob_{n+1}$ we have

$$H(p_{(A,F)}) = \psi(A, F) \circ \Phi(p_{(A,F)}) \circ \psi(A)^{-1}$$

Therefore we need to show that

$$\psi(A, F) \circ \Phi(p_{(A,F)}) = p_{H(A,F)} \circ \psi(A)$$

which is (15). ■

4.5. LEMMA. Let $A \in Ob_m$, $B = (ft(B), F) \in Ob_{n+1}$ and $a : int(A) \rightarrow int(ft(B))$ is a morphism. Then one has

$$H(a^*(B)) = H(a)^*(H(B)) \tag{18}$$

and

$$H(q(a, B)) = q(H(a), H(B)) \tag{19}$$

PROOF. We have

$$H(a^*(ft(B), F)) = H(A, a \circ F) = (H(A), \psi(A) \circ \Phi(a \circ F) \circ \phi)$$

and

$$\begin{aligned} H(a)^*(H(ft(B), F)) &= H(a)^*(H(ft(B)), \psi(ft(B)) \circ \Phi(F) \circ \phi) = \\ &= (H(A), H(a) \circ \psi(ft(B)) \circ \Phi(F) \circ \phi) \end{aligned}$$

Therefore we need to check that

$$\psi(A) \circ \Phi(a) = H(a) \circ \psi(ft(B))$$

which follows from the definition of $H(a)$.

To prove (19) it is sufficient, since $\psi(B)$ is an isomorphism, to show that

$$H(q(a, B)) \circ \psi(B) = q(H(a), H(B)) \circ \psi(B)$$

In view of (18) both sides are morphisms from $\text{int}(H(a^*(B)))$ to

$$\Phi(\text{int}(B)) = \Phi(\text{int}(ft(B)); F)$$

Since the two right squares of (14) for $(ft(B), F)$ are pull-back squares, $\Phi(\text{int}(ft(B)); F)$ is a fiber product with projections $\Phi(Q(F)) \circ \tilde{\phi}$ and $\Phi(p_B)$. Therefore it is sufficient to check two equalities

$$H(q(a, B)) \circ \psi(B) \circ \Phi(Q(F)) \circ \tilde{\phi} = q(H(a), H(B)) \circ \psi(B) \circ \Phi(Q(F)) \circ \tilde{\phi} \quad (20)$$

and

$$H(q(a, B)) \circ \psi(B) \circ \Phi(p_B) = q(H(a), H(B)) \circ \psi(B) \circ \Phi(p_B) \quad (21)$$

Note first that

$$H(q(a, B)) \circ \psi(B) = \psi(a^*(B)) \circ \Phi(q(a, (ft(B), F))) = \psi(A, a \circ F) \circ \Phi(Q(a, F)) \quad (22)$$

where the first equality is by (17) and the second by (4), and

$$\begin{aligned} q(H(a), H(B)) \circ \psi(B) &= q(H(a), (H(ft(B)), \psi(ft(B)) \circ \Phi(F) \circ \phi)) \circ \psi(B) = \\ &= Q(H(a), \psi(ft(B)) \circ \Phi(F) \circ \phi) \circ \psi(ft(B), F) \end{aligned} \quad (23)$$

where the first equality is by (13) and the second equality is by (4).

For (20) we have

$$\begin{aligned} H(q(a, B)) \circ \psi(B) \circ \Phi(Q(F)) \circ \tilde{\phi} &= \psi(A, a \circ F) \circ \Phi(Q(a, F)) \circ \Phi(Q(F)) \circ \tilde{\phi} = \\ &= \psi(A, a \circ F) \circ \Phi(Q(a \circ F)) \circ \tilde{\phi} = Q(\psi(A) \circ \Phi(a \circ F) \circ \phi) \end{aligned}$$

where the first equality is by (22), the second equality is by (1) and the third one by (16), and

$$\begin{aligned} q(H(a), H(B)) \circ \psi(B) \circ \Phi(Q(F)) \circ \tilde{\phi} &= \\ Q(H(a), \psi(ft(B)) \circ \Phi(F) \circ \phi) \circ \psi(ft(B), F) \circ \Phi(Q(F)) \circ \tilde{\phi} &= \\ Q(H(a), \psi(ft(B)) \circ \Phi(F) \circ \phi) \circ Q(\psi(ft(B)) \circ \Phi(F) \circ \phi) &= \\ Q(H(a) \circ \psi(ft(B)) \circ \Phi(F) \circ \phi) &= \\ Q(\psi(A) \circ \Phi(a) \circ \psi(ft(B))^{-1} \circ \psi(ft(B)) \circ \Phi(F) \circ \phi) &= Q(\psi(A) \circ \Phi(a \circ F) \circ \phi) \end{aligned}$$

where the first equality is by (23), the second equality is by (16), the third one by (1), and the fourth one by (17). For (20) we have:

$$H(q(a, B)) \circ \psi(B) \circ \Phi(p_B) = \psi(A, a \circ F) \circ \Phi(Q(a, F)) \circ \Phi(p_B) = \psi(a^*(B)) \circ \Phi(Q(a, F)) \circ p_B =$$

$$\psi(a^*(B)) \circ \Phi(q(a, B) \circ p_B) = \psi(a^*(B)) \circ \Phi(p_{a^*(B)}) \circ \Phi(a)$$

where the first equality is by (22), the second by (3) and the assumption that Φ is a functor, the third one by (4) and the fourth one by the commutativity of the canonical squares and the assumption that Φ is a functor.

For the other side we have:

$$q(H(a), H(B)) \circ \psi(B) \circ \Phi(p_B) = q(H(a), H(B)) \circ p_{H(B)} \circ \psi(ft(B)) =$$

$$p_{H(a^*(B))} \circ H(a) \circ \psi(ft(B)) = p_{H(a^*(B))} \circ \psi(A) \circ \Phi(a) = \psi(a^*(B)) \circ \Phi(p_{a^*(B)}) \circ \Phi(a)$$

Where the first equality is by (15), the second by the commutativity of the canonical squares, the third by (17) and the fourth again by (15). This completes the proof of Lemma 4.5. ■

4.6. PROBLEM. *Let*

$$(\Phi, \phi, \tilde{\phi}) : (\mathcal{C}, p, pt) \rightarrow (\mathcal{C}', p', pt')$$

be a functor of universes categories. To define a homomorphism $H = H(\Phi, \phi, \tilde{\phi})$ from $CC(\mathcal{C}, p)$ to $CC(\mathcal{C}', p')$.

4.7. CONSTRUCTION. We define H_{Ob} as the sum of functions H_n constructed above and for

$$(\Gamma, (\Gamma', a)) \in Mor(CC(\mathcal{C}, p))$$

we set

$$H_{Mor}(\Gamma, (\Gamma', a)) = (H_{Ob}(\Gamma), (H_{Ob}(\Gamma'), H(a)))$$

where $H(a)$ was constructed above.

The fact that H_{Ob} commutes with the length functions is immediate from the construction. The fact that it commutes with the ft functions follows from Lemma 4.2, the fact that H_{Ob} and H_{Mor} form a functor follows from Lemma 4.3. The fact that H_{Mor} satisfies the p -condition follows from Lemma 4.4. The fact that H_{Mor} satisfies the q -condition follows from Lemma 4.5.

Applying Lemma 3.4 we conclude that $H = (H_{Ob}, H_{Mor})$ is a homomorphism of C-systems.

4.8. LEMMA. *Let $(\Phi, \phi, \tilde{\phi})$ be as in Problem 4.6 and let H be the solution given by Construction 4.7. Then one has:*

1. *If Φ is a faithful functor and ϕ is a monomorphism then H is an injection of C-systems.*
2. *If Φ is a fully faithful functor and ϕ is an isomorphism then H is an isomorphism.*

PROOF. Both statements in relation to objects have straightforward proofs by induction on the length. In relation to morphisms the statements follow from the ones about the objects and the fact that int is fully faithful. ■

Lemma 4.8 can be further specialized into the following example.

4.9. EXAMPLE. Let \mathcal{C} be a category and $p : \tilde{U} \rightarrow U$ a morphism in \mathcal{C} . Let now $(p_{X,F}, Q(F))$ and $(p'_{X,F}, Q'(F))$ be two universe structures on p and pt and pt' be two final objects in \mathcal{C} . These data give us two universe categories. Let us denote them by \mathcal{UC} and \mathcal{UC}' . The identity functor $\Phi = Id_{\mathcal{C}}$ on \mathcal{C} together with the identity morphisms $\phi = Id_U$ and $\tilde{\phi} = Id_{\tilde{U}}$ define a universe category functor $\Phi : \mathcal{UC} \rightarrow \mathcal{UC}'$. The corresponding homomorphism of C-systems $H_{\Phi} : CC(\mathcal{UC}) \rightarrow CC(\mathcal{UC}')$ is an isomorphism with the inverse isomorphism given by the same triple considered as a universe functor from \mathcal{UC} to \mathcal{UC}' . This example shows that, up to a “canonical” isomorphism, the C-system defined by a universe category depends only on the category \mathcal{C} and the morphism p .

4.10. PROBLEM. Let (\mathcal{C}, p) be a universe category. Let CC be a C-system. Given the following collection of data:

1. A functor $I : CC \rightarrow \mathcal{C}$ from the underlying category of CC to \mathcal{C} ,
2. For each $\Gamma \in CC$ a function

$$u_{\Gamma} : Ob_1(\Gamma) \rightarrow Hom_{\mathcal{C}}(I(\Gamma), U)$$

3. For each $\Gamma \in CC$, $\Delta \in Ob_1(\Gamma)$ an isomorphism

$$\gamma_{\Delta} : (I(\Gamma); u(\Delta)) \rightarrow I(\Delta)$$

such that

1. the morphism $\pi_{I(pt)} : I(pt) \rightarrow pt$ is an isomorphism
2. for each $f : \Gamma' \rightarrow \Gamma$ and $\Delta \in Ob_1(\Gamma)$ one has $u_{\Gamma'}(f^*(\Delta)) = I(f) \circ u_{\Gamma}(\Delta)$,
3. for each $f : \Gamma' \rightarrow \Gamma$ and $\Delta \in Ob_1(\Gamma)$ one has $p_{I(\Gamma), u(\Delta)} = \gamma_{\Delta} \circ I(p_{\Delta})$,
4. for each $f : \Gamma' \rightarrow \Gamma$ and $\Delta \in Ob_1(\Gamma)$ one has $\gamma_{f^*(\Delta)} \circ I(q(f, \Delta)) = Q(I(f), u(\Delta)) \circ \gamma_{\Delta}$

to construct a C-system homomorphism

$$H(I, u, \gamma) : CC \rightarrow CC(\mathcal{C}, p)$$

In what follows we will often write u instead of u_{Γ} .

4.11. CONSTRUCTION. First we construct by induction on n pairs (H_n, ψ_n) where

$$H_n : Ob_n(CC) \rightarrow Ob_n(\mathcal{C}, p)$$

is a function and ψ_n is a family of isomorphisms of the form

$$\psi_n(\Gamma) : int(H(\Gamma)) \rightarrow I(\Gamma)$$

given for all $\Gamma \in Ob_n(CC)$ as follows (we will sometimes write ψ instead of ψ_n and H instead of H_n):

1. For $n = 0$ we set

$$\begin{aligned} H(pt) &= pt \\ \psi(pt) &= (\pi_{I(pt)})^{-1} : pt \rightarrow I(pt) \end{aligned}$$

2. For the successor of n , Γ such that $H_n(\Gamma) = B$ and $\Delta \in Ob_1(\Gamma)$ we set

$$H_{n+1}(\Delta) = (B, \psi(\Gamma) \circ u(\Delta)) \tag{24}$$

and

$$\psi(\Delta) = Q(\psi(\Gamma), u(\Delta)) \circ \gamma_\Delta \tag{25}$$

The fact that $\psi(\Delta)$ is an isomorphism follows from the inductive assumption, the assumption that γ_Δ is an isomorphism and Lemma 2.4.

The functions H_n define a function

$$H_{Ob} : Ob(CC) \rightarrow Ob(CC(\mathcal{C}, p))$$

where $H_{Ob}(\Gamma) = (l(\Gamma), H_l(\Gamma))$ that commutes with the length functions and functions ft .
For $f : \Gamma' \rightarrow \Gamma$ define

$$H_{Mor}(f) = (H_{Ob}(\Gamma'), (H_{Ob}(\Gamma), \psi(\Gamma') \circ I(f) \circ \psi(\Gamma)^{-1}))$$

This gives us a function

$$H_{Mor} : Mor(CC) \rightarrow Mor(CC(\mathcal{C}, p))$$

Note that we can also define $H_{Mor}(f)$ as the unique morphism such that

$$int(H_{Mor}(f)) = \psi(\Gamma') \circ I(f) \circ \psi(\Gamma)^{-1} \tag{26}$$

Without using any more assumptions on I , γ and u one verifies easily that the pair $H = (H_{Ob}, H_{Mor})$ is a functor from the underlying category of CC to the underlying category of $CC(\mathcal{C}, p)$.

In view of Lemma 3.4 it remains to verify that H satisfies the p -morphism and the q -morphism conditions of Definition 3.1.

For the p -condition we need to verify that $H(p_\Gamma) = p_{H(\Gamma)}$ for all Γ . Since both sides have the same domain and codomain and int is bijective on morphisms with the a given domain and codomain it is sufficient to verify that

$$\text{int}(H(p_\Gamma)) = \text{int}(p_{H(\Gamma)})$$

We proceed by induction on $n = l(\Gamma)$:

1. for $n = 0$

$$\text{int}(H(\text{Id}_{pt})) = \text{int}(\text{Id}_{pt}) = \text{int}(p_{pt}) = \text{int}(p_{H(pt)})$$

2. for the successor of n let $\Delta \in \text{Ob}_{n+1}(CC)$ and $\Gamma = \text{ft}(\Delta)$. Then $\Delta \in \text{Ob}_1(\Gamma)$ and

$$\begin{aligned} \text{int}(H(p_\Delta)) &= \psi(\Delta) \circ I(p_\Delta) \circ \psi(\Gamma)^{-1} = Q(\psi(\Gamma), u(\Delta)) \circ \gamma_\Delta \circ I(p_\Delta) \circ \psi(\Gamma)^{-1} = \\ &= Q(\psi(\Gamma), u(\Delta)) \circ p_{I(\Gamma), u(\Delta)} \circ \psi(\Gamma)^{-1} \end{aligned}$$

where the first equality is by (26), the second one by (25) and the third one by condition (3) of the problem. On the other hand we have

$$\text{int}(p_{H(\Delta)}) = \text{int}(p_{(n+1, (H_n(\Gamma), \psi(\Gamma) \circ u(\Delta)))}) = p_{\text{int}(H(\Gamma), \psi(\Gamma) \circ u(\Delta))}$$

where the first equality is by (24) and the second by (5). Composing with $\psi(\Gamma)$ we get

$$\begin{aligned} \text{int}(H(p_\Delta)) \circ \psi(\Gamma) &= Q(\psi(\Gamma), u(\Delta)) \circ p_{I(\Gamma), u(\Delta)} \\ \text{int}(p_{H(\Delta)}) \circ \psi(\Gamma) &= p_{\text{int}(H(\Gamma), \psi(\Gamma) \circ u(\Delta))} \circ \psi(\Gamma) \end{aligned}$$

and these expressions are equal by commutativity of the squares (2).

To prove the q -condition let us verify first that for $f : \Gamma' \rightarrow \Gamma$ and $\Delta \in \text{Ob}_1(\Gamma)$ one has

$$H(f^*(\Delta)) = H(f)^*(H(\Delta)) \tag{27}$$

Let $H(\Gamma') = (m, A)$ and $H(\Gamma) = (n, B)$. Then

$$H(f^*(\Delta)) = (m + 1, (A, \psi(\Gamma') \circ u(f^*(\Delta))))$$

by (24) and

$$H(f)^*(H(\Delta)) = H(f)^*(n + 1, (B, \psi(\Gamma) \circ u(\Delta))) = (m + 1, (A, \text{int}(H(f)) \circ \psi(\Gamma) \circ u(\Delta)))$$

where the first equality holds by (24) and the second by (6). Next one has

$$\psi(\Gamma') \circ u(f^*(\Delta)) = \psi(\Gamma') \circ I(f) \circ u(\Delta)$$

by condition (2) of the problem and

$$\text{int}(H(f)) \circ \psi(\Gamma) \circ u(\Delta) = \psi(\Gamma') \circ I(f) \circ \psi(\Gamma)^{-1} \circ \psi(\Gamma) \circ u(\Delta) = \psi(\Gamma') \circ I(f) \circ u(\Delta)$$

by (26).

The equality (27) implies that the morphisms $H(q(f, \Delta))$ and $q(H(f), H(\Delta))$ have the same domain and codomain. Therefore to prove that they are equal it is sufficient to prove that they become equal after application of int . We further compose both sides with $\psi(\Delta)$. Then we have

$$\begin{aligned} int(H(q(f, \Delta))) \circ \psi(\Delta) &= \psi(f^*(\Delta)) \circ I(q(f, \Delta)) = \\ &Q(\psi(\Gamma'), I(f) \circ u(\Delta)) \circ \gamma_{f^*(\Delta)} \circ I(q(f, \Delta)) = \\ &Q(\psi(\Gamma'), I(f) \circ u(\Delta)) \circ Q(I(f), u(\Delta)) \circ \gamma_\Delta = Q(\psi(\Gamma') \circ I(f), u(\Delta)) \circ \gamma_\Delta \end{aligned}$$

where the first equality is by (26), the second by (25), the third by condition (4) of the problem and the fourth by Lemma 2.5. On the other hand

$$\begin{aligned} int(q(H(f), H(\Delta))) \circ \psi(\Delta) &= int(q(H(f), (n + 1, (B, \psi(\Gamma) \circ u(\Delta)))) \circ \psi(\Delta) = \\ &Q(int(H(f)), \psi(\Gamma) \circ u(\Delta)) \circ \psi(\Delta) = Q(int(H(f)), \psi(\Gamma) \circ u(\Delta)) \circ Q(\psi(\Gamma'), u(\Delta)) \circ \gamma_\Delta = \\ &Q(int(H(f)) \circ \psi(\Gamma), u(\Delta)) \circ \gamma_\Delta = Q(\psi(\Gamma') \circ I(f), u(\Delta)) \circ \gamma_\Delta \end{aligned}$$

where the first equality is by (24), the second by (7), the third by (25), the fourth by Lemma 2.5 and the fifth by (26). This completes Construction 4.11.

4.12. REMARK. The homomorphisms $H(\Phi, \phi, \tilde{\phi})$ can be obtained as particular cases of the homomorphisms $H(I, u, \gamma)$. More precisely, we can state without a proof that

$$H(\Phi, \phi, \tilde{\phi}) = H(I, u, \gamma)$$

where:

1. $I(\Gamma) = \Phi(int(\Gamma))$ and $I(f) = \Phi(int(f))$,
2. for $\Gamma = (n, B)$ and $\Delta = (n + 1, (B, F))$,

$$u_\Gamma(\Delta) = \Phi(F) \circ \phi$$

3. for $\Gamma = (n, B)$ and $\Delta = (n + 1, (B, F))$, γ_Δ is the “natural” isomorphism from $(\Phi(int(B)); \Phi(F) \circ \phi)$ to $\Phi(int(B); F)$. More precisely

$$\gamma_\Delta = ((p_{\Phi(int(B)), \Phi(F) \circ \phi}) * (Q(\Phi(F) \circ \tilde{\phi})))^{-1}$$

4.13. LEMMA. Let I, u and γ be as in Problem 4.10 and let H be the corresponding homomorphism of Construction 4.11. Then one has:

1. If I is a faithful functor and u_Γ are injective then H is an injection of C -systems.
2. If I is a fully faithful functor and u_Γ are bijective then H is an isomorphism of C -systems.

PROOF. Both statements in relation to objects have straightforward proofs by induction on the length. In relation to morphisms the statements follow from the ones about the objects and the fact that int is fully faithful and formula (26). ■

5. Every C-system is isomorphic to a C-system of the form $CC(\mathcal{C}, p)$

5.1. PROBLEM. Let CC be a C-system. Construct a universe category (\mathcal{C}, p) and an isomorphism $CC \cong CC(\mathcal{C}, p)$.

We will provide three different constructions for this problem - Constructions 5.2, 5.6 and 5.7 with the two latter constructions using the first one.

It is customary in the modern mathematics to use “the” category of sets *Sets*. In fact, every set X in the Zermelo-Fraenkel theory defines a category $S(X)$ where:

$$Ob(S(X)) = X$$

$$Mor(S(X)) = \coprod_{x_1, x_2 \in X} Fun(x_1, x_2)$$

where $Fun(x_1, x_2)$ is the set of functions from x_1 to x_2 . This definition makes sense since elements of Zermelo-Fraenkel sets are themselves Zermelo-Fraenkel sets.

Taking X to be sets satisfying particular conditions, e.g. Grothendieck universes, one obtains categories that can be equipped with various familiar structures such as fiber products, internal *Hom*-objects etc. When one says consider “the” category of sets one presumably means the category $S(GU)$ for a chosen Grothendieck universe GU .

The first construction that we provide assumes that we are working in set theory with a chosen Grothendieck universe (or in type theory with a chosen type theoretic universe) that contains the sets of objects and morphisms of our C-system. In the case of a type theory we will actually need two universes in order to have a type of which the first universe is an object.

In what follows we use the notations

$$\widetilde{Ob}(CC) = \{s \in Mor(CC) \mid s : ft(X) \rightarrow X, l(X) > 0, s \circ p_X = Id_X\}$$

and $\partial : \widetilde{Ob}(CC) \rightarrow Ob(CC)$, $\partial(s) = codom(s)$ that were introduced in [9]. We may sometimes abbreviate $Ob(CC)$ to CC .

We will write $\widetilde{Ob}_1(\Gamma)$ for the subset of \widetilde{Ob} that consists of s such that $ft(\partial(s)) = \Gamma$. For $f : \Gamma' \rightarrow \Gamma$ we have the function $s \mapsto f^*(s, 1)$ (see [9]) that maps s to the element of $\widetilde{Ob}_1(\Gamma)$ that is the pull-back of the section s relative to f . We will denote this function by f^* . It is easy to verify from the definitions that

$$(Id_\Gamma)^*(s) = s \tag{28}$$

and for $g : \Gamma'' \rightarrow \Gamma'$, $f : \Gamma' \rightarrow \Gamma$ and $s \in \widetilde{Ob}_1(\Gamma)$ one has

$$g^*(f^*(s)) = (g \circ f)^*(s). \tag{29}$$

i.e., that the maps f^* define on the family of sets \widetilde{Ob}_1 the structure of a presheaf. We continue using the notation \widetilde{Ob}_1 for this presheaf.

5.2. CONSTRUCTION. Denote by $PreShv(CC)$ the category of presheaves on the precategory underlying CC , i.e., the category of contravariant functors from the precategory underlying CC to $Sets$.

Let Ob_1 be the presheaf that takes an object $\Gamma \in CC$ to the set $Ob_1(\Gamma)$ and a morphism $f : \Gamma' \rightarrow \Gamma$ to the map $\Delta \mapsto f^*(\Delta)$. It is a functor due to the composition and unity axioms for f^* .

Let \widetilde{Ob}_1 be the presheaf that takes Γ to $\widetilde{Ob}_1(\Gamma)$ described above.

Let further $\partial : \widetilde{Ob}_1 \rightarrow Ob_1$ be the morphism that takes s to $\partial(s)$. It is well defined as a morphisms of families of sets and forms a morphism of presheaves since $\partial(f^*(s)) = f^*(\partial(s))$.

The morphism ∂ carries a universe structure that is defined by the standard pull-back squares in the category of presheaves.

We are going to construct a homomorphism $CC \rightarrow CC(PreShv(CC), \partial)$ using Construction 4.11 and to show that it is an isomorphism using Lemma 4.13.

We set Yo to be the Yoneda embedding.

We set

$$v_\Gamma : Ob_1(\Gamma) \rightarrow Hom_{PreShv}(Yo(\Gamma), Ob_1)$$

to be the standard bijections between sections of the presheaf Ob_1 on an object Γ and morphisms from the corresponding representable presheaf $Yo(\Gamma)$ to Ob_1 in the category of presheaves. It follows easily from the definitions that for $f : \Gamma' \rightarrow \Gamma$ and $\Delta \in Ob_1(\Gamma)$ one has

$$v_{\Gamma'}(f^*(\Delta)) = Yo(f) \circ v_\Gamma(\Delta) \tag{30}$$

We also set

$$\tilde{v}_\Gamma : \widetilde{Ob}_1(\Gamma) \rightarrow Hom_{PreShv}(Yo(\Gamma), \widetilde{Ob}_1)$$

to be the bijections of the same form for \widetilde{Ob}_1 . Again, it follows easily from the definitions that for $f : \Gamma' \rightarrow \Gamma$ and $s \in \widetilde{Ob}_1(\Gamma)$ one has

$$\tilde{v}_{\Gamma'}(f^*(s)) = Yo(f) \circ \tilde{v}_\Gamma(s) \tag{31}$$

To construct γ we first need to prove a lemma. Recall that for $\Delta \in CC$ such that $l(\Delta) > 0$ we let $\delta(\Delta) : \Delta \rightarrow p_\Delta^*(\Delta)$ denote the section of $p_{p_\Delta^*(\Delta)}$ given by the diagonal. We have

$$\delta(\Delta) \in \widetilde{Ob}_1(\Delta)$$

5.3. LEMMA. *Let $\Gamma \in Ob(CC)$ and $\Delta \in Ob_1(\Gamma)$. Then the square*

$$\begin{array}{ccc} Yo(\Delta) & \xrightarrow{\tilde{v}(\delta(\Delta))} & \widetilde{Ob}_1 \\ Yo(p_\Delta) \downarrow & & \downarrow \partial \\ Yo(\Gamma) & \xrightarrow{v(\Delta)} & Ob_1 \end{array} \tag{32}$$

is a pull-back square.

PROOF. We have to show that for any $\Gamma' \in CC$ the function

$$Hom(\Gamma', \Delta) \rightarrow Hom(\Gamma', \Gamma) \times_{Ob_1(\Gamma')} \widetilde{Ob}_1(\Gamma') \tag{33}$$

defined by the square (32) is a bijection. Unfolding the definitions we see that this function sends $g : \Gamma' \rightarrow \Delta$ to the pair $(g \circ p_\Delta, g^*(\delta(\Delta)))$ and that the fiber product is relative to the function from $Hom(\Gamma', \Gamma)$ to $Ob_1(\Gamma')$ that sends f to $f^*(\Delta)$ and the function from $\widetilde{Ob}_1(\Gamma')$ to $Ob_1(\Gamma')$ that sends s to $\partial(s)$.

Note that $g^*(\delta(\Delta)) = s_g$ where s is the s -operation of C-systems (see [9, Definition 2.3]) and $g \circ p_\Delta$ is the morphism that we denoted in [9] by $ft(g)$.

Let $f_1, f_2 : \Gamma' \rightarrow \Delta$ be two morphisms such that their images under (33) coincide, i.e., such that $ft(f_1) = ft(f_2)$ and $s_{f_1} = s_{f_2}$. This implies that $f_1 = f_2$ in view of [9, Definition 2.3(3)]. Therefore the function (33) is injective.

Let $f : \Gamma' \rightarrow \Gamma$ be a morphism and $s \in \widetilde{Ob}_1(\Gamma')$ a section such that $\partial(s) = f^*(\Delta)$. Then the composition $s \circ q(f, \Delta)$ is a morphism $f' : \Gamma' \rightarrow \Delta$ such that $f' \circ p_\Delta = f$. We also have

$$s_{f'} = s_{s \circ q(f, \Delta)} = s_s = s$$

which proves that (32) is surjective. This completes the proof of Lemma 5.3. ■

Let $\Gamma \in Ob(CC)$ and $\Delta \in Ob_1(\Gamma)$. By construction, $(Yo(\Gamma); v(\Delta))$ is the standard fiber product of the morphisms $v(\Delta)$ and ∂ in the category of presheaves. On the other hand $Yo(\Delta)$ is a fiber product of the same two morphisms by Lemma 5.3. Therefore there exists a unique isomorphism

$$\gamma_\Delta : (Yo(\Gamma); v(\Delta)) \rightarrow Yo(\Delta)$$

such that

$$\gamma_\Delta \circ \tilde{v}(\delta(\Delta)) = Q(v(\Delta)) \tag{34}$$

and

$$\gamma_\Delta \circ Yo(p_\Delta) = p_{Yo(\Gamma), v(\Delta)} \tag{35}$$

It remains to verify the four conditions of Problem 4.10 since the conditions of Lemma 4.13(2) are obviously satisfied.

We have that $Yo(pt) \rightarrow pt$ is an isomorphism.

The second condition is (30).

The third condition is (35).

It remains to verify the fourth condition. Let $f : \Gamma' \rightarrow \Gamma$ and $\Delta \in Ob_1(\Gamma)$. We need to show that

$$\gamma_{f^*(\Delta)} \circ Yo(q(f, \Delta)) = Q(Yo(f), v(\Delta)) \circ \gamma_\Delta \tag{36}$$

Two of the morphisms that are involved in the condition can be seen on the diagram

$$\begin{array}{ccccccc}
 (Yo(\Gamma'); v(f^*(\Delta))) & \xrightarrow{\gamma_{f^*(\Delta)}} & Yo(f^*(\Delta)) & \xrightarrow{Yo(q(f, \Delta))} & Yo(\Delta) & \xrightarrow{\tilde{v}(\delta(\Delta))} & \widetilde{Ob}_1 \\
 \downarrow p_{Yo(\Gamma'), v(f^*(\Delta))} & & \downarrow Yo(p_{f^*(\Delta)}) & & \downarrow Yo(p_\Delta) & & \downarrow p \\
 Yo(\Gamma') & \xlongequal{\quad} & Yo(\Gamma') & \xrightarrow{Yo(f)} & Yo(\Gamma) & \xrightarrow{v(\Delta)} & Ob_1
 \end{array}$$

By Lemma 5.3, $Yo(\Delta)$ is a fiber product with the projections $Yo(p_\Delta)$ and $\tilde{v}(\delta(\Delta))$. Therefore it is sufficient to verify that the compositions of the two sides of (36) with the projections are equal, i.e., we have to prove two equalities:

$$\gamma_{f^*(\Delta)} \circ Yo(q(f, \Delta)) \circ Yo(p_\Delta) = Q(Yo(f), v(\Delta)) \circ \gamma_\Delta \circ Yo(p_\Delta) \tag{37}$$

and

$$\gamma_{f^*(\Delta)} \circ Yo(q(f, \Delta)) \circ \tilde{v}(\delta(\Delta)) = Q(Yo(f), v(\Delta)) \circ \gamma_\Delta \circ \tilde{v}(\delta(\Delta)) \tag{38}$$

For (37) we have

$$\begin{aligned} \gamma_{f^*(\Delta)} \circ Yo(q(f, \Delta)) \circ Yo(p_\Delta) &= \gamma_{f^*(\Delta)} \circ Yo(p_{f^*(\Delta)}) \circ Yo(f) = \\ p_{Yo(\Gamma'), v(f^*(\Delta))} \circ Yo(f) &= p_{Yo(\Gamma'), Yo(f) \circ v(\Delta)} \circ Yo(f) \end{aligned}$$

where the first equality is by the commutativity of the canonical squares in CC and the fact that Yo is a functor, the second by (34) and the third one by (30). On the other hand

$$Q(Yo(f), v(\Delta)) \circ \gamma_\Delta \circ Yo(p_\Delta) = Q(Yo(f), v(\Delta)) \circ p_{Yo(\Gamma), v(\Delta)} = p_{Yo(\Gamma'), Yo(f) \circ v(\Delta)} \circ Yo(f)$$

where the first equality is by (34) and the second by the commutativity of the squares (2).

For (38) we have

$$\begin{aligned} \gamma_{f^*(\Delta)} \circ Yo(q(f, \Delta)) \circ \tilde{v}(\delta(\Delta)) &= \gamma_{f^*(\Delta)} \circ \tilde{v}(q(f, \Delta)^*(\delta(\Delta))) = \\ \gamma_{f^*(\Delta)} \circ \tilde{v}(\delta(f^*(\Delta))) &= Q(v(f^*(\Delta))) \end{aligned}$$

where the first equality is by (31), the second follows from a simple computation in CC and the third one is by (34). On the other hand one has

$$Q(Yo(f), v(\Delta)) \circ \gamma_\Delta \circ \tilde{v}(\delta(\Delta)) = Q(Yo(f), v(\Delta)) \circ Q(v(\Delta)) = Q(Yo(f) \circ v(\Delta)) = Q(f^*(\Delta))$$

where the first equality is by (34), the second one by Lemma 2.5 and the third one by (30).

This completes Construction 5.2.

The second construction that we provide for Problem 5.1 is as follows. For a set M let $Rp_n(M)$ be the set of subsets of $(\dots(M \times M) \times \dots) \times M$ where M occurs in the expression $n + 1$ times. Let $Rp(M)$ be the category with

$$Ob(Rp(M)) = \coprod_{n \geq 0} Rp_n(M)$$

$$Mor(Rp(M)) = \coprod_{(m,X),(n,Y)} Fun(X, Y)$$

where $Fun(X, Y)$ is the set of functions from X to Y and the identity morphisms and compositions of morphisms are given in the obvious way.

If our theory has a universe then the category *Sets* is defined and there is a functor $FF : Rp(M) \rightarrow Sets$ that sends (n, X) to X and acts on morphisms in the obvious way. This functor is fully faithful. (Note that it is not an inclusion of categories since, for example, the empty subset is an element of each of $Rp_n(M)$ so that there are objects (n, \emptyset) for all n which are all mapped by FF to the one empty set of *Sets*).

Let CC be a C-system. Consider $\mathcal{C} = Funct(CC^{op}, Rp(Mor(CC)))$. Define a universe $(p : \tilde{O} \rightarrow O, p_-, Q(-))$ in \mathcal{C} as follows.

For $\Gamma \in CC$ let

$$Ob'_1(\Gamma) = \{f \in Mor(CC) \mid f = Id_{codom(f)}\}$$

and let $\iota_\Gamma : Ob'_1(\Gamma) \rightarrow Ob_1(\Gamma)$ be the bijection defined by the codomain function *codom*. These bijections together with the structure of a presheaf on the family of sets $Ob_1(\Gamma)$ define a structure of a presheaf on the family of sets $Ob'_1(\Gamma)$. Let

$$O(\Gamma) = (0, Ob'_1(\Gamma)) \in Rp(Mor(CC))$$

The structure of a presheaf of sets on Ob'_1 defines a structure of an element of \mathcal{C} on O .

Next let

$$\tilde{O}(\Gamma) = (0, \tilde{Ob}_1(\Gamma)) \in Rp(Mor(C))$$

The structure of a presheaf on \tilde{Ob}_1 provide \tilde{O} with a structure of an object of \mathcal{C} . The morphism of presheaves ∂ defines in an obvious way a morphism $p : \tilde{O} \rightarrow O$ in \mathcal{C} . Let us construct a universe structure on p .

Let $X \in \mathcal{C}$ and $F : X \rightarrow O$. For $\Gamma \in CC$ let $X(\Gamma) = (n, X_0(\Gamma))$ where $X_0(\Gamma)$ is an element of $Rp_n(Mor(CC))$. Then

$$(X; F)_0(\Gamma) = \{(x_0, s) \in X_0(\Gamma) \times \tilde{Ob}_1(\Gamma) \mid F(x_0) = Id_{codom(s)}\} \in Rp_{n+1}(Mor(CC))$$

is an element of $Rp_{n+1}(Mor(CC))$ and $(X; F)(\Gamma) = (n + 1, (X; F)_0(\Gamma))$ is an element of $Rp(Mor(CC))$.

It is easy to equip the family $(X; F)(\Gamma)$ of elements of $Rp(Mor(CC))$ with a structure of an object of \mathcal{C} and equally easy to define morphisms $p_{X,F} : (X; F) \rightarrow X$ and $Q(F) : (X; F) \rightarrow \tilde{O}$.

We have also a functor $I : CC \rightarrow \mathcal{C}$ that extends the family of sets

$$X \mapsto (Y \mapsto (0, Mor(X, Y)))$$

The image of the final object of CC under this functor is a final object in \mathcal{C} which completes the description of a universe category structure on \mathcal{C} .

5.4. LEMMA. For any $F : X \rightarrow O$ the square

$$\begin{array}{ccc} (X; F) & \xrightarrow{Q(F)} & \tilde{O} \\ p_{X,F} \downarrow & & \downarrow p \\ X & \xrightarrow{F} & O \end{array} \tag{39}$$

is a pull-back square in \mathcal{C} .

PROOF. One can either give a direct proof which would not require an extra universe or one can argue that the functor FF defines a functor $\Phi : \mathcal{C} \rightarrow PreShv(CC)$ which is fully faithful and which maps squares (39) to standard pull-back squares in the category of presheaves of sets. ■

5.5. PROBLEM. *To construct an isomorphism $H : CC \rightarrow CC(\mathcal{C}, p)$ where (\mathcal{C}, p) is the universe category constructed above.*

There are two constructions for this problem. One we don't describe here because giving its detailed description would take a lot of space and add little understanding. It is a direct construction based on Construction 4.11 and Lemma 4.13 that parallels Construction 5.2. This direct construction would not use any extra universes and, in combination with the construction of (\mathcal{C}, p) based on the direct proof of Lemma 5.4 would provide a construction for Problem 5.1 that does not require any additional universes.

The construction that we give below uses Construction 5.2 and therefore requires an extra universe.

5.6. CONSTRUCTION. Let $\Phi : \mathcal{C} \rightarrow PreShv(CC)$ be the functor defined by FF . Since FF is fully faithful so is Φ . We have, by definition

$$\begin{aligned} \Phi(O) &= Ob'_1 \\ \Phi(\tilde{O}) &= \tilde{Ob}_1 \end{aligned}$$

The bijections ι_Γ give us an isomorphism of presheaves

$$\iota : \Phi(O) = Ob'_1 \rightarrow Ob_1$$

which commute with $\Phi(p)$ and ∂ and together with Φ form a universe category functor $(\Phi, \iota, Id_{\tilde{Ob}_1})$. This universe category functor satisfies the conditions of Lemma 4.8(2) and therefore the homomorphism $H(\Phi, \iota, Id_{\tilde{Ob}_1})$ is an isomorphism. Composing the isomorphism of Construction 5.2 the inverse to this isomorphism we obtain a solution to Problem 5.5.

The direct construction of the universe category (\mathcal{C}, p) does not increase the universe level but it uses the operation of taking the set of subsets that in type theory requires the propositional resizing rule in order to be defined inside a given universe. Here is an outline of a third construction that gives an even “tighter” universe category (\mathcal{C}, p) with an isomorphism $CC \rightarrow CC(\mathcal{C}, p)$.

5.7. CONSTRUCTION. Define by induction on n pairs (C_n, Φ_n) where C_n is a set and $\Phi_n : C_n \rightarrow PreShc(CC)$ is a function as follows:

1. for $n = 0$ we set $C_0 = \{pt, U, \tilde{U}\}$ and

$$\begin{aligned} \Phi_0(pt) &= pt \\ \Phi_0(U) &= Ob_1 \end{aligned}$$

$$\Phi_0(\tilde{U}) = \tilde{Ob}_1$$

where on the right hand side of the first equality is the final object of $PreShv(CC)$,

2. for the successor of n we set

$$C_{n+1} = \coprod_{X \in C_n} Hom_{PreShv(CC)}(\Phi_n(X), Ob_1)$$

and

$$\Phi_{n+1}(X, F) = (\Phi_n(X); F)$$

where $(X; F)$ is defined using standard fiber products in $PreShv(CC)$.

We then define

$$Ob(\mathcal{C}) = \coprod_{n \geq 0} C_n$$

$$Mor(\mathcal{C}) = \coprod_{(m,X),(n,Y) \in Ob(\mathcal{C})} Hom_{PreShv(CC)}(\Phi_m(X), \Phi_n(Y))$$

The composition and the identity morphisms are defined in such a way as to make the pair of maps

$$\Phi_{Ob} = \coprod_n \Phi_n$$

$$\Phi_{Mor} = \coprod_{(m,X),(n,Y) \in Ob(\mathcal{C})} i(\Phi_m(X), \Phi_n(Y))$$

where $i(F, G)$ is the inclusion of $Hom_{PreShv(CC)}(F, G)$ into $Mor(PreShv(CC))$, into a functor. This functor, which we denote by Φ , is then fully faithful.

One proves easily that pt is a final object of \mathcal{C} . One defines the universe morphism in \mathcal{C} as the morphism $p : \tilde{U} \rightarrow U$ that is mapped by Φ to ∂ . Given $(m, X) \in Ob(\mathcal{C})$ and a morphism $F : (m, X) \rightarrow U$ one defines $((m, X); F)$ as $(m + 1, (X, \Phi(F)))$. This object is a vertex of the square

$$\begin{CD} ((m, X); F) @>Q(F)>> \tilde{U} \\ @Vp(m,X,F)VV @VVpV \\ (m, X) @>F>> U \end{CD} \tag{40}$$

that is defined by the condition that it is mapped by Φ to the square

$$\begin{CD} (\Phi_m(X); \Phi(F)) @>Q(\Phi(F))>> \tilde{Ob}_1 \\ @Vp_{\Phi_m(X), \Phi(F)}VV @VVpV \\ \Phi_m(X) @>\Phi(F)>> \tilde{Ob}_1 \end{CD} \tag{41}$$

Since Φ is fully faithful and the square (41) is a pull-back square, the square (40) is a pull-back square. This provides us with a universe structure on p and completes the construction of the universe category (\mathcal{C}, p) .

The functor Φ together with two identity morphisms forms a universe category functor $\mathbf{\Phi} = (\Phi, Id_{Ob_1}, Id_{\widetilde{Ob_1}})$ that satisfies the conditions of Lemma 4.13(2). Therefore $\mathbf{\Phi}$ defines an isomorphism

$$CC(\mathcal{C}, p) \rightarrow CC(PreShv(CC), \partial)$$

composing the isomorphism of Construction 5.2 with the inverse to this isomorphism we obtain a solution to Problem 5.1. This completes Construction 5.7.

5.8. REMARK. The category \mathcal{C} of Construction 5.7 has all of the structures of a C-system and these structures satisfy all of the required properties except for the property that $l^{-1}(0) = \{pt\}$. We would like to call such objects “generalized C-systems”. They seem to appear also in other examples and may play an important role in the future.

6. A universe category defined by a precategory

The following problem was inspired by a question from an anonymous referee of [9]. Here we have to use the word precategory as in the definition of a C-system since the construction for this problem is not invariant under equivalences. Let us recall the following definition that also introduces the notations to be used below.

6.1. DEFINITION. *A category with fiber products is a category together with, for all pairs of morphisms of the form $f : X \rightarrow Z, g : Y \rightarrow Z$, fiber squares*

$$\begin{array}{ccc} (X, f) \times_Z (Y, g) & \xrightarrow{pr_2^{(X,f),(Y,g)}} & Y \\ pr_1^{(X,f),(Y,g)} \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

We will often abbreviate these main notations in various ways. The morphism $pr_2 \circ g = pr_1 \circ f$ from $(X, f) \times (Y, g)$ to Z is denoted by $f \diamond g$.

6.2. PROBLEM. *Let C be a precategory with a final object pt and fiber products. To construct a C-system CC and an equivalence of categories $J_* : CC \rightarrow C, J^* : C \rightarrow CC$.*

6.3. REMARK. In the univalent formalization an extra condition that C is a set-category, i.e., that the type of objects of C is a set is required for the construction that we provide to work.

6.4. REMARK. The construction that we provide increases the universe level. It might be possible to provide another construction for Problem 6.2 that does not increase the universe level.

6.5. REMARK. Note that if we required an isomorphism $CC \rightarrow C$ then the problem would have no solution since, for example, there is no C -system whose set of objects is the set with two elements. Indeed, one of these elements, let us denote it by X , will have to have length $n > 0$. Then $l(p_X^*(X)) = l(X) + 1$. Therefore $p_X^*(X) \neq X$ and $p_X^*(X) \neq pt$ which contradicts the assumption that CC has only two objects.

We start with a general construction that does not require C to have fiber products or a final object. The parts of it that do not concern C -systems must certainly have been known for a long time but we do not know where it was originally introduced.

For a precategory C let U_C be the presheaf such that

$$U_C(X) = \{(f, g) \text{ where } f : X \rightarrow Y \text{ and } g : Z \rightarrow Y\}$$

and for $a : X' \rightarrow X$,

$$U_C(a)(f, g) = (a \circ f, g)$$

One proves easily that this presheaf data defines a presheaf.

Let \tilde{U}_C be the presheaf such that

$$\tilde{U}_C(X) = \{(f', g) \text{ where } f' : X \rightarrow Z \text{ and } g : Z \rightarrow Y\}$$

and for $a : X' \rightarrow X$,

$$\tilde{U}_C(a)(f', g) = (a \circ f', g)$$

Again one proves easily that \tilde{U}_C this presheaf data defines a presheaf.

Let $p_C : \tilde{U}_C \rightarrow U_C$ be the morphism given by

$$(p_C)_X(f', g) = (f' \circ g, g)$$

One proves easily that this family of maps of sets is a morphism of presheaves.

As in Construction 5.2 let Yo be the Yoneda embedding and let

$$v_X : U_C(X) \rightarrow Hom_{PreShv}(Yo(X), U_C)$$

$$\tilde{v}_X : \tilde{U}_C(X) \rightarrow Hom_{PreShv}(Yo(X), \tilde{U}_C)$$

be the standard bijections which we will often write as v and \tilde{v} .

6.6. LEMMA. For $(f, g) \in U_C(X)$, $(f', g') \in \tilde{U}_C(X')$ and $u : X' \rightarrow X$ the square

$$\begin{array}{ccc} Yo(X') & \xrightarrow{\tilde{v}(f', g')} & \tilde{U}_C \\ Yo(u) \downarrow & & \downarrow p_C \\ Yo(X) & \xrightarrow{v(f, g)} & U_C \end{array} \quad (42)$$

commutes if and only if $g' = g$ and the square

$$\begin{array}{ccc}
 X' & \xrightarrow{f'} & Z \\
 u \downarrow & & \downarrow g \\
 X & \xrightarrow{f} & Y
 \end{array} \tag{43}$$

commutes. The square (42) is a pull-back square if and only if $g = g'$ and the square (43) is a pull-back square.

PROOF. The assertion about commutativity is obvious. The proof of the assertion about being a pull-back square is as follows. The square (6.6) is a pull-back square if and only if for all X'' the corresponding square of sections on X'' is a pull-back square of sets. This square of sections is of the form

$$\begin{array}{ccc}
 \text{Hom}(X'', X') & \xrightarrow{r_1} & \{(f'_0 : X'' \rightarrow Z_0, g_0 : Z_0 \rightarrow Y_0)\} \\
 s_1 \downarrow & & \downarrow r_2 \\
 \text{Hom}(X'', X) & \xrightarrow{s_2} & \{(f_0 : X'' \rightarrow Y_0, g_0 : Z_0 \rightarrow Y_0)\}
 \end{array} \tag{44}$$

where $r_1(a') = (a' \circ f', g')$, $r_2(f'_0, g'_0) = (f'_0 \circ g'_0, g'_0)$, $s_1(a) = a \circ u$, $s_2(a) = (a \circ f, g)$.

To check that (44) is a pull-back square it is sufficient to check that for every $a \in \text{Hom}(X'', X)$ the map $s_1^{-1}(a) \rightarrow r_2^{-1}(s_2(a))$ defined by r_1 is a bijection. We have

$$s_1^{-1}(a) = \{a' : X'' \rightarrow X' \mid a' \circ u = a\}$$

and

$$r_2^{-1}(s_2(a)) = \{(v : X'' \rightarrow Z, g) \mid v \circ g = a \circ f\} \tag{45}$$

and the map defined by r_1 maps a' to $(a' \circ v, g)$.

Applying the same reasoning to the condition that the square is (43) is pull-back we see that it is equivalent to the condition that for all X'' and all $a : X'' \rightarrow X$ the map from the set

$$(- \circ u)^{-1}(a) = \{a' : X'' \rightarrow X' \mid a' \circ u = a\}$$

to the set

$$(- \circ g)^{-1}(a \circ f) = \{v : X'' \rightarrow Z \mid v \circ g = a \circ f\} \tag{46}$$

given by $a' \mapsto a' \circ f'$, is a bijection. Since g in (45) the sets on the right hand sides of (45) and (46) are in the obvious bijection that is compatible with the functions from $\{a' : X'' \rightarrow X' \mid a' \circ u = a\}$ and therefore these two conditions are equivalent. ■

Applying our main construction to $(PreShv(CC), p_C)$ we obtain, for any precategory C , a C -system $CC(C) = CC(PreShv(CC), p_C)$.

6.7. **PROBLEM.** *Let C be a precategory with a final object pt . To construct a function $J_1^* : Ob(C) \rightarrow Ob_1(PreShv(C), p_C)$ and a family of isomorphisms $j_X : Yo(X) \rightarrow int(J^*(X))$.*

6.8. **CONSTRUCTION.** Let $X \in Ob(C)$. The pull-back square

$$\begin{array}{ccc} X & \xrightarrow{Id_X} & X \\ \pi_X \downarrow & & \downarrow \pi_X \\ pt & \xrightarrow{Id_{pt}} & pt \end{array}$$

defines by Lemma 6.6 a pull-back square

$$\begin{array}{ccc} Yo(X) & \xrightarrow{\tilde{v}(Id_X, \pi_X)} & \tilde{U}_C \\ Yo(\pi_X) \downarrow & & \downarrow p_C \\ Yo(pt) & \xrightarrow{v(Id_{pt}, \pi_X)} & U_C \end{array} \quad (47)$$

Let $\psi : pt \rightarrow Yo(pt)$ be the unique isomorphism. Set

$$J^*(X) = (pt, \psi \circ v(Id_{pt}, \pi_X))$$

Then

$$\pi_{Yo(X)} \circ \psi = Yo(\pi_X)$$

and therefore $\pi_{Yo(X)} * \tilde{v}(Id_X, \pi_X)$ is a well defined morphism from $Yo(X)$ to $J^*(X)$. It is easy to prove now that since (47) is a pull-back square this morphism is an isomorphism.

6.9. **REMARK.** For C with a final object pt , the set $Ob_1(PreShv(CC), p_C)$ is in a constructive bijection with the set of pairs $(f : pt \rightarrow Y, g : Z \rightarrow Y)$ which is given, in the notation of Construction 6.8, by the map $(f, g) \mapsto \psi \circ v(f, g)$. After composition with this bijection the function J^* takes X to $(Id : pt \rightarrow pt, \pi_X : X \rightarrow pt)$. The function $(f, g) \mapsto dom(g)$ defines a one-sided inverse to J_1^* so that J_1^* is always a split monomorphism.

6.10. **PROBLEM.** *Suppose that C is a category with a final object pt and fiber products. To construct a function $J_* : Ob(CC(C)) \rightarrow Ob(C)$ and for every $\Gamma \in Ob(CC(C))$ an isomorphism $\sigma_\Gamma : Yo(J_*(\Gamma)) \rightarrow int(\Gamma)$.*

6.11. **CONSTRUCTION.** We first construct by induction on n , pairs (J_n, σ_n) where

$$J_n : Ob_n(PreShv, p_C) \rightarrow Ob(C)$$

and σ_n is a family of isomorphisms

$$\sigma_n(A) : Yo(J_n(A)) \rightarrow int_n(A)$$

given for all $A \in Ob_n(PreShv, p_C)$ as follows (we write J instead of J_n and σ instead of σ_n):

1. For $n = 0$ we set $J(A) = pt$ and $\sigma(A) : Yo(pt) \rightarrow pt$ is the unique isomorphism,
2. For the successor of n we proceed as follows. Let $(B, F) \in Ob_{n+1}$ where $B \in Ob_n$ and $F : int(B) \rightarrow U_C$. Then

$$v^{-1}(\sigma_B \circ F) \in U_C(J(B))$$

is of the form

$$v^{-1}(\sigma_B \circ F) = (f : J(B) \rightarrow Z, g : Y \rightarrow Z)$$

Let

$$J(B, F) = (J(B), f) \times_Z (Y, g)$$

To define $\sigma(B, F)$ consider the diagram

$$\begin{array}{ccccccc} Yo(J(B, F)) & \xrightarrow{\iota} & (Yo(J(B)), \sigma(B) \circ F) & \xrightarrow{Q(\sigma(B), F)} & (int(B); F) & \xrightarrow{Q(F)} & \tilde{U}_C \\ Yo(pr_1) \downarrow & & p_{Yo(J(B)), \sigma(B) \circ F} \downarrow & & p_{int(B), F} \downarrow & & \downarrow p_C \\ Yo(J(B)) & \xlongequal{\quad} & Yo(J(B)) & \xrightarrow{\sigma(B)} & int(B) & \xrightarrow{F} & U_C \end{array}$$

where ι is the morphism $Yo(pr_1) * \tilde{v}(pr_2, g)$. This morphism is defined because

$$Yo(pr_1) \circ \sigma(B) \circ F = Yo(pr_1) \circ v(f, g) = (pr_1 \circ f, g)$$

and

$$\tilde{v}(pr_2, g) \circ p_C = (pr_2 \circ g, g)$$

and $pr_1 \circ f = pr_2 \circ g$. By Lemma 6.6 the square

$$\begin{array}{ccc} Yo(J(B, F)) & \xrightarrow{\tilde{v}(pr_2, g)} & \tilde{U}_C \\ Yo(pr_1) \downarrow & & \downarrow p_C \\ Yo(J(B)) & \xrightarrow{v(f, g)} & U_C \end{array}$$

is a pull-back square which implies that ι is an isomorphism. We define

$$\sigma(B, F) = \iota \circ Q(\sigma_B, F)$$

The morphism $Q(\sigma(B), F)$ is an isomorphism by Lemma 2.4 and therefore $\sigma(B, F)$ is an isomorphism.

We now define J_* as the sum over $n \in \mathbf{N}$ of J_n . This completes Construction 6.11.

6.12. REMARK. Using the bijection of Remark 6.9 we can look at the function from pairs $(f : pt \rightarrow Z, g : Z \rightarrow Y)$ to $Ob(C)$ corresponding to J_1 . This function is given by

$$(f, g) \mapsto (pt, f) \times_Z (Y, g)$$

When we compose it with J^* and consider $J_*(J^*(X))$ we obtain $(pt, Id_{pt}) \times_p t(X, \pi_X)$. Depending on the choice of the fiber product this element of $Ob(C)$ may be equal to X or not but in any case there is a natural in X isomorphism $X \rightarrow J_*(J^*(X))$.

We can now provide the following construction for Problem 6.2.

6.13. CONSTRUCTION. Let $J^* : Ob(C) \rightarrow Ob(CC(C))$ be the composition of the function J_1^* of Construction 6.8 with the inclusion of $Ob_1(PreShv, p_C)$ into $Ob(CC(PreShv, p_C))$. We can extend it to the data for a functor setting:

$$J_{Mor}^*(f : X \rightarrow Y) = int_{J^*(X), J^*(Y)}^{-1}(j_X^{-1} \circ Yo(f) \circ j_Y)$$

where $int_{\Gamma, \Gamma'}$ is the bijection

$$Hom_{CC(C)}(\Gamma, \Gamma') \rightarrow Hom(int(\Gamma), int(\Gamma'))$$

defined by the functor int . It is easy to prove from definitions that it is a functor and, using the fact that both Yo and int are fully faithful, that J^* is fully faithful.

Similarly we can extend J_* of Construction 6.11 to the data for a functor setting

$$(J_*)_{Mor}(f : \Gamma' \rightarrow \Gamma) = Yo_{J_*(\Gamma'), J_*(\Gamma)}^{-1}(\sigma_{\Gamma'} \circ int(f) \circ \sigma_{\Gamma}^{-1})$$

where $Yo_{X, Y}$ is the bijection

$$Hom_C(X, Y) \rightarrow Hom_{PreShv}(Yo(X), Yo(Y))$$

defined by the Yoneda embedding. Again it is easy to prove from definitions that this functor data is a functor and using the fact that both int and Yo are fully faithful that J_* is fully faithful.

After J_* and J^* have been extended to morphisms it makes sense to ask whether the families of isomorphisms j_X and σ_{Γ} are natural in X and Γ respectively and one verifies easily that they indeed are.

Let $X \in Ob(C)$ then we have an isomorphism

$$Yo(J_*(J^*(X))) \xrightarrow{\sigma_{J^*(X)}} int(J^*(X)) \xrightarrow{j_X^{-1}} Yo(X)$$

which is natural in X and applying to it $Yo_{J_*(J^*(X)), X}^{-1}$ we get an isomorphism

$$J_*(J^*(X)) \rightarrow X$$

which is again natural in X , i.e., we obtained a functor isomorphism $J_* \circ J^* \rightarrow Id$.

Similarly, starting with,

$$int(\Gamma) \xrightarrow{\sigma_{\Gamma}^{-1}} Yo(J_*(\Gamma)) \xrightarrow{j_{J_*(\Gamma)}} int(J^*(J_*(\Gamma)))$$

one obtains a functor isomorphism $Id \rightarrow J^* \circ J_*$. This completes the Construction 6.13.

6.14. REMARK. The construction of the C-system $CC(C)$ does not require a choice of a final object or fiber products in C and in particular does not depend on such choices.

The functor $J^* : C \rightarrow CC(C)$ requires a choice of a final object for its construction and depends on this choice. Let B be the bijection

$$Ob_1(PreShv(CC), p_C) \rightarrow \{(f : pt_1 \rightarrow Z, g : Y \rightarrow Z)\}$$

of Remark 6.9 defined by the choice of a final object pt_1 . Let $J_{1,1}^*$ and $J_{1,2}^*$ be the functions $Ob(C) \rightarrow Ob_1(PreShv, p_C)$ of Construction 6.7 defined by the choice of the final object pt_1 and a final object pt_2 respectively. Then one has

$$B(J_{1,1}^*(X)) = (Id : pt_1 \rightarrow pt_1, \pi_{X,1} : X \rightarrow pt_1)$$

and

$$B(J_{1,2}^*(X)) = (a : pt_1 \rightarrow pt_2, \pi_{X,2} : X \rightarrow pt_2)$$

where $a : pt_1 \rightarrow pt_2$ is the unique morphism. This shows that $J_{1,1}^* \neq J_{1,2}^*$ if $pt_1 \neq pt_2$ and in particular that J^* depends on the choice of the final object.

The fact that J_* depends on the choice of fiber products is seen from the formula for $J_*(J^*(X))$ given in Remark 6.12.

6.15. CONJECTURE. Let \mathcal{C} be a category, CC be a C-system and $M : CC \rightarrow \mathcal{C}$ a functor such that $M(pt_{CC})$ is a final object of \mathcal{C} and M maps distinguished squares of CC to pull-back squares of \mathcal{C} . Then there exists a universe $p_M : \tilde{U}_M \rightarrow U_M$ in $PreShv(\mathcal{C})$ and a C-system homomorphism $M' : CC \rightarrow CC(PreShv(\mathcal{C}), p_M)$ such that the square

$$\begin{array}{ccc} CC & \xrightarrow{M} & \mathcal{C} \\ \downarrow M' & & \downarrow \\ CC(PreShv(\mathcal{C}), p_M) & \xrightarrow{int} & PreShv(\mathcal{C}) \end{array}$$

where the right hand side vertical arrow is the Yoneda embedding, commutes up to a functor isomorphism.

References

- [1] Peter Aczel. A general Church-Rosser theorem. <http://www.ens-lyon.fr/LIP/REWRITING/MISC/AGeneralChurch-RosserTheorem.pdf>, 1978.
- [2] Benedikt Ahrens, Krzysztof Kapulkin, and Michael Shulman. Univalent categories and the Rezk completion. *Math. Structures Comput. Sci.*, 25(5):1010–1039, 2015.
- [3] John Cartmell. Generalised algebraic theories and contextual categories. *Ph.D. Thesis, Oxford University*, 1978. <https://uf-ias-2012.wikispaces.com/Semantics+of+type+theory>.

- [4] John Cartmell. Generalised algebraic theories and contextual categories. *Ann. Pure Appl. Logic*, 32(3):209–243, 1986.
- [5] Marcelo Fiore, Gordon Plotkin, and Daniele Turi. Abstract syntax and variable binding (extended abstract). In *14th Symposium on Logic in Computer Science (Trento, 1999)*, pages 193–202. IEEE Computer Soc., Los Alamitos, CA, 1999.
- [6] André Hirschowitz and Marco Maggesi. Modules over monads and linearity. In *Logic, language, information and computation*, volume 4576 of *Lecture Notes in Comput. Sci.*, pages 218–237. Springer, Berlin, 2007.
- [7] Vladimir Voevodsky. Notes on type systems. https://github.com/vladimirias/old_notes_on_type_systems, 2009–2012.
- [8] Vladimir Voevodsky. C-system of a module over a monad on sets. *arXiv 1407.3394*, submitted, pages 1–20, 2014.
- [9] Vladimir Voevodsky. Subsystems and regular quotients of C-systems. In *Conference on Mathematics and its Applications, (Kuwait City, 2014)*, number to appear, pages 1–11, 2015.

School of Mathematics, Institute for Advanced Study, Princeton NJ, USA.

Email: vladimir@math.ias.edu

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available from the journal's server at <http://www.tac.mta.ca/tac/>. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is \TeX , and $\text{\LaTeX}2\text{e}$ is required. Articles in PDF format may be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

\TeX EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT \TeX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva: valeria.depaiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Ieke Moerdijk, Radboud University Nijmegen: i.moerdijk@math.ru.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Université du Québec à Montréal : tierney.myles4@gmail.com

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca