

ALGEBRAIC EXPONENTIATION FOR LIE ALGEBRAS

XABIER GARCÍA-MARTÍNEZ AND JAMES R. A. GRAY

ABSTRACT. It is known that the category of Lie algebras over a ring admits *algebraic exponents*. The aim of this paper is to show that the same is true for the category of internal Lie algebras in an additive, cocomplete, symmetric, closed, monoidal category. In this way, we add some new examples to the brief list of known locally algebraically cartesian closed categories, including the categories of Lie superalgebras and differentially graded Lie algebras amongst others. Note that we are mainly interested in the case where the underlying category is abelian, as is the case in all our examples, but do not impose this condition since not requiring it adds no complexity to our arguments.

1. Introduction

Let \mathbb{C} be a finitely complete category. Given an object B of \mathbb{C} , we write $\mathbf{Pt}_B(\mathbb{C})$ for the *category of points over B* in which an object (A, p, s) is a split epimorphism $p: A \rightarrow B$ in \mathbb{C} , together with a chosen section $s: B \rightarrow A$, so that $p \circ s = 1_B$. A morphism of $\mathbf{Pt}_B(\mathbb{C})$ from (A, p, s) to (A', p', s') is a morphism $f: A \rightarrow A'$ in \mathbb{C} such that $p' \circ f = p$ and $f \circ s = s'$. For each morphism $q: E \rightarrow B$, there is a *change-of-base* functor $q^*: \mathbf{Pt}_B(\mathbb{C}) \rightarrow \mathbf{Pt}_E(\mathbb{C})$, that takes a point (A, p, s) to the point $(E \times_B A, \pi_1, \langle \text{id}, sq \rangle)$ obtained from the pullback

$$\begin{array}{ccc}
 E \times_B A & \xrightarrow{\pi_2} & A \\
 \langle \text{id}, sq \rangle \uparrow \pi_1 & \lrcorner & \uparrow s \\
 E & \xrightarrow{q} & B \\
 & & \downarrow p
 \end{array}$$

If for each morphism q in \mathbb{C} , the change-of-base functor q^* reflects isomorphisms, the category is said to be *Bourn-protomodular* [2]. If \mathbb{C} is pointed, this is equivalent to the same condition restricted to morphisms whose domain is the zero object. Note that the composition of the change-of-base functor $i_B^*: \mathbf{Pt}_B(\mathbb{C}) \rightarrow \mathbf{Pt}_0(\mathbb{C})$, where $i_B: 0 \rightarrow B$ is the unique morphism from the zero object to B , with the isomorphism $\mathbf{Pt}_0(\mathbb{C}) \cong \mathbb{C}$ is the kernel functor Ker_B sending each point (A, p, s) to the kernel of p . Therefore, a finitely

The first author was supported by Ministerio de Economía y Competitividad (Spain), with grant number MTM2016-79661-P and is a Postdoctoral Fellow of the Research Foundation–Flanders (FWO). The second author was supported in part by the National Research Foundation of South Africa (Grant Number 109203)

Received by the editors 2020-06-13 and, in final form, 2021-05-01.

Published on 2021-05-27 in the Rosebrugh Festschrift.

2020 Mathematics Subject Classification: 18E13, 16W25, 17A32, 18M05.

Key words and phrases: locally algebraically cartesian closed, semi-abelian category, algebraic exponentiation, Lie algebra.

© Xabier García-Martínez and James R. A. Gray, 2021. Permission to copy for private use granted.

complete pointed category is Bourn-protomodular if and only if for each B in \mathbb{C} the kernel functor reflects isomorphisms, or equivalently, the split short five lemma holds. Recall that a pointed protomodular category which is Barr-exact and has finite coproducts is called a *semi-abelian* category [20].

If the change-of-base functor q^* has a right adjoint for each morphism q , then \mathbb{C} is said to be *locally algebraically cartesian closed* ((LACC) for short) [5, 17, 19]. If \mathbb{C} is also protomodular and pointed, then by [19, Theorem 5.1] it is sufficient to check if for each B in \mathbb{C} the kernel functor $\text{Ker}_B: \text{Pt}_B(\mathbb{C}) \rightarrow \mathbb{C}$ has a right adjoint.

If \mathbb{C} is a (LACC), semi-abelian category, it implies several categorical-algebraic properties are satisfied by \mathbb{C} , such as *peri-abelianness* [4], *strong protomodularity* [3], the *Smith is Huq* condition [25], *normality of Higgins commutators* [8], and *algebraic coherence*.

As we can see, (LACC) is a strong condition where the short list of known semi-abelian examples includes groups, Lie algebras, crossed modules, cocommutative Hopf algebras over a field and all abelian categories. In fact, it was shown in [13] that (LACC) characterizes Lie algebras amongst all varieties of non-associative algebras over an infinite field of characteristic not equal to two. Furthermore, for an infinite field of characteristic two there are exactly two non-abelian (LACC) sub-varieties: Lie algebras and quasi-Lie algebras; that is, the variety of algebras obtained from Lie algebras by replacing the identity $[x, x] = 0$ by $[x, y] = -[y, x]$ (which is of course the same as $[x, y] = [y, x]$).

It is not known whether there is any relation between (LACC) and *action representability* [1]. They coincide in non-associative algebras over an infinite field (they both characterize Lie algebras [11]), but it is no longer true over finite fields, since Boolean algebras are action representable but not (LACC) [1, 19].

The aim of the present paper is to show that, under certain conditions, the category of Lie algebras in a monoidal category is (LACC). In doing so, we will add the categories of Lie superalgebras, \mathbb{Z} -graded Lie algebras and differentially graded Lie algebras, amongst others, to the brief list of known (LACC) examples. Another interesting example, is the category of Lie algebras in the Loday-Pirashvili category [23]. In this way, we will show that the category of Leibniz algebras (which is not (LACC) [12]) is a full reflective subcategory of a (LACC) category.

Let us briefly explain our approach, which is motivated by [18] where it is shown that the category of Lie algebras over a commutative ring is (LACC). Let \mathbf{C} be a symmetric monoidal closed category with underlying category additive and small complete. We show that for an object B in $\text{Lie}(\mathbf{C})$, the category of internal Lie algebras in \mathbf{C} , the category $\text{Pt}_B(\text{Lie}(\mathbf{C}))$ is equivalent to the category $\text{Lie}(\mathbf{D})$ of internal Lie algebras in the symmetric monoidal category \mathbf{D} with underlying category the category of *internal monoid actions* of $U(B)$, the *universal enveloping monoid* of B , acting on the objects of \mathbf{C} , and with monoidal structure induced by \mathbf{C} together with a natural comonoid structure on $U(B)$. Moreover, we show that the kernel functor $\text{Ker}_B: \text{Pt}_B(\text{Lie}(\mathbf{C})) \rightarrow \text{Lie}(\mathbf{C})$ factors, via this isomorphism, through a forgetful functor $V: \text{Lie}(\mathbf{D}) \rightarrow \text{Lie}(\mathbf{C})$. This reduces the problem of finding a right adjoint of Ker_B to finding a right adjoint of V , which can be found by an adjoint lifting theorem.

Note that although, when developing our theory, we don't assume that the underlying category is abelian, this is the case in all our examples. This means that semi-abelianess of the respective categories of Lie algebras in our examples amounts to the existence of coproducts, which can be constructed in each case in essentially the same way as for classical Lie algebras.

The paper is organized as follows: In Section 2 after recalling the definition of various relevant types of internal algebras we construct adjoint functors between certain categories of such internal algebras, which we will use in Section 4. In Section 3 we recall the necessary background and then prove the adjoint lifting theorem which we use to produce the right adjoint of the functor V mentioned above. The adjoint functors previously defined are used in Section 4 to produce the symmetric monoidal category \mathbf{D} as well as the equivalence of categories between $\text{Pt}_B(\text{Lie}(\mathbf{C}))$ and $\text{Lie}(\mathbf{D})$ mentioned before. Finally, in Section 5 we study some applications of the obtained result.

2. Internal algebras

Throughout the rest of paper we denote by $\mathbf{C} = (\mathbb{C}, \otimes, I, \alpha, \lambda, \rho, \sigma)$ a *symmetric monoidal category*. We refer to [24] for basic knowledge about this topic. Recall that \mathbf{C} is said to be *closed* if for each object X in \mathbb{C} the endofunctor $X \otimes - : \mathbb{C} \rightarrow \mathbb{C}$ is a left adjoint. A chosen right adjoint will be denoted by $(-)^X$ and the counit by ev . As the main non-cartesian example we have the symmetric monoidal category $\text{Vect}_{\mathbb{K}}$ of vector spaces with the tensor product $\otimes_{\mathbb{K}}$. The canonical isomorphism between $\text{Hom}_{\text{Vect}_{\mathbb{K}}}(X \otimes_{\mathbb{K}} Z, Y)$ and $\text{Hom}_{\text{Vect}_{\mathbb{K}}}(Z, \text{Hom}_{\text{Vect}_{\mathbb{K}}}(X, Y))$ gives us the right adjoint $(-)^X = \text{Hom}_{\text{Vect}_{\mathbb{K}}}(X, -)$. In this particular case, the counit of the adjunction $\text{ev}_Y : Y^X \otimes X \rightarrow Y$ maps $f \otimes x$ to the evaluation $f(x)$.

The setting we need in this manuscript is the following: \mathbf{C} is a symmetric monoidal closed category with underlying category \mathbb{C} additive and cocomplete.

Recall that a *monoid* (A, m, u) in \mathbf{C} is a triple, where A is an object of \mathbb{C} , $m : A \otimes A \rightarrow A$ and $u : I \rightarrow A$ are morphisms of \mathbb{C} , such that the diagrams

$$\begin{array}{ccc}
 A \otimes (A \otimes A) & \xrightarrow{\alpha} & (A \otimes A) \otimes A \\
 \text{id} \otimes m \downarrow & & \downarrow m \otimes \text{id} \\
 A \otimes A & \xrightarrow{m} & A \leftarrow m \quad A \otimes A
 \end{array}
 \qquad
 \begin{array}{ccc}
 A \otimes I & \xrightarrow{\text{id} \otimes u} & A \otimes A \xleftarrow{u \otimes \text{id}} I \otimes A \\
 \rho \searrow & & \downarrow m \\
 & & A \xleftarrow{\lambda}
 \end{array}$$

commute [24]. If in addition the diagram

$$\begin{array}{ccc}
 A \otimes A & \xrightarrow{\sigma} & A \otimes A \\
 m \searrow & & \swarrow m \\
 & & A
 \end{array}$$

commutes, then it is said to be a *commutative monoid*. The categories of such objects will be denoted by $\text{Mon}(\mathbf{C})$ and $\text{Mon}^c(\mathbf{C})$, respectively. Recall that since \mathbf{C} is a symmetric

monoidal category, given two monoids (A, m, u) and (A', m', u') in \mathbf{C} the triple $(A \otimes A', (m \otimes m')i, (u \otimes u')\lambda)$, where i is the *middle interchange isomorphism*, is a monoid which we denote by $(A, m, u) \otimes (A', m', u')$ (which is commutative whenever both (A, m, u) and (A', m', u') are). Moreover, with this definition of tensor product the symmetric monoidal structure of \mathbf{C} lifts to both $\mathbf{Mon}(\mathbf{C})$ and $\mathbf{Mon}^c(\mathbf{C})$. Let us write $\underline{\mathbf{Mon}}\mathbf{C}$ and $\underline{\mathbf{Mon}}^c\mathbf{C}$ for the respective symmetric monoidal categories. The duals of the notions of monoid and commutative monoid are *comonoid* and *cocommutative comonoid*, respectively, and the categories of such objects are denoted $\mathbf{CoMon}(\mathbf{C})$ and $\mathbf{CoMon}^{\text{coc}}(\mathbf{C})$, respectively. By duality the symmetric monoidal structure on \mathbf{C} lifts to both $\mathbf{CoMon}(\mathbf{C})$ and $\mathbf{CoMon}^{\text{coc}}(\mathbf{C})$, producing the symmetric monoidal categories $\underline{\mathbf{CoMon}}\mathbf{C}$ and $\underline{\mathbf{CoMon}}^{\text{coc}}\mathbf{C}$, respectively.

A *bimonoid* can be defined as a comonoid in the symmetric monoidal category $\mathbf{Mon}(\mathbf{C})$ (or equivalently as a monoid in $\mathbf{CoMon}(\mathbf{C})$). We present a bimonoid as a quintuple (A, m, u, d, e) where (A, m, u) is an monoid, (A, d, e) is a comonoid and $d: (A, m, u) \rightarrow (A, m, u) \otimes (A, m, u)$ and $e: (A, m, u) \rightarrow (I, \lambda, \text{id})$ are monoid morphisms, or equivalently m and u are comonoid morphisms. The category of bimonoids will be denoted by $\mathbf{BiMon}(\mathbf{C})$. A bimonoid is called *commutative* if the monoid part is commutative, and it is called *cocommutative* if the comonoid part is cocommutative.

Under the conditions assumed on \mathbf{C} it is well known that free monoids exist, in fact much milder assumptions are needed (see e.g. [22] and the references therein).

2.1. PROPOSITION. *The forgetful functor $G: \mathbf{Mon}(\mathbf{C}) \rightarrow \mathbf{C}$ has a left adjoint which we will denote by F .*

A *Lie algebra* in \mathbf{C} is a pair (X, b) where X is an object and $b: X \otimes X \rightarrow X$ is a morphism in \mathbf{C} making the diagrams

$$\begin{array}{ccc}
 X \otimes X & \xrightarrow{\text{id} + \sigma} & X \otimes X \\
 \downarrow 0 & \swarrow b & \\
 X & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 X \otimes (X \otimes X) & \xrightarrow{\text{id} + \sigma\alpha + (\sigma\alpha)^2} & X \otimes (X \otimes X) \\
 \downarrow 0 & & \downarrow \text{id} \otimes b \\
 X & \xleftarrow{b} & X \otimes X
 \end{array}
 \tag{1}$$


commute. Note that we can sum morphisms since we are assuming that \mathbf{C} is additive. The category of Lie algebras in \mathbf{C} will be denoted by $\mathbf{Lie}(\mathbf{C})$. As expected, when \mathbf{C} is the category of vector spaces (with the usual monoidal structure) Lie algebras in \mathbf{C} are usual Lie algebras.

Throughout this paper we will denote the direct sum of A and B by $A \oplus B$ with projections $\pi_1: A \oplus B \rightarrow A$ and $\pi_2: A \oplus B \rightarrow B$ and inclusions $\iota_1: A \rightarrow A \oplus B$ and $\iota_2: B \rightarrow A \oplus B$. For morphisms $f: W \rightarrow A$, $g: W \rightarrow B$ and $h: A \rightarrow C$ and $i: B \rightarrow C$ we will denote by $\langle f, g \rangle: W \rightarrow A \oplus B$ and $[h, i]: A \oplus B \rightarrow C$ the unique morphisms with $\pi_1 \langle f, g \rangle = f$, $\pi_2 \langle f, g \rangle = g$, $[h, i] \iota_1 = h$ and $[h, i] \iota_2 = i$.

Given a monoid (A, m, u) in \mathbf{C} , the pair $(A, m(\text{id} - \sigma))$ is an object in $\mathbf{Lie}(\mathbf{C})$ and this assignment determines the object map of a functor $L: \mathbf{Mon}(\mathbf{C}) \rightarrow \mathbf{Lie}(\mathbf{C})$ which is the identity on morphisms (see e.g. [18]). The following proposition is probably known, but we couldn't find a reference so we included a proof.

2.2. PROPOSITION. *The functor $L: \mathbf{Mon}(\mathbf{C}) \rightarrow \mathbf{Lie}(\mathbf{C})$ has a left adjoint, which we denote by U .*

PROOF. Let us write η for the unit of the adjunction $F \dashv G$ from Proposition 2.1. Let (B, b) be an object in $\mathbf{Lie}(\mathbf{C})$ and (A, m, u) an object in $\mathbf{Mon}(\mathbf{C})$. For a morphism $f: B \rightarrow A$ in \mathbf{C} let \bar{f} be the corresponding monoid morphism from $F(B) = (T(B), m_B, u_B)$ to (A, m, u) obtained via the adjunction $F \dashv G$, that is the unique monoid morphism such that $G(\bar{f})\eta = f$. By considering the diagram



ones sees that the sub-diagram $[1]$ commutes if and only if the outer arrows commute. This implies that f is a morphism from (B, b) to $L(A, m, u)$ in $\mathbf{Lie}(\mathbf{C})$ if and only if the diagram

$$B^{\otimes 2} \begin{array}{c} \xrightarrow{\eta b} \\ \xrightarrow{m_B(\eta \otimes \eta)(id - \sigma)} \end{array} T(B) \xrightarrow{\bar{f}} A$$

is a fork in \mathbf{C} . However, the previous diagram is a fork if and only if the diagram

$$B \oplus B^{\otimes 2} \begin{array}{c} \xrightarrow{[\eta, \eta b]} \\ \xrightarrow{[\eta, m_B(\eta \otimes \eta)(id - \sigma)]} \end{array} T(B) \xrightarrow{\bar{f}} A$$

is a fork. Therefore, writing $r, s: F(B \oplus B^{\otimes 2}) \rightarrow F(B)$ for the corresponding monoid morphisms induced by $[\eta, \eta b]$ and $[\eta, m_B(\eta \otimes \eta)(id - \sigma)]$, respectively, we see that previous diagram is a fork if and only if the diagram

$$F(B \oplus B^{\otimes 2}) \begin{array}{c} \xrightarrow{r} \\ \xrightarrow{s} \end{array} F(B) \xrightarrow{\bar{f}} (A, m, u)$$

is a fork in $\mathbf{Mon}(\mathbf{C})$. Since the tensor \otimes preserves coequalizers in each argument, it is well-known that it preserves reflexive coequalizers, and the forgetful functor $\mathbf{Mon}(\mathbf{C}) \rightarrow \mathbf{C}$ creates reflexive coequalizers. Noting that the parallel pair of morphisms in the previous diagram is reflexive it easily follows that if $q: F(B) \rightarrow U(B, b)$ is the coequalizer of r and s in $\mathbf{Mon}(\mathbf{C})$, then $U(B, b)$ together with the morphism $q\eta$ from (B, b) to $L(U(B, b))$ in $\mathbf{Lie}(\mathbf{C})$ is the universal morphism from (B, b) to L . ■

Next we show that for any B in $\text{Lie}(\mathbf{C})$ the monoid $U(B)$ is the underlying monoid of a cocommutative bimonoid (in fact of a cocommutative Hopf monoid) $\tilde{U}(B)$, and the assignment $B \mapsto \tilde{U}(B)$ is the object map of a functor which is left adjoint of a functor P which we construct below. This functor P generalizes the construction of the primitive Lie algebra of a bialgebra.

Recall, as mentioned above, that since \mathbf{C} is symmetric monoidal the monoidal structure lifts to $\text{Mon}(\mathbf{C})$.

2.3. PROPOSITION. *For each monoid $A = (A, m, u)$ the morphism $\delta_A: A \rightarrow A \otimes A$ defined by $\delta_A = (u \otimes \text{id})\lambda^{-1} + (\text{id} \otimes u)\rho^{-1}$ is a morphism from $L(A)$ to $L(A \otimes A)$ in $\text{Lie}(\mathbf{C})$, natural in A .*

PROOF. Let $A = (A, m, u)$ be a monoid in \mathbf{C} and let $\delta_A: A \rightarrow A \otimes A$ be the morphism $\delta_A = l_A + r_A$ where $l_A = (u \otimes \text{id})\lambda^{-1}$ and $r_A = (\text{id} \otimes u)\rho^{-1}$. The naturality of δ easily follows from the naturality of l and r . Therefore, it only remains to show that δ_A is a morphism from $L(A)$ to $L(A \otimes A)$. We have

$$\begin{aligned} (m \otimes m)i(l \otimes r) &= (m \otimes m)i((u \otimes \text{id}) \otimes (\text{id} \otimes u))(\lambda^{-1} \otimes \rho^{-1}) \\ &= (m \otimes m)((u \otimes \text{id}) \otimes (\text{id} \otimes u))i(\lambda^{-1} \otimes \rho^{-1}) \\ &= (\lambda \otimes \rho)i(\lambda^{-1} \otimes \rho^{-1}) \\ &= \sigma \end{aligned}$$

and

$$\begin{aligned} (m \otimes m)i\sigma(l \otimes r) &= (m \otimes m)i\sigma((u \otimes \text{id}) \otimes (\text{id} \otimes u))(\lambda^{-1} \otimes \rho^{-1}) \\ &= (m \otimes m)((\text{id} \otimes u) \otimes (u \otimes \text{id}))i\sigma(\lambda^{-1} \otimes \rho^{-1}) \\ &= (\rho \otimes \lambda)i\sigma(\lambda^{-1} \otimes \rho^{-1}) \\ &= \sigma \end{aligned}$$

and hence

$$(m \otimes m)i(\text{id} - \sigma)(l \otimes r) = 0$$

and

$$(m \otimes m)i(\text{id} - \sigma)(r \otimes l) = (m \otimes m)i(\sigma - \text{id})\sigma(r \otimes l) = -(m \otimes m)i(\text{id} - \sigma)(l \otimes r)\sigma = 0.$$

On the other hand it is easy to check that l_A and r_A are monoid morphisms from A to $A \otimes A$ and hence they are morphisms from $L(A)$ to $L(A \otimes A)$ in $\text{Lie}(\mathbf{C})$. Therefore

$$\begin{aligned} (m \otimes m)i(\text{id} - \sigma)(\delta_A \otimes \delta_A) &= (m \otimes m)i(\text{id} - \sigma)(l_A \otimes l_A) + (m \otimes m)i(\text{id} - \sigma)(l_A \otimes r_A) \\ &\quad + (m \otimes m)i(\text{id} - \sigma)(r_A \otimes l_A) + (m \otimes m)i(\text{id} - \sigma)(r_A \otimes r_A) \\ &= l_A m(\text{id} - \sigma) + 0 + 0 + r_A m(\text{id} - \sigma) \\ &= \delta_A m(\text{id} - \sigma) \end{aligned}$$

as desired. ■

Recall that a bimonoid (A, m, u, d, e) is called a *Hopf monoid* if there is a morphism $s: A \rightarrow A$ making the diagram

$$\begin{array}{ccccc}
 & & A \otimes A & \xrightarrow{\text{id} \otimes s} & A \otimes A & & \\
 & d \nearrow & & & & m \searrow & \\
 A & & & & & & A \\
 & e \longrightarrow & I & \xrightarrow{u} & & & \\
 & d \searrow & & & & m \nearrow & \\
 & & A \otimes A & \xrightarrow{s \otimes \text{id}} & A \otimes A & &
 \end{array}$$

commute. Recall also that such a morphism s is unique (whenever it exists) and is called the *antipode* of the Hopf monoid (A, m, u, d, e) . Note that Goyvaerts and Vercruyssen have studied a generalization of the functor P , below, in [15]. In fact, the following proposition can be found in [16, Proposition 2.8]

2.4. PROPOSITION. *Let H be the forgetful functor from $\mathbf{BiMon}(\mathbf{C}) \rightarrow \mathbf{Mon}(\mathbf{C})$. For a bimonoid A , with comultiplication $d_A: A \rightarrow A \otimes A$, the morphism denoted by $j_A: P(A) \rightarrow L(H(A))$ forming part of the equalizer diagram*

$$P(A) \xrightarrow{j_A} L(H(A)) \begin{array}{c} \xrightarrow{\delta_{H(A)}} \\ \xrightarrow{L(d_A)} \end{array} L(H(A) \otimes H(A)),$$

is the component of a natural transformation from a functor $P: \mathbf{BiMon}(\mathbf{C}) \rightarrow \mathbf{Lie}(\mathbf{C})$ to the functor LH . Moreover, the functor P is part of an adjunction with left adjoint \tilde{U} and unit $\tilde{\nu}$ such that: $H\tilde{U} = U$ and $(j \circ \tilde{U})\tilde{\nu} = \nu$ where ν is the unit of the adjunction $U \dashv L$. Furthermore, for each object X in $\mathbf{Lie}(\mathbf{C})$ the bimonoid $\tilde{U}(X)$ is a cocommutative Hopf monoid. ■

3. Monoidal functors and lifting of adjoints

Given two monoidal categories $\mathbf{C} = (\mathbf{C}, \otimes, I, \alpha, \lambda, \rho)$ and $\mathbf{C}' = (\mathbf{C}', \otimes', \alpha', \lambda', \rho')$ a *lax monoidal functor* from \mathbf{C} to \mathbf{C}' is a triple (F, θ, ϕ) where $F: \mathbf{C} \rightarrow \mathbf{C}'$ is a functor, $\theta: I' \rightarrow F(I)$ is a morphism in \mathbf{C}' , and $(\phi_{A,B}: F(A) \otimes' F(B) \rightarrow F(A \otimes B))_{A,B \in \mathbf{C}}$ is a natural transformation, such that for all A, B and C in \mathbf{C} the diagrams

$$\begin{array}{ccc}
 F(A) \otimes' (F(B) \otimes' F(C)) & \xrightarrow{\alpha'} & (F(A) \otimes' F(B)) \otimes' F(C) \\
 \text{id} \otimes' \phi \downarrow & & \downarrow \phi \otimes' \text{id} \\
 F(A) \otimes' F(B \otimes C) & & (F(A \otimes B) \otimes' F(C)) \\
 \phi \downarrow & & \downarrow \phi \\
 F(A \otimes (B \otimes C)) & \xrightarrow{F(\alpha)} & F((A \otimes B) \otimes C)
 \end{array}$$

$$\begin{array}{ccc}
 F(A) & \xrightarrow{\lambda'} & I' \otimes' F(A) \\
 & \searrow^{F(\lambda)} & \downarrow \theta \otimes' \text{id} \\
 & & F(I) \otimes' F(A) \\
 & & \downarrow \phi \\
 & & F(I \otimes A)
 \end{array}
 \qquad
 \begin{array}{ccc}
 F(A) \otimes' I' & \xleftarrow{\rho'} & F(A) \\
 \text{id} \otimes' \theta \downarrow & & \downarrow F(\rho) \\
 F(A) \otimes F(I) & & \\
 \phi \downarrow & & \\
 F(A \otimes I) & &
 \end{array}$$

commute. If \mathbf{C} and \mathbf{C}' are symmetric monoidal categories with *symmetry* isomorphisms σ and σ' respectively, then a lax monoidal functor (F, θ, ϕ) between \mathbf{C} and \mathbf{C}' is called a lax symmetric monoidal functor as soon as the diagram

$$\begin{array}{ccc}
 F(A) \otimes' F(B) & \xrightarrow{\sigma'} & F(B) \otimes' F(A) \\
 \phi \downarrow & & \downarrow \phi \\
 F(A \otimes B) & \xrightarrow{F(\sigma)} & F(B \otimes A)
 \end{array}$$

commutes, for every A and B in \mathbf{C} . A lax symmetric monoidal functor (F, θ, ϕ) is called strong when θ and ϕ are isomorphisms, and strict when they are identity morphisms.

Recall also that (symmetric) monoidal structures on categories lift to both functor categories and products of categories (componentwise). In particular, if \mathbf{C} is a (symmetric) monoidal category, then by $\mathbf{C} \times \mathbf{C}$ we will mean the (symmetric) monoidal category with underlying category $\mathbf{C} \times \mathbf{C}$, with tensor product defined by $(A, B) \otimes (A', B') = (A \otimes A', B \otimes B')$, and with remaining structure defined componentwise. For a symmetric monoidal category \mathbf{C} there is via the coherence theorem a unique natural isomorphism $i_{A,B,C,D}: (A \otimes B) \otimes (C \otimes D) \rightarrow (A \otimes C) \otimes (B \otimes D)$ built from α, λ, ρ and σ which we will call (middle) interchange isomorphism. Note that this produces a strong symmetric monoidal functor $(\otimes, \lambda, i): \mathbf{C} \times \mathbf{C} \rightarrow \mathbf{C}$.

It is well known that if $\mathbf{C} = (\mathbb{C}, \otimes, I, \alpha, \lambda, \rho)$ and $\mathbf{C}' = (\mathbb{C}', \otimes', \alpha', \lambda', \rho')$ are monoidal categories, and (F, θ, ϕ) is a lax monoidal functor $\mathbf{C} \rightarrow \mathbf{C}'$, then the assignment $(A, m, u) \mapsto (F(A), F(m)\phi, F(u)\theta)$ is the object map of a functor $\text{Mon}(\mathbf{C}) \rightarrow \text{Mon}(\mathbf{C}')$ which sends a morphism f to $F(f)$. In a similar way, when \mathbf{C}, \mathbf{C}' and F are also additive, the assignment $(B, b) \mapsto (F(B), F(b)\phi)$ is the object map of a functor $\text{Lie}(\mathbf{C}) \rightarrow \text{Lie}(\mathbf{C}')$ which sends a morphism f to $F(f)$. Let us denote this induced functor by $\text{Lie}(F)$. The following fact may be known, but we couldn't find a reference:

3.1. PROPOSITION. *Let $\mathbf{C} = (\mathbb{C}, \otimes, I, \alpha, \lambda, \rho)$ and $\mathbf{C}' = (\mathbb{C}', \otimes', \alpha', \lambda', \rho')$ be monoidal categories such that \mathbf{C} and \mathbf{C}' are additive, and let $F: \mathbf{C} \rightarrow \mathbf{C}'$ be a strict monoidal functor which is additive. If $F: \mathbf{C} \rightarrow \mathbf{C}'$ has a right adjoint, then $\text{Lie}(F)$ has a right adjoint.*

PROOF. Suppose F has a right adjoint G , and the unit and counit of the associated adjunction are denoted by η and ϵ , respectively. According to [21] we know that defining $\theta = \eta_I$, and for each A' and B' in \mathbf{C}' , $\phi_{A',B'}: G(A') \otimes G(B') \rightarrow G(A' \otimes' B')$ to be the

composite $G(\epsilon_{A'} \otimes' \epsilon_{B'})\eta_{G(A')\otimes G(B')}$ produces a lax monoidal functor $(G, \theta, \phi): \mathbf{C}' \rightarrow \mathbf{C}$, and furthermore, these data make the triangle in (2), below, commute. As explained above this lax functor determines a functor $\text{Lie}(G): \text{Lie}(\mathbf{C}') \rightarrow \text{Lie}(\mathbf{C})$. The claim now follows by observing that for objects (X, b) and (X', b') in $\text{Lie}(\mathbf{C})$ and $\text{Lie}(\mathbf{C}')$, respectively, and for $f: X \rightarrow G(X')$ a morphism in \mathbb{C} , either (and hence both) of the statements: $f: (X, b) \rightarrow (G(X), G(b')\phi)$ is a morphism in $\text{Lie}(\mathbf{C})$; and $\epsilon F(f): (F(X), F(b)) \rightarrow (X', b')$ is a morphism in $\text{Lie}(\mathbf{C}')$; are equivalent to the commutativity of the diagram

$$\begin{array}{ccccc}
 F(X) \otimes' F(X) & \xrightarrow{F(f)\otimes' F(f)} & FG(X') \otimes' FG(X') & & \\
 \parallel & & \parallel & \searrow^{\epsilon_{X'}\otimes'\epsilon_{X'}} & \\
 F(X \otimes X) & \xrightarrow{F(f\otimes f)} & F(G(X') \otimes G(X')) & & \\
 \downarrow F(b) & & \downarrow F(\phi_{X', X'}) & & \\
 F(X) & \xrightarrow{F(f)} & FG(X' \otimes' X') & \xrightarrow{\epsilon_{X'}\otimes' X'} & X' \otimes' X' \\
 & & \downarrow FG(b') & & \downarrow b' \\
 & & FG(X') & \xrightarrow{\epsilon_{X'}} & X'
 \end{array} \tag{2}$$

■

4. Actions

Recall that if $A = (A, m, u)$ is a monoid in \mathbf{C} and X is an object in \mathbb{C} an action of A on X is a morphism $\phi: A \otimes X \rightarrow X$ making the diagrams

$$\begin{array}{ccc}
 (A \otimes A) \otimes X & \xrightarrow{\alpha^{-1}} & A \otimes (A \otimes X) \\
 m \otimes \text{id} \downarrow & & \downarrow \text{id} \otimes \phi \\
 A \otimes X & \xrightarrow{\phi} & X \longleftarrow_{\phi} A \otimes X
 \end{array}
 \qquad
 \begin{array}{ccc}
 I \otimes X & \xrightarrow{u \otimes \text{id}} & A \otimes X \\
 & \searrow \lambda & \downarrow \phi \\
 & & X
 \end{array}$$

commute. Given an object $B = (B, b)$ in $\text{Lie}(\mathbf{C})$ and an object X in \mathbb{C} we say that a morphism $\theta: B \otimes X \rightarrow X$ is an action of B on X if the diagram

$$\begin{array}{ccc}
 (B \otimes B) \otimes X & \xrightarrow{\alpha^{-1}((\text{id}-\sigma)\otimes \text{id})} & B \otimes (B \otimes X) \\
 b \otimes \text{id} \downarrow & & \downarrow \text{id} \otimes \theta \\
 B \otimes X & \xrightarrow{\theta} & X \longleftarrow_{\theta} B \otimes X
 \end{array} \tag{3}$$

commutes. Let us write $E(X)$ for the monoid with underlying object X^X and with multiplication $m: X^X \otimes X^X \rightarrow X^X$ and $u: I \rightarrow X^X$ the unique morphisms making the

diagrams

$$\begin{array}{ccc}
 (X^X \otimes X^X) \otimes X & \xrightarrow{\alpha^{-1}} & X^X \otimes (X^X \otimes X) & I \otimes X & \xrightarrow{u \otimes \text{id}} & X^X \otimes X \\
 m \otimes \text{id} \downarrow & & \downarrow \text{id} \otimes \text{ev} & \searrow \lambda & & \downarrow \text{ev} \\
 X^X \otimes X & \xrightarrow{\text{ev}} & X & \xleftarrow{\text{ev}} & X^X \otimes X & \\
 & & & & &
 \end{array}$$

commute. Let $f: A' \rightarrow A$ be a morphism in $\text{Mon}(\mathbf{C})$, $g: B' \rightarrow B$ be a morphism in $\text{Lie}(\mathbf{C})$, $\phi: A \otimes X \rightarrow X$ be an action of A on X , and $\theta: B \otimes X \rightarrow X$ be an action of B on X , then $\phi(f \otimes \text{id})$ is an action of A' on X and $\theta(g \otimes \text{id})$ is an action of B' on X . We have

4.1. PROPOSITION. For an object X in \mathbf{C} there are natural isomorphisms between the following functors:

1. the functor assigning to each B in $\text{Lie}(\mathbf{C})^{\text{op}}$ the set of actions of B on X ;
2. $\text{hom}(-, L(E(X))): \text{Lie}(\mathbf{C})^{\text{op}} \rightarrow \text{Set}$;
3. $\text{hom}(U(-), E(X)): \text{Lie}(\mathbf{C})^{\text{op}} \rightarrow \text{Set}$;
4. the functor assigning to each B in $\text{Lie}(\mathbf{C})^{\text{op}}$ the set of monoid actions of $U(B)$ on X .

For an object B in $\text{Lie}(\mathbf{C})$, under these natural isomorphisms, an action θ of B on X , a morphism $\bar{\theta}: B \rightarrow L(E(X))$ in $\text{Lie}(\mathbf{C})$, a morphism $\bar{\phi}: U(B) \rightarrow E(X)$ in $\text{Mon}(\mathbf{C})$, and a monoid action ϕ of $U(B)$ on X are related when: $\theta = \text{ev}(\bar{\theta} \otimes \text{id})$, $\bar{\theta} = \bar{\phi} \nu$, $\phi = \text{ev}(\bar{\phi} \otimes \text{id})$, and (hence) $\theta = \phi(\nu \otimes \text{id})$, where ν is the unit of the adjunction $U \dashv L$.

PROOF. The natural isomorphism between the functors in (3) and (4) is standard, while the natural isomorphism between the functors in (2) and (3) is obtained from the adjunction $U \dashv L$. Given a morphism $\theta: B \otimes X \rightarrow X$, let $\bar{\theta}: B \rightarrow X^X$ be the unique morphism such that $\text{ev}(\bar{\theta} \otimes \text{id}) = \theta$. One easily observes that if the map $\bar{\theta}: B \rightarrow L(E(X))$ is a morphism in $\text{Lie}(\mathbf{C})$, then the diagram 1 below is commutative, making the whole diagram

$$\begin{array}{ccccc}
 (B \otimes B) \otimes X & \xrightarrow{(\text{id} - \sigma) \otimes \text{id}} & (B \otimes B) \otimes X & \xrightarrow{\alpha^{-1}} & B \otimes (B \otimes X) \\
 \downarrow b \otimes \text{id} & \searrow (\bar{\theta} \otimes \bar{\theta}) \otimes \text{id} & \downarrow (\text{id} - \sigma) \otimes \text{id} & \downarrow \bar{\theta} \otimes (\bar{\theta} \otimes \text{id}) & \searrow \text{id} \otimes \theta \\
 & (X^X \otimes X^X) \otimes X & (X^X \otimes X^X) \otimes X & \xrightarrow{\alpha^{-1}} & X^X \otimes (X^X \otimes X) & \rightarrow & B \otimes X \\
 & \downarrow m \otimes \text{id} & \downarrow m \otimes \text{id} & \downarrow \text{id} \otimes \text{ev} & \downarrow \text{id} \otimes \text{ev} & & \\
 B \otimes X & \xrightarrow{\bar{\theta} \otimes \text{id}} & X^X \otimes X & \xrightarrow{\text{ev}} & X & \xleftarrow{\theta} & B \otimes X \\
 & \searrow \theta & \downarrow \text{ev} & \downarrow \text{ev} & \downarrow \text{ev} & & \\
 & & X & & X & &
 \end{array}$$

commute and hence θ is an action of B on X . The converse follows immediately from the universal property of ev . ■

Let $\mathbf{2}$ denote the category with two objects 0 and 1 and one non-identity morphism $0 \rightarrow 1$. Recall that the functor category $\mathbf{C}^{\mathbf{2}}$ can be made into a symmetric monoidal category $\mathbf{C}^{\mathbf{2}}$ with monoidal structure defined componentwise. Recall also, that when \mathbf{C} has pullbacks the symmetric monoidal category $\mathbf{C}^{\mathbf{2}}$ is monoidal closed. Since the functor $L: \text{Lie}(\mathbf{C}^{\mathbf{2}}) \rightarrow \text{Mon}(\mathbf{C}^{\mathbf{2}})$ is, up to isomorphism, the same as the functor $L^{\mathbf{2}}: \text{Lie}(\mathbf{C})^{\mathbf{2}} \rightarrow \text{Mon}(\mathbf{C})^{\mathbf{2}}$, it follows from the previous proposition applied to $\mathbf{C}^{\mathbf{2}}$ that for a morphism $g: B \rightarrow B'$ in $\text{Lie}(\mathbf{C})$, a morphism $f: X \rightarrow X'$ in \mathbf{C} , and monoid actions ϕ and ϕ' of $U(B)$ on X and $U(B')$ on X' respectively, the outer arrows of the diagram

$$\begin{array}{ccccc} B \otimes X & \xrightarrow{\nu \otimes \text{id}} & U(B) \otimes X & \xrightarrow{\phi} & X \\ g \otimes f \downarrow & & \downarrow U(g) \otimes f & & \downarrow f \\ B' \otimes X' & \xrightarrow{\nu \otimes \text{id}} & U(B') \otimes X' & \xrightarrow{\phi'} & X' \end{array}$$

commute if and only if the right hand square commutes. In particular, writing $\text{LieAct}(B, \mathbf{C})$ and $\text{MonAct}(A, \mathbf{C})$ for the categories of B actions and A actions respectively (for some objects B in $\text{Lie}(\mathbf{C})$ and A in $\text{Mon}(\mathbf{C})$), this implies:

4.2. PROPOSITION. *For an object B in $\text{Lie}(\mathbf{C})$ the assignment $(X, \phi) \mapsto (X, \phi(\nu \otimes \text{id}))$ is the object map of an isomorphism*

$$\text{MonAct}(U(B), \mathbf{C}) \rightarrow \text{LieAct}(B, \mathbf{C})$$

which is identity on morphisms.

Now suppose that A is a bimonoid with comultiplication $d: A \rightarrow A \otimes A$. If ϕ and ϕ' are actions of the underlying monoid of A on X and Y , respectively, then $\phi \otimes^d \phi': A \otimes (X \otimes Y) \rightarrow (X \otimes Y)$ defined by $\phi \otimes^d \phi' = (\phi \otimes \phi')i(d \otimes \text{id})$ is a monoid action. To see why, note that the monoidal functor $(\otimes, \lambda, i): \mathbf{C} \times \mathbf{C} \rightarrow \mathbf{C}$ sends the action (ϕ, ϕ') of the monoid (A, A) on (X, Y) to the action $(\phi \otimes \phi')i$ of the monoid $A \otimes A$ on $X \otimes Y$. However, since $d: A \rightarrow A \otimes A$ is a monoid morphism it follows that $\phi \otimes^d \phi' = (\phi \otimes \phi')i(d \otimes \text{id})$ is an action of A on $X \otimes Y$.

Noting that the counit of the bimonoid A determines a monoid action of A on I one easily establishes that the symmetric monoidal structure of \mathbf{C} lifts to $\text{MonAct}(A, \mathbf{C})$ and hence the forgetful functor $G_A: \text{MonAct}(A, \mathbf{C}) \rightarrow \mathbf{C}$ is strict monoidal. Let us write $\underline{\text{MonAct}}(A, \mathbf{C})$ for this monoidal category. It is well known that when \mathbf{C} is monoidal closed G_A has a right adjoint. The object map of this right adjoint assigns to each object X the object X^A with action $\phi: A \otimes X^A \rightarrow X^A$, the unique morphism making the diagram

$$\begin{array}{ccc} (A \otimes X^A) \otimes A & \xrightarrow{\phi \otimes \text{id}} & X^A \otimes A \\ (\text{id} \otimes \sigma)\alpha(\sigma \otimes \text{id}) \downarrow & & \downarrow \text{ev} \\ X^A \otimes (A \otimes A) & \xrightarrow{\text{id} \otimes m} & X^A \otimes A \xrightarrow{\text{ev}} X \end{array}$$

commute. Therefore, since the categories \mathbb{C} and $\mathbf{MonAct}(A, \mathbf{C})$ are additive and so is the functor G_A , by Proposition 3.1, it follows that:

4.3. PROPOSITION. *The induced forgetful functor $\mathbf{Lie}(G_A): \mathbf{Lie}(\mathbf{MonAct}(A, \mathbf{C})) \rightarrow \mathbf{Lie}(\mathbf{C})$ has a right adjoint.*

Suppose B is an object in $\mathbf{Lie}(\mathbf{C})$. Let us write $\mathbf{LieAct}(B, \mathbf{C})$ for the monoidal category obtained by translating the monoidal structure from $\mathbf{MonAct}(U(B), \mathbf{C})$ via the isomorphism given in Proposition 4.2. Trivially:

4.4. PROPOSITION. *For an object B in $\mathbf{Lie}(\mathbf{C})$ the assignment given by $((X, \phi), b) \mapsto ((X, \phi(\nu \otimes \text{id})), b)$ is the object map of an isomorphism*

$$\mathbf{Lie}(\mathbf{MonAct}(U(B), \mathbf{C})) \rightarrow \mathbf{Lie}(\mathbf{LieAct}(B, \mathbf{C}))$$

which is the identity on morphisms.

Let us calculate explicitly what the tensor product in $\mathbf{LieAct}(B, \mathbf{C})$ is. Suppose that (X, θ) and (X', θ') are objects in $\mathbf{LieAct}(B, \mathbf{C})$. Letting (X, ϕ) and (X', ϕ') be the corresponding objects in $\mathbf{MonAct}(U(B), \mathbf{C})$ we see that the diagram

$$\begin{array}{ccccc}
 & & U(B)^{\otimes 2} \otimes (X \otimes X') & & \\
 & & \begin{array}{c} \downarrow i \\ (U(B) \otimes X) \otimes (U(B) \otimes X') \\ \downarrow \phi \otimes \phi' \\ X \otimes X' \end{array} & & \\
 & \begin{array}{c} \nearrow (\nu \otimes u) \otimes \text{id} \\ (B \otimes I) \otimes (X \otimes X') \\ \downarrow i \\ (B \otimes X) \otimes (I \otimes X') \\ \downarrow \text{id} \otimes \lambda \\ (B \otimes X) \otimes X' \end{array} & & \begin{array}{c} \nwarrow ((u \otimes \nu) \otimes \text{id}) \\ (I \otimes B) \otimes (X \otimes X') \\ \downarrow i \\ (I \otimes X) \otimes (B \otimes X') \\ \downarrow \lambda \otimes \text{id} \\ X \otimes (B \otimes X') \end{array} & & \\
 & \begin{array}{c} \nearrow \rho^{-1} \otimes \text{id} \\ B \otimes (X \otimes X') \\ \downarrow \alpha \\ (B \otimes X) \otimes X' \end{array} & & \begin{array}{c} \nwarrow \lambda^{-1} \otimes \text{id} \\ B \otimes (X \otimes X') \\ \downarrow \sigma \alpha(\text{id} \otimes \sigma) \\ X \otimes (B \otimes X') \end{array} & & \\
 & & X \otimes X' & & \\
 & \begin{array}{c} \nearrow \theta \otimes \text{id} \\ (B \otimes X) \otimes X' \\ \downarrow \theta \otimes \text{id} \\ X \otimes X' \end{array} & & \begin{array}{c} \nwarrow \text{id} \otimes \theta' \\ X \otimes (B \otimes X') \\ \downarrow \text{id} \otimes \theta' \\ X \otimes X' \end{array} & &
 \end{array}$$

commutes. This implies that

$$\begin{aligned}
 (\phi \otimes^d \phi')(\nu \otimes \text{id}) &= (\phi \otimes \phi')i(d\nu \otimes \text{id}) \\
 &= (\phi \otimes \phi')i(\delta\nu \otimes \text{id}) \\
 &= (\phi \otimes \phi')i((u \otimes \nu)\lambda^{-1} + (\nu \otimes u)\rho^{-1}) \otimes \text{id} \\
 &= (\theta \otimes \text{id})\alpha + (\text{id} \otimes \theta')\sigma\alpha(\text{id} \otimes \sigma)
 \end{aligned}$$

and hence the translated tensor product of $\mathbf{LieAct}(B, \mathbf{C})$ is defined by

$$(X, \theta) \otimes (X', \theta') = (X \otimes X', \theta * \theta')$$

where $\theta * \theta' = (\theta \otimes \text{id})\alpha + (\text{id} \otimes \theta')\sigma\alpha(\text{id} \otimes \sigma)$. One easily observes that this means that an object in $\text{Lie}(\underline{\text{LieAct}}(B, \mathbf{C}))$ can be identified with a pair $((X, b), \theta)$ where (X, b) is an object in $\text{Lie}(\mathbf{C})$, and θ is an action of B on X making the diagram

$$\begin{array}{ccc}
 B \otimes (X \otimes X) & \xrightarrow{\text{id} \otimes b} & B \otimes X \\
 \theta * \theta \downarrow & & \downarrow \theta \\
 X \otimes X & \xrightarrow{b} & X
 \end{array} \tag{4}$$

commute. Writing $\overline{G}_B: \underline{\text{LieAct}}(B, \mathbf{C}) \rightarrow \mathbf{C}$ for the forgetful functor we have:

4.5. PROPOSITION. *For an object B in $\text{Lie}(\mathbf{C})$ the map assigning to each object (A, p, s) in $\text{Pt}_B(\text{Lie}(\mathbf{C}))$ the pair (X, θ) where X is the domain of $k: X \rightarrow A$ the kernel of p and $\theta: B \otimes X \rightarrow X$ is the unique morphism in \mathbf{C} with $k\theta = b_A(s \otimes k)$, is the object map of a functor $W: \text{Pt}_B(\text{Lie}(\mathbf{C})) \rightarrow \text{Lie}(\underline{\text{LieAct}}(B, \mathbf{C}))$ which is part of an equivalence of categories making the diagram*

$$\begin{array}{ccc}
 \text{Pt}_B(\text{Lie}(\mathbf{C})) & \xrightarrow{W} & \text{Lie}(\underline{\text{LieAct}}(B, \mathbf{C})) \\
 \text{Ker}_B \searrow & & \swarrow \text{Lie}(\overline{G}_B) \\
 & \text{Lie}(\mathbf{C}) &
 \end{array} \tag{5}$$

commute.

PROOF. Let B be an object in $\text{Lie}(\mathbf{C})$, for an object (A, p, s) in $\text{Pt}_B(\text{Lie}(\mathbf{C}))$, defining $W(A, p, s) = (X, \theta)$ where X and θ are defined as above. One can check that the commutativity of diagrams (3) and (4) follow from commutativity of diagrams (1) and coherence.

If $f: (A, p, s) \rightarrow (A', p', s')$ is a morphism in $\text{Pt}_B(\text{Lie}(\mathbf{C}))$, then an easy calculation shows that the induced map between the kernels of p and p' lifts to a morphism $W(A, p, s) \rightarrow W(A', p', s')$ producing a functor $W: \text{Pt}_B(\text{Lie}(\mathbf{C})) \rightarrow \text{Lie}(\underline{\text{LieAct}}(B, \mathbf{C}))$ making diagram (5) commute. On the other hand, given (X, θ) in $\text{Lie}(\underline{\text{LieAct}}(B, \mathbf{C}))$ letting $b: (B \oplus X) \otimes (B \oplus X) \rightarrow B \oplus X$ be the unique morphism with $b(\iota_1 \otimes \iota_1) = \iota_1 b_B$, $b(\iota_1 \otimes \iota_2) = \theta$, $b(\iota_2 \otimes \iota_1) = -\theta\sigma$ and $b(\iota_2 \otimes \iota_2) = \iota_2 b_X$ makes $((B \oplus X, b), \pi_1, \iota_1)$ into an object of $\text{Pt}_B(\text{Lie}(\mathbf{C}))$ such that $W((B \oplus X, b), \pi_2, \iota_2) = (X, \theta)$. Therefore, noting that W is faithful, to show that it is part of an equivalence of categories we need only show it is full.

To that end suppose that (A, p, s) and (A', p', s') are objects in $\text{Pt}_B(\text{Lie}(\mathbf{C}))$, $k: X \rightarrow A$ and $k': X' \rightarrow A'$ are the kernels of p and p' , respectively, $\theta: B \otimes X \rightarrow X$ and $\theta': B \otimes X' \rightarrow X'$ are morphisms such that $W(A, p, s) = (X, \theta)$ and $W(A', p', s') = (X', \theta')$, and f is a morphism from $W(A, p, s)$ to $W(A', p', s')$. Letting $l: A \rightarrow X'$ be the unique morphism in \mathbf{C} such that $kl = \text{id} - sp$, using the fact that k and s are jointly epimorphic, one can easily check that if $g = k'fl + s'p$ then g is a morphism from (A, p, s) to (A', p', s') such that $W(g) = f$. ■

Combining the previous proposition with Proposition 4.4 we obtain:

4.6. PROPOSITION. *For each B in $\text{Lie}(\mathbf{C})$ the kernel functor $\text{Ker}_B: \text{Pt}_B(\text{Lie}(\mathbf{C})) \rightarrow \mathbf{C}$ factors through the forgetful functor $\text{Lie}(G_{U(B)}): \text{Lie}(\text{MonAct}(U(B), \mathbf{C})) \rightarrow \text{Lie}(\mathbf{C})$ via an equivalence of categories.*

We are now ready to prove our main theorem:

4.7. THEOREM. *For a symmetric monoidal closed category $\mathbf{C} = (\mathbf{C}, \otimes, I, \alpha, \lambda, \rho, \sigma)$ such that \mathbf{C} is cocomplete and additive, the category $\text{Lie}(\mathbf{C})$ is (LACC).*

PROOF. Combining the previous proposition and Proposition 4.3 we see that for each B in $\text{Lie}(\mathbf{C})$ the functor $\text{Ker}_B: \text{Pt}_B(\text{Lie}(\mathbf{C})) \rightarrow \text{Lie}(\mathbf{C})$ has a right adjoint. The claim now follows by [19, Theorem 5.1]. ■

5. Examples

5.1. EXAMPLE. *Let R be a commutative ring and consider the category of chain complexes of R -modules. If (V, d) and (V', d') are chain complexes, the tensor product of (V, d) and (V', d') is the chain complex $(V \otimes V', \delta)$ where*

$$(V \otimes V')_i = \bigoplus_{j+k=i} V_j \otimes V'_k,$$

and δ is defined for all $v \in V_j$ and $v' \in V'_k$ by

$$\delta(v \otimes v') = d(v) \otimes v' + (-1)^j v \otimes d'(v').$$

The symmetry isomorphism $\sigma: V \otimes V' \rightarrow V' \otimes V$ is defined for all $v \in V_j$ and $v' \in V'_k$ by $\sigma(v \otimes v') = (-1)^{ij} v' \otimes v$. With this tensor product, symmetry isomorphism and the remaining structure defined canonically, it forms a symmetric monoidal closed category whose underlying category is cocomplete and abelian.

Its category of internal Lie algebras is equivalent to the category whose objects are differential graded Lie algebras, i.e., a \mathbb{Z} -graded R -module V , with a linear map $d: V \rightarrow V$ of degree -1 such that $d \circ d = 0$, and a bilinear map $[-, -]: V \otimes V \rightarrow V$ of degree zero satisfying: for all homogeneous x, y, z in V

1. $[x, y] = -(-1)^{|x||y|}[y, x]$,
2. $(-1)^{|x||z|}[x, [y, z]] + (-1)^{|y||x|}[y, [z, x]] + (-1)^{|z||y|}[z, [x, y]] = 0$,
3. $d([x, y]) = [d(x), y] + (-1)^{|x|}[x, d(y)]$

where $|v|$ denotes the degree of v . By Theorem 4.7 it follows that the category of differential graded Lie algebras is (LACC).

5.2. EXAMPLE. *If in the previous example we consider the full subcategory formed by chain complexes where $d = 0$, internal Lie algebras are essentially \mathbb{Z} -graded Lie algebras. Applying Theorem 4.7 again we know that the category of \mathbb{Z} -graded Lie algebras is (LACC).*

In a similar way, the category of Lie superalgebras over a commutative ring can be seen as the category of internal Lie algebras in the symmetric monoidal closed category of super R -modules and hence it is (LACC).

5.3. EXAMPLE. *Let us consider now the category whose objects are linear maps $f: V \rightarrow W$ and morphisms are homomorphisms of R -modules making the diagram commutative*

$$\begin{array}{ccc} V & \longrightarrow & V' \\ f \downarrow & & \downarrow f' \\ W & \longrightarrow & W' \end{array}$$

There is a tensor product, called the infinitesimal tensor product defined as

$$\left(\begin{array}{c} V \\ \downarrow f \\ W \end{array} \right) \otimes \left(\begin{array}{c} V' \\ \downarrow f' \\ W' \end{array} \right) = \left(\begin{array}{c} (V \otimes W') \oplus (W \otimes V') \\ \downarrow [f \otimes \text{id}, \text{id} \otimes f'] \\ W \otimes W' \end{array} \right)$$

This monoidal category was introduced in [23] as the tensor category of linear maps, and in more recent papers was renamed to the Loday-Pirashvili category [9, 10, 26], denoted by \mathcal{LP} . An internal Lie algebra in this category is a linear map $f: M \rightarrow \mathfrak{g}$, where \mathfrak{g} is a Lie algebra, M is a right \mathfrak{g} -module and f preserves the \mathfrak{g} -action. Once more, as a consequence of Theorem 4.7, the category of Lie algebras in \mathcal{LP} is (LACC). Furthermore, the category of Leibniz algebras can be found as a full reflective subcategory of internal Lie algebras in \mathcal{LP} , but it is known that the category of Leibniz algebras is not (LACC) [12].

5.4. EXAMPLE. *Whereas the categories of Hom-Lie algebras and multiplicative Hom-Lie algebras are known to be semi-abelian categories satisfying very few categorical-algebraic properties (see [7]), it was shown in [6, 14] that regular Hom-Lie algebras (Hom-Lie algebras whose twist is an isomorphism preserving the bracket) can be found as internal Lie algebras in a certain symmetric monoidal category, and therefore, again by Theorem 4.7, it is (LACC).*

Acknowledgements

The first author is grateful to Stellenbosch University for their kind hospitality during his stay there. The authors also want to thank the referee for the very useful comments that helped to improve this paper.

References

- [1] F. Borceux, G. Janelidze, and G. M. Kelly, *On the representability of actions in a semi-abelian category*, Theory Appl. Categ. **14** (2005), no. 11, 244–286.
- [2] D. Bourn, *Normalization equivalence, kernel equivalence and affine categories*, Category Theory, Proceedings Como 1990 (A. Carboni, M. C. Pedicchio, and G. Rosolini, eds.), Lecture Notes in Math., vol. 1488, Springer, 1991, pp. 43–62.
- [3] D. Bourn, *Normal functors and strong protomodularity*, Theory Appl. Categ. **7** (2000), no. 9, 206–218.
- [4] D. Bourn, *The cohomological comparison arising from the associated abelian object*, preprint [arXiv:1001.0905](https://arxiv.org/abs/1001.0905), 2010.
- [5] D. Bourn and J. R. A. Gray, *Aspects of algebraic exponentiation*, Bull. Belg. Math. Soc. Simon Stevin **19** (2012), 823–846.
- [6] S. Caenepeel and I. Goyvaerts, *Monoidal Hom-Hopf algebras*, Comm. Algebra **39** (2011), no. 6, 2216–2240.
- [7] J. M. Casas and X. García-Martínez, *Abelian extensions and crossed modules of Hom-Lie algebras*, J. Pure Appl. Algebra **224** (2020), no. 3, 987–1008.
- [8] A. S. Cigoli, J. R. A. Gray, and T. Van der Linden, *On the normality of Higgins commutators*, J. Pure Appl. Algebra **219** (2015), 897–912.
- [9] R. Fernández-Casado, X. García-Martínez, and M. Ladra, *A natural extension of the universal enveloping algebra functor to crossed modules of Leibniz algebras*, Appl. Categ. Structures **25** (2017), no. 6, 1059–1076.
- [10] X. García-Martínez and M. Ladra, *Universal central extensions of Leibniz superalgebras over superdialgebras*, Mediterr. J. Math. **14** (2017), 14: 63.
- [11] X. García-Martínez, M. Tsishyn, T. Van der Linden, and C. Vienne, *Algebras with representable representations*, To appear in Proc. Edinburgh Math. Soc., 2021.
- [12] X. García-Martínez and T. Van der Linden, *A characterisation of Lie algebras amongst anti-commutative algebras*, J. Pure Appl. Algebra **223** (2019), no. 11, 4857–4870.
- [13] X. García-Martínez and T. Van der Linden, *A characterisation of Lie algebras via algebraic exponentiation*, Adv. Math. **341** (2019), 92–117.
- [14] I. Goyvaerts and J. Verduyck, *A note on the categorification of Lie algebras*, Lie theory and its applications in physics, Springer Proc. Math. Stat., vol. 36, Springer, Tokyo, 2013, pp. 541–550.

- [15] I. Goyvaerts and J. Vercauteren, *Lie monads and dualities*, J. Algebra **414** (2014), 120–158.
- [16] I. Goyvaerts and J. Vercauteren, *On the duality of generalized Lie and Hopf algebras*, Adv. Math. **258** (2014), 154–190.
- [17] J. R. A. Gray, *Algebraic exponentiation in general categories*, Ph.D. thesis, University of Cape Town, 2010.
- [18] J. R. A. Gray, *Algebraic exponentiation for categories of Lie algebras*, J. Pure Appl. Algebra **216** (2012), 1964–1967.
- [19] J. R. A. Gray, *Algebraic exponentiation in general categories*, Appl. Categ. Structures **20** (2012), 543–567.
- [20] G. Janelidze, L. Márki, and W. Tholen, *Semi-abelian categories*, J. Pure Appl. Algebra **168** (2002), no. 2–3, 367–386.
- [21] G. M. Kelly, *Doctrinal adjunction*, Category Seminar (Proc. Sem., Sydney, 1972/1973), 1974, pp. 257–280. Lecture Notes in Math., Vol. 420.
- [22] Stephen Lack, *Note on the construction of free monoids*, Applied Categorical Structures **18** (2010), no. 1, 17–29.
- [23] J. L. Loday and T. Pirashvili, *The tensor category of linear maps and Leibniz algebras*, Georgian Math. J. **5** (1998), no. 3, 263–276.
- [24] S. Mac Lane, *Categories for the working mathematician*, second ed., Grad. Texts in Math., vol. 5, Springer, 1998.
- [25] N. Martins-Ferreira and T. Van der Linden, *A note on the “Smith is Huq” condition*, Appl. Categ. Structures **20** (2012), no. 2, 175–187.
- [26] A. Rovi, *Lie algebroids in the Loday-Pirashvili category*, SIGMA Symmetry Integrability Geom. Methods Appl. **11** (2015), Paper 079, 16.

*Departamento de Matemáticas,
Esc. Sup. de Enx. Informática
Campus de Ourense, Universidade de Vigo
E-32004, Ourense, Spain
and
Faculty of Engineering
Vrije Universiteit Brussel
Pleinlaan 2
B-1050 Brussel Belgium*

*Mathematics Division
Department of Mathematical Sciences
Stellenbosch University
Private Bag X1
7602 Matieland
South Africa*

Email: `xabier.garcia.martinez@uvigo.gal`
`jamesgray@sun.ac.za`

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS \LaTeX 2 ϵ is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Geoff Cruttwell, Mount Allison University: gcruttwell@mta.ca

TEXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT TEX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Gabriella Böhm, Wigner Research Centre for Physics: bohm.gabriella@wigner.mta.hu

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Dirk Hofmann, Universidade de Aveiro: dirk@ua.pt

Pieter Hofstra, Université d' Ottawa: phofstra@uottawa.ca

Anders Kock, University of Aarhus: kock@math.au.dk

Joachim Kock, Universitat Autònoma de Barcelona: kock@mat.uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Ieke Moerdijk, Utrecht University: i.moerdijk@uu.nl

Susan Niefield, Union College: niefiels@union.edu

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: ross.street@mq.edu.au

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be