

# A GELFAND DUALITY FOR CONTINUOUS LATTICES

RUIYUAN CHEN

ABSTRACT. We prove that the category of continuous lattices and meet- and directed join-preserving maps is dually equivalent, via the hom functor to  $[0, 1]$ , to the category of complete Archimedean meet-semilattices equipped with a finite meet-preserving action of the monoid of continuous monotone maps of  $[0, 1]$  fixing 1. We also prove an analogous duality for completely distributive lattices. Moreover, we prove that these are essentially the only well-behaved “sound classes of joins  $\Phi$ , dual to a class of meets” for which “ $\Phi$ -continuous lattice” and “ $\Phi$ -algebraic lattice” are different notions, thus for which a 2-valued duality does not suffice.

## 1. Introduction

The classical Gelfand duality asserts that a compact Hausdorff space  $X$  may be recovered from its ring of continuous functions  $C(X)$ , and moreover such rings are up to isomorphism precisely the commutative  $C^*$ -algebras. From a categorical perspective,  $C(X)$  is best regarded as having “underlying set” given by its (positive) unit ball, i.e., consisting of continuous  $\mathbb{I} := [0, 1]$ -valued functions, so that Gelfand duality falls under the umbrella of Stone-type dualities induced by two “commuting” structures on  $\mathbb{I}$ ; see [Joh82, VI §4]. Namely,  $\mathbb{I}$  is equipped with its usual compact Hausdorff topology, and also with all operations  $\mathbb{I}^{\kappa} \rightarrow \mathbb{I}$  “commuting” with the topology, i.e., which are continuous. Thus, for another object in either category, the hom functor into  $\mathbb{I}$  yields a dual in the other category, and this gives a dual adjunction, which Gelfand duality asserts is an equivalence. An explicit axiomatization of the dual operations on the  $\mathbb{I}$ -valued  $C(X)$  was recently given in [MR17]; see there for a detailed history of  $\mathbb{I}$ -valued Gelfand duality. In [HNN18], [Abb19],  $\mathbb{I}$ -valued Gelfand duality was further extended to compact partially ordered spaces (*a la* Nachbin).

In this note, we prove analogous Gelfand-type dualities for compact pospaces equipped with lattice operations. Recall that a **continuous lattice** is a compact topological meet-semilattice obeying a “local convexity under meets” condition, that each point has a neighborhood basis of subsemilattices. Equivalently, they can be defined purely order-theoretically as posets with arbitrary meets distributing over directed joins. An analog of Urysohn’s lemma, sometimes known as the Urysohn–Lawson lemma, states that every continuous lattice  $X$  admits enough morphisms to  $\mathbb{I}$ , i.e., the canonical evaluation map  $X \rightarrow \mathbb{I}^{\text{Hom}(X, \mathbb{I})}$  is an embedding; see [G<sup>+</sup>03, IV-3.3], [Joh82, VII 3.2]. It is thus natural

---

Received by the editors 2023-01-09 and, in final form, 2024-01-11.

Transmitted by Dirk Hofmann. Published on 2024-01-16.

2020 Mathematics Subject Classification: 06B35, 06D10, 18F70, 18A35.

Key words and phrases: continuous lattice, completely distributive lattice, duality, free cocompletion.

© Ruiyuan Chen, 2024. Permission to copy for private use granted.

to ask whether, by equipping  $\text{Hom}(X, \mathbb{I})$  with suitable structure commuting with the continuous lattice structure on  $\mathbb{I}$ , we may recover  $X$  as the double dual.

Let  $\widehat{\mathbb{U}}$  denote the monoid of continuous monotone maps  $\mathbb{I} \rightarrow \mathbb{I}$  fixing 1, i.e., all unary operations on  $\mathbb{I}$  commuting with the continuous lattice structure. Note that finite meets do as well. By a  $\widehat{\mathbb{U}}$ -**module**, we mean a unital meet-semilattice equipped with an action of  $\widehat{\mathbb{U}}$  preserving finite meets in both variables. In every  $\widehat{\mathbb{U}}$ -module  $A$ , we have a canonical pseudoquasimetric

$$\rho(a, b) := \bigwedge \{r \in \mathbb{I} \mid a \leq b \dot{+} r\}$$

where  $b \dot{+} r$  denotes the result of the action on  $b$  of the truncated addition  $(-) \dot{+} r \in \widehat{\mathbb{U}}$ . We say  $A$  is **Archimedean** if  $\rho(a, b) = 0 \implies a \leq b$ , and **complete** if  $A$  is Archimedean and complete with respect to the induced metric  $d(a, b) := \rho(a, b) \vee \rho(b, a)$ . We prove

1.1. **THEOREM.** [Corollary 5.9] *Hom into  $\mathbb{I}$  yields a dual equivalence of categories between continuous lattices and complete  $\widehat{\mathbb{U}}$ -modules.*

There is a generalization of continuous lattice theory, with the role of directed joins replaced by an arbitrary “class of joins  $\Phi$ ” obeying suitable axioms; see [WWT78], [BE83], [Xu95], as well as [AK88], [ABLR02], [KS05] for a further extension in enriched category theory. Other than  $\Phi =$  “directed joins”, the most well-known case is  $\Phi =$  “all joins”, for which  $\Phi$ -continuous lattices are completely distributive lattices. As for continuous lattices, there is a Urysohn-type lemma, stating that all completely distributive lattices admit enough morphisms to  $\mathbb{I}$ ; see [G<sup>+</sup>03, IV-3.31–32], [Joh82, 1.10–14]. We likewise boost this to a Gelfand-type duality as follows.

Let  $\mathbb{U} \subseteq \widehat{\mathbb{U}}$  denote the monoid of complete lattice morphisms, i.e., monotone surjections. A  $\mathbb{U}$ -**poset** is a poset with a monotone action of  $\mathbb{U}$ . There is a canonical way of defining a pseudoquasimetric on a  $\mathbb{U}$ -poset, agreeing with the above definition in  $\widehat{\mathbb{U}}$ -modules; see Definition 4.2. A  $\mathbb{U}$ -poset  $A$  is **stackable** if, intuitively speaking, an element  $a \in A$  may be specified via its “restrictions to sublevel and superlevel sets  $a^{-1}([0, r]), a^{-1}([r, 1])$ ” for any  $0 < r < 1$ ; see Definition 4.12.

1.2. **THEOREM.** [Corollary 5.5] *Hom into  $\mathbb{I}$  yields a dual equivalence of categories between completely distributive lattices and complete stackable  $\mathbb{U}$ -posets.*

In fact, we prove a single result underlying Theorems 1.1 and 1.2, for a “class of joins  $\Phi$  dual to a class of meets  $\Psi^{\text{op}}$ ”, more precisely for a *sound* class of joins in the sense of [ABLR02], [KS05]; see Section 3. This general result, Theorem 5.2, says that  $\Phi$ -continuous lattices are dual to complete stackable  $\mathbb{U}$ - $\Psi^{\text{op}}$ -inflattices, *provided that not all  $\Phi$ -continuous lattices are  $\Phi$ -algebraic*, i.e., already admit enough morphisms into 2. This is a reasonable restriction, since for these other  $\Phi$ , we instead have a simple 2-valued duality generalizing the classical Hofmann–Mislove–Stralka duality [HMS74] between algebraic lattices and meet-semilattices (see Corollary 3.7).

Part of the reason we work with general  $\Phi$  is to hint at the possibility of generalizing to quantale-enriched posets, or even to enriched categories, which we plan to pursue in

future work. However, in the original context of mere posets, it turns out that essentially the only  $\Phi$  are the classical ones:

1.3. THEOREM. [Theorem 3.9] *There are precisely 4 sound classes of joins  $\Phi$  for which not every  $\Phi$ -continuous lattice is  $\Phi$ -algebraic: “directed joins”, “all joins”, and the minor variations including/excluding empty joins.*

ACKNOWLEDGMENTS. I would like to thank the anonymous referee for numerous helpful comments and suggestions that improved the presentation of the paper. Research partially supported by NSF grant DMS-2224709.

## 2. $\Phi$ -continuous lattices

We assume familiarity with basic category theory. For a category  $\mathbf{C}$ ,  $\mathbf{C}(X, Y)$  will denote the hom-set of morphisms from  $X$  to  $Y$ , while  $\mathbf{C}^{\text{op}}$  will denote the opposite category; this includes opposite posets. We let  $\mathbf{Pos}$  denote the category of posets,  $\mathbf{Sup}$  denote the category of suplattices (i.e., complete lattices with join-preserving maps as morphisms),  $\mathbf{Inf}$  denote the category of inflattices, and  $\mathbf{CLat} = \mathbf{Sup} \cap \mathbf{Inf}$  denote the category of complete lattices. These are all locally ordered categories: each hom-set is partially ordered pointwise, and composition is monotone on both sides. For  $f : X \rightarrow Y \in \mathbf{Pos}$  left adjoint to  $g : Y \rightarrow X$ , we will write  $f = g^+$  and  $g = f^\times$ . We will frequently use the “mate calculus”: for monotone  $h, k$ , we have  $h \circ g \leq k \iff h \leq k \circ f$ .

For a poset  $X$ , we let  $\mathcal{L}(X)$  denote the poset of lower sets  $\phi \subseteq X$ , ordered via  $\subseteq$ . Then  $\mathcal{L} : \mathbf{Pos} \rightarrow \mathbf{Pos}$  is the free suplattice monad, where the monad structure consists of:

- unit  $\downarrow = \downarrow_X : X \rightarrow \mathcal{L}(X)$ , where  $\downarrow x = \{y \mid y \leq x\}$  is the principal ideal below  $x$ ;
- multiplication  $\bigcup : \mathcal{L}(\mathcal{L}(X)) \rightarrow \mathcal{L}(X)$ ;
- $f : X \rightarrow Y \in \mathbf{Pos}$  inducing  $f_* = \mathcal{L}(f) : \mathcal{L}(X) \rightarrow \mathcal{L}(Y) \in \mathbf{Sup}$ , where  $f_*(\phi) = \bigcup_{x \in \phi} \downarrow f(x)$ .

We now review the theory of “relative” suplattices for a “class of joins”  $\Phi$ . This is a special case of the theory of “classes of colimits” in enriched category theory [AK88], [ABLR02], [KS05], and has also been well-studied in the order theory literature as “ $Z$ -completeness” [WWT78], [BE83]. We will use notation and terminology based on that from enriched categories.

2.1. DEFINITION. A **join doctrine** is a class  $\Phi$  of posets  $\phi$ , thought of as indexing posets for certain joins  $\bigvee_{x \in \phi} f(x)$  of monotone  $f : \phi \rightarrow Y$ . We require  $\Phi$  to obey the following “saturation” conditions:

- (i) The singleton poset  $\mathbf{1}$  is in  $\Phi$ .
- (ii) If  $\phi$  is a poset which is a union  $\bigcup \Psi$  of a set  $\Psi \subseteq \Phi$  of subposets  $\psi \subseteq \phi$  which are in  $\Phi$ , and also  $\Psi$  (as a poset under  $\subseteq$ ) is in  $\Phi$ , then  $\phi \in \Phi$ .

(iii) If  $f : \phi \rightarrow \psi$  is a monotone map with cofinal image, and  $\phi \in \Phi$ , then  $\psi \in \Phi$ .

(iv) If  $\phi \subseteq \psi$  is a cofinal subposet, and  $\psi \in \Phi$ , then  $\phi \in \Phi$ .

A  $\Phi$ -**join** in a poset  $X$  is a join of a subset  $\phi \subseteq X$  such that  $\phi \in \Phi$ . A  $\Phi$ -**suplattice** is a poset with all  $\Phi$ -joins; we denote the category of all such (and monotone  $\Phi$ -join-preserving maps) by  $\Phi\text{Sup}$ . A  $\Phi$ -**ideal** in a  $\Phi$ -suplattice is a lower sub- $\Phi$ -suplattice. The **free  $\Phi$ -suplattice** generated by a poset  $X$  is the subset  $\Phi(X) \subseteq \mathcal{L}(X)$  of all lower subsets of  $X$  in  $\Phi$ . Note that for a poset  $\phi$ , we have  $\phi \in \Phi \iff \phi \in \Phi(\phi)$ ; we thereby identify the class of posets  $\Phi$  with the submonad  $\Phi \subseteq \mathcal{L}$ .

## 2.2. EXAMPLE.

- The “class of directed joins” is given by the join doctrine  $\Phi :=$  all directed posets, for which a  $\Phi$ -suplattice is a directed-complete poset (DCPO), a  $\Phi$ -ideal is a Scott-closed subset, and  $\Phi(X)$  is the ideal completion of  $X$  (note: *not* “ $\Phi$ -ideal completion”).
- The “class of finite joins” is given by  $\Phi :=$  all posets with finite cofinality.
- The “class of all joins” is given by  $\Phi :=$  all posets.
- The least join doctrine, of “trivial joins”, is given by  $\Phi :=$  posets with a greatest element.

2.3. REMARK. In [AK88] and [KS05], a more general notion of “class of colimits” is considered, consisting in the posets case of an arbitrary submonad  $\Phi \subseteq \mathcal{L}$ , i.e., an assignment to each poset  $X$  of a set of lower sets  $\Phi(X) \subseteq \mathcal{L}(X)$  closed under the monad operations on  $\mathcal{L}$ .

The precise connection with our definition of “join doctrine” as a class of posets is as follows. Each join doctrine  $\Phi$  induces a free  $\Phi$ -suplattice submonad as above; this yields an order-embedding

$$\{\text{join doctrines}\} \hookrightarrow \{\text{submonads of } \mathcal{L}\},$$

whose image consists of those submonads  $\Phi \subseteq \mathcal{L}$  obeying the additional “saturation” condition

$$(*) \text{ for each order-embedding between posets } f : X \hookrightarrow Y, \text{ we have } \Phi(X) = f_*^{-1}(\Phi(Y)).$$

This condition is implied by condition (iv) in Definition 2.1 of join doctrine, and conversely, ensures that  $\{\phi \in \text{Pos} \mid \phi \in \Phi(\phi)\}$  is a join doctrine inducing the submonad  $\Phi$ .

An example of a submonad not obeying (\*) is

$$\Phi(X) := \{\phi \in \mathcal{L}(X) \mid \phi \text{ has an upper bound in } X\},$$

which yields the “class of bounded joins”. However, (\*) is automatic for the  $\Phi$  suitable for our duality purposes, which is why we use the simpler definition of “join doctrine”; see Remark 3.2.

2.4. DEFINITION. Let  $\Phi$  be a join doctrine,  $X$  be a  $\Phi$ -suplattice. We define, for  $x, y \in X$ ,

$$\begin{aligned} \downarrow &= \downarrow_X^\Phi : X \longrightarrow \mathcal{L}(X) \\ x &\longmapsto \bigcap \{ \phi \in \Phi(X) \mid x \leq \bigvee \phi \}, \\ x \ll y &:\iff x \ll^\Phi y :\iff x \in \downarrow y. \end{aligned}$$

We call  $x \in X$   **$\Phi$ -compact** ( $\Phi$ -atomic in [KS05]) if  $x \ll^\Phi x$ , i.e., whenever  $\bigvee_i y_i$  is a  $\Phi$ -join  $\geq x$ , then some  $y_i \geq x$ , i.e., the indicator function of  $\uparrow x : X \rightarrow 2$  preserves  $\Phi$ -joins. Denote these by

$$X_\Phi := \{x \in X \mid x \ll^\Phi x\}.$$

We call  $X$   **$\Phi$ -algebraic** if it is generated under  $\Phi$ -joins by  $X_\Phi \subseteq X$ . In that case, it is easy to see that in fact, for each  $x \in X$  the set  $X_\Phi \cap \downarrow x$  belongs to  $\Phi(X_\Phi)$  and has join  $x$ ; and this yields an order-isomorphism  $X \cong \Phi(X_\Phi)$ . Conversely, for any poset  $Y$ , we easily have that  $\Phi(Y)$  is  $\Phi$ -algebraic, with  $\Phi(Y)_\Phi = \{\text{principal ideals}\} \cong Y$ .

2.5. PROPOSITION. Let  $\Phi$  be a join doctrine,  $X$  be a  $\Phi$ -suplattice. The following are equivalent:

- (i) For each  $x \in X$ , there is a  $\phi \in \Phi(X)$  such that  $\phi \subseteq \downarrow x$  and  $x \leq \bigvee \phi$ , whence in fact  $\phi = \downarrow x$ .
- (ii)  $\bigvee : \Phi(X) \rightarrow X$  has a left adjoint, namely  $\downarrow$ .

If  $X$  is a complete lattice, these are further equivalent to:

- (iii)  $\bigvee : \Phi(X) \rightarrow X$  preserves meets.
- (iv) Arbitrary meets distribute over  $\Phi$ -joins: if  $\bigvee_{j \in J_i} x_{i,j}$  is a  $\Phi$ -join for each  $i \in I$ , then

$$\bigwedge_{i \in I} \bigvee_{j \in J_i} x_{i,j} = \bigvee_{(j_i)_{i \in I} \in \prod_i J_i} \bigwedge_{i \in I} x_{i,j_i}.$$

All of these hold if  $X$  is algebraic, with  $\downarrow = \downarrow_* : \Phi(X_\Phi) \rightarrow \Phi(\Phi(X_\Phi))$ , i.e.,

$$x \ll y \iff \exists z \in X_\Phi (x \leq z \leq y).$$

If (i), (ii) hold for a  $\Phi$ -suplattice  $X$ , we call  $X$   **$\Phi$ -continuous**. If furthermore  $X$  is a complete lattice, we call  $X$  a  **$\Phi$ -continuous lattice**, or a  **$\Phi$ -algebraic lattice** if  $X$  is algebraic.

PROOF. (i)  $\iff$  (ii) since it is easily seen that  $\phi$  in (i) must be  $\downarrow x$ .

(ii)  $\iff$  (iii) by the adjoint functor theorem.

(iii)  $\iff$  (iv) because the latter says  $\bigwedge_{i \in I} \bigvee \bigcup_{j \in J_i} \downarrow x_{i,j_i} = \bigvee \bigcap_{i \in I} \bigcup_{j \in J_i} \downarrow x_{i,j_i}$ . ■

2.6. PROPOSITION. *In every  $\Phi$ -suplattice,*

$$(a) \downarrow x \subseteq \downarrow x, \text{ i.e., } y \ll x \implies y \leq x.$$

$$(b) x' \leq x \ll y \leq y' \implies x' \ll y'.$$

*In a  $\Phi$ -continuous  $\Phi$ -suplattice,*

$$(c) \text{ (interpolation) } \downarrow = \bigcup \downarrow_* \downarrow, \text{ i.e., } \downarrow x = \bigcup_{y \ll x} \downarrow y, \text{ i.e.,}$$

$$z \ll x \iff \exists y (z \ll y \ll x).$$

PROOF. The first two are obvious. For interpolation: since  $X$  is an algebra of the monad  $\Phi$ , we have  $\bigvee \bigcup = \bigvee \bigvee_* : \Phi(\Phi(X)) \rightarrow X$ ; taking left adjoints yields  $\downarrow_* \downarrow = \downarrow_* \downarrow$ ; now take  $\bigcup$ . ■

A **morphism of  $\Phi$ -continuous lattices** is a meet-preserving,  $\Phi$ -join-preserving map between  $\Phi$ -continuous lattices. Let  $\Phi\text{CtsLat}$  denote the category of  $\Phi$ -continuous lattices and morphisms, and  $\Phi\text{AlgLat} \subseteq \Phi\text{CtsLat}$  denote the full subcategory of  $\Phi$ -algebraic lattices.

2.7. PROPOSITION. *Let  $f : X \rightarrow Y$  be a right adjoint between  $\Phi$ -continuous  $\Phi$ -suplattices, with left adjoint  $f^+ : Y \rightarrow X$ . Then  $f$  preserves  $\Phi$ -joins iff  $f^+$  preserves  $\ll$ . Thus*

$$\begin{aligned} \Phi\text{CtsLat}(X, Y)^{\text{op}} &\cong \ll^{\Phi}\text{Sup}(Y, X) := \{f^+ : Y \rightarrow X \mid f^+ \text{ preserves } \ll, \bigvee\} \\ &f \mapsto f^+. \end{aligned}$$

PROOF.  $f \bigvee = \bigvee f_* : \Phi(X) \rightarrow Y$  iff, taking left adjoints,  $\downarrow f^+ = (f^+)_* \downarrow : Y \rightarrow \Phi(X)$ . ■

2.8. PROPOSITION. *Let  $\Phi$  be a join doctrine. The following are equivalent:*

- (i) *For every complete lattice  $X$ ,  $\Phi(X) \subseteq \mathcal{L}(X)$  is closed under meets.*
- (ii) *For every poset  $X$ ,  $\Phi(\mathcal{L}(X)) \subseteq \mathcal{L}(\mathcal{L}(X))$  is closed under meets.*
- (iii) *For every poset  $X$ ,  $\mathcal{L}(X)$  is  $\Phi$ -continuous.*

If these conditions hold, we call  $\Phi$  a **continuous join doctrine**.

PROOF. (i)  $\implies$  (ii) is obvious.

(ii)  $\implies$  (iii) since  $\bigcup : \Phi(\mathcal{L}(X)) \rightarrow \mathcal{L}(X)$  is the composite of the inclusion  $\Phi(\mathcal{L}(X)) \hookrightarrow \mathcal{L}(\mathcal{L}(X))$  and  $\bigcup : \mathcal{L}(\mathcal{L}(X)) \rightarrow \mathcal{L}(X)$ , which both preserve meets, i.e., have left adjoints.

(iii)  $\implies$  (i) since the composite  $\mathcal{L}(X) \xrightarrow{\downarrow_{\mathcal{L}(X)}} \Phi(\mathcal{L}(X)) \xrightarrow{\bigvee_*} \Phi(X)$  yields the  $\Phi(X)$ -closure of each lower set  $\psi$ : we have  $1_{\mathcal{L}(X)} \leq \bigvee_* \downarrow_{\mathcal{L}(X)}$  because  $\bigcup \leq \bigvee_* : \Phi(\mathcal{L}(X)) \rightarrow \Phi(X) \subseteq \mathcal{L}(X)$ , while  $\bigvee_* \downarrow_{\mathcal{L}(X)}$  restricted to  $\Phi(X) \subseteq \mathcal{L}(X)$  becomes  $\bigvee_* \downarrow_* = 1_{\Phi(X)}$ . ■

The following are the two main examples of continuous join doctrines:

2.9. **EXAMPLE.** If  $\Phi$  is the “class of directed joins”, i.e., the class of all directed posets, so that  $\Phi(X)$  for  $X \in \mathbf{Pos}$  is the ideal completion of  $X$ , then  $\ll$  is the classical way-below relation, and  $\Phi$ -continuity and  $\Phi$ -algebraicity become classical continuity and algebraicity for DCPOs.

Similarly, for any infinite regular cardinal  $\kappa$ , one can consider  $\kappa$ -directed joins. But it turns out that for uncountable  $\kappa$ , continuity and algebraicity coincide; see Corollary 2.13.

2.10. **EXAMPLE.** If  $\Phi$  is the “class of all joins”, i.e., the class of all posets, so that  $\Phi(X) = \mathcal{L}(X)$ , then a  $\Phi$ -continuous lattice is a completely distributive lattice, and  $\ll$  is the “way-way-below” relation sometimes denoted  $\lll$ ; see e.g., [G<sup>+</sup>03, IV-3.31].

Minor variations are to include/exclude empty joins, which only affects  $\Phi$ -compactness of  $\perp$ .

2.11. **EXAMPLE.** [the unit interval] For any join doctrine  $\Phi$ ,  $\mathbb{I} := [0, 1]$  is a  $\Phi$ -continuous lattice. Indeed,  $\ll$  contains  $<$ , since any  $\phi \in \mathcal{L}(\mathbb{I})$  with  $r \leq \bigvee \phi$  must clearly contain  $[0, r]$ ; thus  $r = \bigvee \downarrow r$ .

We now completely characterize the  $\ll^\Phi$  relation on  $\mathbb{I}$ , by determining which  $r \in \mathbb{I}$  are  $\Phi$ -compact.

2.12. **PROPOSITION.** *Let  $\Phi$  be a join doctrine.*

- (a) *For every  $\Phi$ -suplattice  $X$ ,  $\perp \in X$  is  $\Phi$ -compact iff  $\emptyset \notin \Phi$ . In particular, this holds for  $0 \in \mathbb{I}$ .*
- (b) *If  $\omega \in \Phi$  (where  $\omega$  has the usual linear order), then no  $r > 0$  is  $\Phi$ -compact in  $\mathbb{I}$ . Otherwise:*
  - (i) *For every  $\phi \in \Phi$  and  $x_0, x_1, \dots \in \phi$ , there are  $i_0 < i_1 < \dots$  such that  $x_{i_0}, x_{i_1}, \dots$  have an upper bound in  $\phi$ . In particular, every  $x_0 \leq x_1 \leq \dots \in \phi$  has an upper bound.*
  - (ii) *Every  $\Phi$ -continuous  $\Phi$ -suplattice  $X$  which also has countable increasing joins is  $\Phi$ -algebraic, with the join of any  $x_0 \ll x_1 \ll \dots \in X$  being  $\Phi$ -compact. In particular, every  $r > 0$  is  $\Phi$ -compact in  $\mathbb{I}$ .*

**PROOF.** (a) is clear from the definition of  $\Phi$ -compact.

(b) If  $\omega \in \Phi$ , then no  $r > 0$  is  $\Phi$ -compact, since  $r$  is the join of a sequence in  $[0, r)$ . Now suppose  $\omega \notin \Phi$ . Then for  $\phi \in \Phi$  and  $x_0, x_1, \dots \in \phi$ , if no infinite subfamily has an upper bound, then we have a monotone map  $\phi \rightarrow \omega$  taking  $\phi \setminus \bigcup_n \uparrow x_n$  to 0 and each  $\uparrow x_n \setminus \bigcup_{m>n} \uparrow x_m$  to  $n + 1$ ; since  $\omega \notin \Phi$ , this map must have finite image, whence there are  $i_0 < i_1 < \dots$  with  $x_{i_0} \geq x_{i_1} \geq \dots$ , a contradiction, which proves (i). It follows that for a  $\Phi$ -continuous  $\Phi$ -suplattice  $X$  with countable increasing joins, every  $\downarrow x \in \Phi(X)$  is closed under countable increasing joins. In particular, for  $x_0 \ll x_1 \ll \dots \in X$ ,  $x := \bigvee_n x_n$  has  $x_n \ll x$  for each  $n$ , whence  $x \ll x$ . Now for any  $y \in X$  and  $x_0 \ll y$ , by interpolation

(Proposition 2.6(c)) we may find  $x_0 \ll x_1 \ll \cdots \ll y$ , whence  $x := \bigvee_n x_n$  is  $\Phi$ -compact with  $x_0 \leq x \ll y$ ; since  $y = \bigvee \downarrow y$ , it follows that  $X$  is  $\Phi$ -algebraic, proving (ii). ■

2.13. COROLLARY. *For a join doctrine  $\Phi$ , the following are equivalent:*

(i)  $\omega \notin \Phi$ .

(ii)  $\mathbb{I}$  is  $\Phi$ -algebraic.

(iii) Every  $\Phi$ -continuous lattice is  $\Phi$ -algebraic. ■

### 3. Commuting meets and joins

We are interested in recovering  $\Phi$ -continuous lattices from their dual algebras of morphisms (to  $\mathbf{2}$  or  $\mathbb{I}$ ). In order to do so, by general duality theory, the dual algebras must be equipped with all operations which commute with the  $\Phi$ -continuous lattice operations of arbitrary meets and  $\Phi$ -joins. Thus, we now review the theory of classes of commuting meets and joins, again due in the general enriched categories context to [KS05], although the posets case is much simpler.

It is convenient to treat a “class of meets” as simply the order-dual of a “class of joins”. Thus, given a join doctrine  $\Phi$ , we will refer to  $\Phi^{\text{op}} := \{\phi^{\text{op}} \mid \phi \in \Phi\}$  as a **meet doctrine**, and a meet indexed by  $\phi^{\text{op}} \in \Phi^{\text{op}}$  as a  $\Phi^{\text{op}}$ -**meet**. A poset with all  $\Phi^{\text{op}}$ -meets is a  $\Phi^{\text{op}}$ -**inflattice**, with the category of all such denoted  $\Phi^{\text{op}}\text{Inf}$ . A  $\Phi^{\text{op}}$ -**filter** is an upper sub- $\Phi^{\text{op}}$ -inflattice. The free  $\Phi^{\text{op}}$ -inflattice generated by a poset  $X$  is  $\Phi(X^{\text{op}})^{\text{op}}$ .

3.1. DEFINITION. [see [KS05]] *For two join doctrines  $\Phi, \Psi$ , where we regard  $\Psi^{\text{op}}$  as a meet doctrine, to say that  $\Psi^{\text{op}}$ -**meets commute with  $\Phi$ -joins in  $\mathbf{2}$**  means that for any posets  $X, Y$ ,*

$$\forall \phi \in \Phi(Y) \forall \psi \in \Psi(X) \forall F : X^{\text{op}} \times Y \rightarrow \mathbf{2} \left( \bigwedge_{x \in \psi} \bigvee_{y \in \phi} F(x, y) = \bigvee_{y \in \phi} \bigwedge_{x \in \psi} F(x, y) \right)$$

(where  $F$  runs over monotone maps). By currying  $F$ , this is equivalent to

$$\begin{aligned} \forall \phi \in \Phi(Y) \forall \psi \in \Psi(X) \forall f : Y \rightarrow \mathcal{L}(X) \left( \psi \subseteq \bigcup_{y \in \phi} f(y) \iff \exists y \in \phi (\psi \subseteq f(y)) \right) \\ \iff \forall \psi \in \Psi(X) (\psi \in \mathcal{L}(X) \text{ is } \Phi\text{-compact}). \end{aligned}$$

We write  $\Phi^*(X) := \mathcal{L}(X)_{\Phi}$  for the  $\Phi$ -compact lower sets  $\psi \subseteq X$ , i.e., those indexing meets commuting with  $\Phi$ -joins in  $\mathbf{2}$ . Note that by order-duality, the roles of  $\Phi, \Psi$  may be swapped. Thus

$$\Psi^{\text{op}}\text{-meets commute with } \Phi\text{-joins in } \mathbf{2} \iff \Psi \subseteq \Phi^* \iff \Phi \subseteq \Psi^* \text{ (as submonads of } \mathcal{L}\text{)}.$$

3.2. **REMARK.** The above definition of  $\Phi^*$ , which follows [KS05], yields *a priori* a submonad of  $\mathcal{L}$ . But such a submonad automatically obeys the saturation condition  $(*)$  of Remark 2.3, since given an order-embedding  $i : X \hookrightarrow X'$  and poset  $Y$ , a monotone  $F : X^{\text{op}} \times Y \rightarrow 2$  may be extended along  $i$  to  $F' : X'^{\text{op}} \times Y \rightarrow 2$  (e.g., the left Kan extension  $F'(x', y) := \bigvee_{x \in i^{-1}(\uparrow x')} F(x, y)$ ), so that for  $\psi \in \mathcal{L}(X)$ , the  $\psi^{\text{op}}$ -meet of  $F$  commutes with all  $\Phi$ -joins iff the  $i_*(\psi)^{\text{op}}$ -meet of  $F'$  does. Thus by Remark 2.3, we may equally well regard  $\Phi^*$  as a class of posets. Namely, for a poset  $\psi$ ,

$$\begin{aligned} \psi \in \Phi^* &\iff \psi \in \Phi^*(\psi) = \mathcal{L}(\psi)_{\Phi} \\ &\iff \text{whenever } \psi \text{ is a } \Phi\text{-union of lower subsets, one of them is } \psi. \end{aligned}$$

Note moreover that this reasoning applies to  $\Phi^*$  even if  $\Phi$  is only a submonad of  $\mathcal{L}$  to begin with; this justifies our claim from Remark 2.3 that for our duality-theoretic purposes, it suffices to consider “join doctrines” which are classes of posets, rather than arbitrary submonads of  $\mathcal{L}$  as in [KS05].

3.3. **REMARK.**  $\Phi$ -joins commute with  $\Psi^{\text{op}}$ -meets in  $2$  iff they do in the unit interval  $\mathbb{I}$ . This follows from the facts that  $2$  is a complete sublattice of  $\mathbb{I}$ , while  $\mathbb{I}$  is a complete lattice homomorphic image via  $\bigvee : \mathcal{L}(\mathbb{I}) \rightarrow \mathbb{I}$  (by complete distributivity, Example 2.11) of a complete sublattice  $\mathcal{L}(\mathbb{I}) \subseteq 2^{\mathbb{I}}$ .

3.4. **REMARK.** If  $\phi \in \Psi^*(X)$  for a  $\Psi$ -suplattice  $X$ , then by considering the indicator function of  $\leq \subseteq X^{\text{op}} \times X$ , we get that  $\phi$  must be a  $\Psi$ -ideal. (The converse is false in general: for  $\Psi =$  directed posets, a  $\Psi$ -ideal is a Scott-closed subset, but only finite meets commute with directed joins.)

3.5. **PROPOSITION.** [KS05, 8.9, 8.11, 8.13] *Let  $\Phi, \Psi$  be two join doctrines such that  $\Psi^{\text{op}}$ -meets commute with  $\Phi$ -joins in  $2$ . The following are equivalent:*

- (i) *For every poset  $X$ ,  $\mathcal{L}(X)$  is generated under  $\Phi$ -joins by  $\Psi(X) \subseteq \mathcal{L}(X)_{\Phi}$ .*
- (ii) *For every  $\Psi$ -suplattice  $X$ ,  $\Phi(X)$  consists precisely of all  $\Psi$ -ideals in  $X$ .*
- (iii) *For every poset  $X$ , there is a sub- $\Psi$ -suplattice  $\Psi'(X) \subseteq \mathcal{L}(X)$  containing all principal ideals  $\downarrow x$  (e.g.,  $\Psi'(X) = \mathcal{L}(X)$  or  $\Psi'(X) = \Psi(X)$ ) such that  $\Phi(\Psi'(X))$  contains all  $\Psi$ -ideals in  $\Psi'(X)$ .*

*If these hold, then in fact  $\Psi(X) = \mathcal{L}(X)_{\Phi} = \Phi^*(X)$ , whence  $\mathcal{L}(X) \cong \Phi(\Psi(X))$  is  $\Phi$ -algebraic, whence in particular  $\Phi$  is a continuous join doctrine; and similarly  $\Phi = \Psi^*$ .*

If these hold, we call  $\Phi$  a **sound join doctrine, dual to the sound meet doctrine  $\Psi^{\text{op}}$** . Thus,  $\Phi$  is a sound join doctrine iff  $\mathcal{L}(X) \cong \Phi(\Phi^*(X))$ , iff  $\Phi(X)$  contains every  $\Phi^*$ -ideal in a  $\Phi^*$ -suplattice  $X$ . (Warning: this notion is *not* preserved under swapping  $\Phi, \Psi$ , in contrast to Definition 3.1.)

PROOF. (ii)  $\implies$  (iii) is obvious.

(iii)  $\implies$  (i): For any  $\theta \in \mathcal{L}(X)$ , clearly  $\Psi'(X) \cap \downarrow\theta = \{\psi \in \Psi'(X) \mid \psi \subseteq \theta\}$  is a  $\Psi$ -ideal in  $\Psi'(X)$ , thus by (iii) is in  $\Phi(\Psi'(X))$ ; and its union is  $\theta$ , which is thus a  $\Phi$ -join of elements of  $\Psi(X)$ .

(i)  $\implies$  (ii): For every  $\theta \in \mathcal{L}(X)$ , the  $\Psi$ -ideal  $\langle \theta \rangle$  it generates is in  $\Phi(X)$ : this is true for  $\theta \in \Psi(X)$  since  $\langle \theta \rangle = \downarrow \bigvee \theta$ , and is true for a  $\Phi$ -join  $\theta = \bigcup_i \theta_i$  if it is true for each  $\theta_i$  since  $\langle \theta \rangle = \bigcup_i \langle \theta_i \rangle$  (using that  $\Psi^{\text{op}}$ -meets commute with  $\Phi$ -joins in 2), thus is true for all  $\theta \in \mathcal{L}(X)$  by (i). Conversely, as noted above, every  $\phi \in \Phi(X)$  is a  $\Psi$ -ideal.

The last sentence follows from (i), (ii), and Remark 3.4, which imply that  $\Phi(X) = \Psi^*(X)$  for a  $\Psi$ -suplattice  $X$ , hence for every poset  $X$  by applying  $(*)$  in Remark 2.3 to  $\downarrow : X \rightarrow \Psi(X)$ .  $\blacksquare$

3.6. LEMMA. *For any join doctrine  $\Phi$ , we have  $\omega \in \Phi$  iff  $\omega \notin \Phi^*$ .*

PROOF.  $\omega \notin \Phi \cap \Phi^*$  since  $\omega$ -joins do not commute with  $\omega^{\text{op}}$ -meets in 2. If  $\omega \notin \Phi^*$ , i.e.,  $\omega \in \mathcal{L}(\omega)$  is not  $\Phi$ -compact, then  $\omega$  is a  $\Phi$ -union of proper lower subsets of  $\omega$ ; the order-type of this union must clearly be  $\omega$ . (This argument is due to the referee; my original proof assumed soundness of  $\Phi$ .)  $\blacksquare$

3.7. COROLLARY. [generalized Hofmann–Mislove–Stralka duality] *Let  $\Phi$  be a sound join doctrine, dual to the meet doctrine  $\Psi^{\text{op}} = \Phi^{*\text{op}}$ . We have a dual equivalence of categories*

$$\Phi\text{AlgLat}^{\text{op}} \begin{array}{c} \xrightarrow{\Phi\text{AlgLat}(-,2)} \\ \xleftarrow{\Psi^{\text{op}}\text{Inf}(-,2)} \end{array} \Psi^{\text{op}}\text{Inf}.$$

*We may replace  $\Phi\text{AlgLat}$  with  $\Phi\text{CtsLat}$  iff  $\omega \notin \Phi$ , i.e.,  $\omega \in \Psi$ .*

PROOF. For a  $\Phi$ -algebraic lattice  $X$ , a morphism  $X \rightarrow 2$  is the indicator function of  $\uparrow x$  for  $\Phi$ -algebraic  $x$ . For a  $\Psi^{\text{op}}$ -inflattice  $A$ , a morphism  $A \rightarrow 2$  is the indicator function of a  $\Psi^{\text{op}}$ -filter. So we have

$$\Phi\text{AlgLat}(X, 2) \cong X_{\Phi}^{\text{op}}, \quad \Psi^{\text{op}}\text{Inf}(A, 2) \cong \Phi(A^{\text{op}}).$$

Now the adjunction (co)unit on the left is given by, for  $X \in \Phi\text{AlgLat}$ , the evaluation map

$$\begin{aligned} X &\longrightarrow \Psi^{\text{op}}\text{Inf}(\Phi\text{AlgLat}(X, 2), 2) \\ x &\longmapsto (f \mapsto f(x)), \end{aligned}$$

which via the above isomorphisms becomes the canonical isomorphism  $X \cong \Phi(X_{\Phi})$  characterizing algebraicity. Similarly, for  $A \in \Psi^{\text{op}}\text{Inf}$ , the unit  $A \rightarrow \Phi\text{AlgLat}(\Psi^{\text{op}}\text{Inf}(A, 2), 2)$  is the canonical isomorphism  $A^{\text{op}} \cong \Phi(A^{\text{op}})_{\Phi}$ . By Corollary 2.13,  $\Phi\text{AlgLat} = \Phi\text{CtsLat}$  iff  $\mathbb{I}$  is  $\Phi$ -algebraic, iff  $\omega \notin \Phi$ .  $\blacksquare$

3.8. **EXAMPLE.**  $\Phi = \text{directed posets}$  forms a sound join doctrine, dual to  $\Psi^{\text{op}} = \text{“finite meets”}$ , i.e.,  $\Psi = \text{the class of posets with finite cofinality}$ . In this case, Corollary 3.7 becomes the classical Hofmann–Mislove–Stralka duality [HMS74] between (unital) meet-semilattices and algebraic lattices.

Similarly, the join doctrine  $\Phi$  of  $\kappa$ -directed posets for an uncountable regular cardinal  $\kappa$  is sound, dual to  $\kappa$ -ary meets. But since  $\omega \notin \Phi$  for uncountable  $\kappa$ , we get a duality between  $\kappa$ -meet-semilattices and  $\kappa$ -continuous lattices.

We now show that there are very few sound join doctrines  $\Phi \ni \omega$ , for which  $\Phi\text{AlgLat} \neq \Phi\text{CtsLat}$ : essentially, they are only the classical cases of continuous and completely distributive lattices (Examples 2.9 and 2.10), plus the minor variations including/excluding empty joins.

3.9. **THEOREM.** *There are precisely 4 sound join doctrines  $\Phi \ni \omega$ , dual to  $\Psi^{\text{op}}$ :*

- (i)  $\Phi = \text{directed posets}$ ,  $\Psi = \text{posets with finite cofinality}$ ;
- (ii)  $\Phi = \text{empty or directed posets}$ ,  $\Psi = \text{nonempty posets with finite cofinality}$ ;
- (iii)  $\Phi = \text{nonempty posets}$ ,  $\Psi = \text{posets which are empty or have greatest element}$ ;
- (iv)  $\Phi = \text{all posets}$ ,  $\Psi = \text{posets with greatest element}$ .

**PROOF.** It is well-known and easily seen that each of these 4 cases is sound; we show the converse.

First, we show that  $\Phi$  must contain every directed poset, i.e., every poset in  $\Psi$  must have finite cofinality. For every set  $X$ ,  $\Phi$  contains the finite powerset  $\mathcal{P}_\omega(X)$ , since this is a  $\Psi$ -ideal in the full powerset  $\mathcal{P}(X)$ , since by Proposition 2.12(i) (applied to  $\Psi \not\ni \omega$ ), every  $\psi \in \Psi(\mathcal{P}_\omega(X))$  can have neither a strictly increasing sequence nor infinitely many maximal elements, thus must be finite. Now for every join-semilattice  $X$ , we have a monotone surjection  $\bigvee : \mathcal{P}_\omega(X) \rightarrow X$ , whence  $X \in \Phi$ . Since every directed poset  $\phi$  is cofinal in the free join-semilattice it generates, it follows that  $\phi \in \Phi$ .

So  $\Psi$  is determined by the finite antichains  $n$  in it. If some  $n > 1$  is in  $\Psi$ , then by induction so is each  $n^k \cong \bigsqcup_{i \in n} n^{k-1}$ ; now every  $m \geq 1$  admits a surjection  $n^k \rightarrow m$ , whence  $m \in \Psi$ . ■

#### 4. $\mathbb{U}$ -posets

Henceforth, we assume  $\Phi \ni \omega$  is a sound join doctrine, dual to  $\Psi^{\text{op}}$ , so one of the cases in Theorem 3.9. Then Hofmann–Mislove–Stralka duality does not apply to all  $\Phi$ -continuous lattices, and so we would like to formulate a duality based on morphisms to  $\mathbb{I}$  instead of  $\mathbb{2}$ .

By Remark 3.3, the dual algebra  $\Phi\text{CtsLat}(X, \mathbb{I})$  will still be equipped with  $\Psi^{\text{op}}$ -meets. But these are not all the operations on  $\mathbb{I}$  commuting with the  $\Phi$ -continuous lattice operations: clearly any complete lattice homomorphism  $\mathbb{I} \rightarrow \mathbb{I}$  does as well. We thus introduce the following notions:

4.1. DEFINITION. Let  $\mathbb{U} := \mathbf{CLat}(\mathbb{I}, \mathbb{I})$  denote the partially ordered monoid of all complete lattice homomorphisms  $\mathbb{I} \rightarrow \mathbb{I}$ , i.e., surjective monotone maps.

A  $\mathbb{U}$ -**poset** is a poset equipped with a monotone (in both variables) action of the monoid  $\mathbb{U}$ . Denote the category of these (and equivariant monotone maps) by  $\mathbf{UPos}$ .

A  $\mathbb{U}$ - $\Psi^{\text{op}}$ -**inflattice** is a  $\mathbb{U}$ -poset which is also a  $\Psi^{\text{op}}$ -inflattice such that the action of each  $u \in \mathbb{U}$  preserves  $\Psi^{\text{op}}$ -meets. Denote the category of these by  $\mathbf{U}\Psi^{\text{op}}\mathbf{Inf}$ .

4.2. DEFINITION. Let  $\dot{+}, \dot{-}$  denote truncated  $+, -$  on  $\mathbb{I}$ ; note that they obey the adjunction

$$(4.3) \quad r \dot{-} s \leq t \iff r \leq s \dot{+} t.$$

For a  $\mathbb{U}$ -poset  $A$  and  $a, b \in A$ , define

$$\begin{aligned} a \leq_r b &: \iff \forall u, v \in \mathbb{U} (u((-) \dot{+} r) \leq v \implies u(a) \leq v(b)), \\ \rho(a, b) &:= \bigwedge \{r \in \mathbb{I} \mid a \leq_r b\}, \\ d(a, b) &:= \rho(a, b) \vee \rho(b, a). \end{aligned}$$

4.4. REMARK. In the definition of  $\leq_r$ , instead of testing  $\forall u, v$ , it is enough to test any particular  $u \in \mathbb{U}$  which restricts to an order-isomorphism  $u : [r, 1] \cong [0, 1]$  (e.g., the linear such isomorphism extended by 0 on  $[0, r]$ ), so that  $v := u((-) \dot{+} r) \in \mathbb{U}$ . Indeed, for any other  $u', v' \in \mathbb{U}$  with  $u'((-) \dot{+} r) \leq v'$ , there is  $w \in \mathbb{U}$  with  $u' = w \circ u$ , whence  $u'(a) = w(u(a)) \leq w(v(a)) \leq v'(a)$ .

4.5. REMARK. There is an evident order-duality for  $\mathbb{U}$ -posets  $A$ : let  $u \in \mathbb{U}$  act on the order-dual  $A^{\text{op}}$  via  $1 - u(1 - (-))$ ; this reverses each  $\leq_r$ , and turns  $\rho$  into  $\rho^{\text{op}}(a, b) := \rho(b, a)$ .

Intuitively,  $a \leq_r b$  means “ $a \leq b \dot{+} r$ ”. The following properties justify this intuition:

4.6. PROPOSITION. In  $\mathbb{I}$ , we have  $a \leq_r b \iff a \leq b \dot{+} r$ , whence  $\rho(a, b) = a \dot{-} b$  and  $d(a, b) = |a - b|$ .

PROOF. If  $a \leq b \dot{+} r$ , then for every  $u, v \in \mathbb{U}$  with  $u((-) \dot{+} r) \leq v$ , we have  $u(a) \leq u(b \dot{+} r) \leq v(b)$ .

For the converse, the case  $r = 1$  is vacuous; thus we may assume  $r < 1$ . Note that  $(-) \dot{+} r : \mathbb{I} \rightarrow \mathbb{I}$  can be written as  $u^\times \circ v$  where  $v := 1 \wedge (-)/(1 - r)$ ,  $u := v((-) \dot{-} r)$ , and  $u^\times$  is the right adjoint of  $u$ . Now from  $a \leq_r b$  and  $u((-) \dot{+} r) = v$ , we get  $u(a) \leq v(b)$ , whence  $a \leq u^\times(v(b)) = b \dot{+} r$ .  $\blacksquare$

4.7. LEMMA. In every  $\mathbb{U}$ -poset  $A$ , we have the following, for  $r, s, t \in \mathbb{I}$ ,  $u, v \in \mathbb{U}$ ,  $a, b, c \in A$ :

$$(a) \quad r \leq s \text{ and } a \leq_r b \implies a \leq_s b.$$

$$(b) \quad \leq_0 \text{ is the same as } \leq.$$

$$(c) \quad a \leq_r b \leq_s c \implies a \leq_{r+s} c.$$

(d)  $\rho$  is a pseudoquasimetric:  $\rho(a, a) = 0$ , and  $\rho(a, b) + \rho(b, c) \geq \rho(a, c)$ . Thus,  $d$  is a pseudometric.

- (e)  $u((-) \dot{+} r) \leq v \dot{+} s$  and  $a \leq_r b \implies u(a) \leq_s v(b)$ . Thus,  $\rho(u(a), v(a)) \leq \rho(u, v) := \bigvee (u \dot{-} v)$ , i.e., the  $\mathbb{U}$ -action is 1-Lipschitz in the first variable with respect to the  $\ell^\infty$ -quasimetric on  $\mathbb{U}$ . Moreover, if  $u \in \mathbb{U}$  is uniformly continuous with modulus  $\mu : \mathbb{I} \rightarrow \mathbb{I}$ , i.e.,  $u(r) \dot{-} u(s) \leq \mu(r \dot{-} s)$ , then the action of  $u$  is uniformly continuous with the same modulus:  $\rho(u(a), u(b)) \leq \mu(\rho(a, b))$ .
- (f)  $u^\times((-) \dot{+} r) \leq v \dot{+} s$  and  $u(a) \leq_r b \implies a \leq_s v(b)$  (where  $u^\times =$  right adjoint of  $u$ ).

In a  $\mathbb{U}$ - $\Psi^{\text{op}}$ -inflattice, we moreover have, for  $\psi, \psi' \in \Psi(A^{\text{op}})$ :

- (g)  $a \leq_r \bigwedge \psi \iff \forall b \in \psi (a \leq_r b)$ . Thus,  $\rho(\bigwedge \psi, \bigwedge \psi') \leq \bigwedge_{a \in \psi} \bigvee_{b \in \psi'} \rho(a, b)$ .

PROOF. (a) and (b) are straightforward, as is (d) given the previous parts.

(c) For  $u, w \in \mathbb{U}$  with  $u((-) \dot{+} (r \dot{+} s)) \leq w$ , we have  $v := u((-) \dot{+} r) \in \mathbb{U}$  with  $u((-) \dot{+} r) \leq v$  and  $v((-) \dot{+} s) \leq w$ , whence  $u(a) \leq v(b) \leq w(c)$ .

(e) For  $u', v' \in \mathbb{U}$  with  $u'((-) \dot{+} s) \leq v'$ , we have  $u'(u((-) \dot{+} r)) \leq u'(v(-) \dot{+} s) \leq v' \circ v$ , whence  $u'(u(a)) \leq v'(v(b))$ . For the last assertion:  $u(r) \dot{-} u(s) \leq \mu(r \dot{-} s)$  means  $u((-) \dot{+} r) \leq u(-) \dot{+} \mu(r)$ .

(f) The assumption is equivalent to  $(-) \dot{-} s \leq v(u(-) \dot{-} r)$ ; thus for  $u', v' \in \mathbb{U}$  with  $u'((-) \dot{+} s) \leq v'$ , we have  $u' \leq v'((-) \dot{-} s) \leq v'(v(u(-) \dot{-} r))$ , whence  $u'(a) \leq v'(v(u(a) \dot{-} r)) \leq v'(v(b))$ .

(g)  $\implies$  and the last assertion follow from (c). For  $\iff$ : for  $u, v \in \mathbb{U}$  with  $u((-) \dot{+} r) \leq v$ , we have  $u(a) \leq \bigwedge_{b \in \psi} v(b) = v(\bigwedge \psi)$ .  $\blacksquare$

For general background on (pseudo)quasimetrics, see e.g., [Kün09]. A pseudoquasimetric  $\rho$  as above induces a topology, where a basic neighborhood of  $a \in A$  is  $\{b \in A \mid \rho(a, b) < r\}$  for some  $r > 0$ . Thus the closure of  $B \subseteq A$  is the set of all  $a \in A$  such that

$$\rho(a, B) = \bigwedge_{b \in B} \rho(a, b) = 0,$$

which is in particular a lower set. To avoid confusion, we will call a closed set in this topology a  $\rho$ -**closed lower set**, and denote the set of all such by  $\overline{\mathcal{L}}(A) \subseteq \mathcal{L}(A)$ . We will also say  $\rho^{\text{op}}$ -**closed upper set**  $B \subseteq A$  for the order-dual notion, i.e., if  $\rho(B, a) = 0$  then  $a \in B$ ; the set of all such is thus  $\overline{\mathcal{L}}(A^{\text{op}})$ . For a  $\mathbb{U}$ - $\Psi^{\text{op}}$ -inflattice  $A$ , recalling that  $\Phi(A^{\text{op}})$  consists of  $\Psi^{\text{op}}$ -filters by soundness, let

$$\overline{\Phi}(A^{\text{op}}) := \Phi(A^{\text{op}}) \cap \overline{\mathcal{L}}(A^{\text{op}})$$

denote the  $\rho^{\text{op}}$ -**closed  $\Psi^{\text{op}}$ -filters** in  $A$ .

4.8. LEMMA. *If  $\phi \in \overline{\Phi}(A^{\text{op}})$  is a  $\Psi^{\text{op}}$ -filter, then so is the  $\rho^{\text{op}}$ -closure  $\overline{\phi}$ .*

PROOF. This follows from the facts that  $\Psi^{\text{op}}$  is a class of finite meets by Theorem 3.9, and that  $\Psi^{\text{op}}$ -meets are Lipschitz by Lemma 4.7(g).  $\blacksquare$

As usual for actions, a subset  $B \subseteq A$  of a  $\mathbb{U}$ -poset is  **$\mathbb{U}$ -invariant** if it is closed under the action. For a class of sets  $\Gamma(A)$ , we write  $\Gamma^{\mathbb{U}}(A)$  for the  $\mathbb{U}$ -invariant members, e.g.,  $\mathcal{L}^{\mathbb{U}}(A), \overline{\Phi}^{\mathbb{U}}(A)$ .

4.9. LEMMA. *If  $\phi \in \mathcal{P}^{\mathbb{U}}(A)$  is a  $\mathbb{U}$ -invariant filter base, then its  $\rho^{\text{op}}$ -closure  $\overline{\phi}$  is a  $\mathbb{U}$ -invariant  $\Psi^{\text{op}}$ -filter, hence is the  $\mathbb{U}$ -invariant  $\rho$ -closed  $\Psi^{\text{op}}$ -filter generated by  $\phi$ .*

PROOF. By uniform continuity of the action of each  $u$  (Lemma 4.7(e)),  $\overline{\phi}$  is  $\mathbb{U}$ -invariant. It is also upper, since every  $\rho^{\text{op}}$ -closed set is, thus it is also the  $\rho^{\text{op}}$ -closure of the upward closure of  $\phi$ , which is a  $\Psi^{\text{op}}$ -filter since  $\Psi^{\text{op}}$ -meets are finite by Theorem 3.9, whence so is  $\overline{\rho}$  by the preceding lemma.  $\blacksquare$

4.10. PROPOSITION. *For a  $\mathbb{U}$ - $\Psi^{\text{op}}$ -inflattice  $A$ , we have an order-isomorphism*

$$\begin{aligned} \mathbb{U}\Psi^{\text{op}}\text{Inf}(A, \mathbb{I}) &\cong \overline{\Phi}^{\mathbb{U}}(A^{\text{op}}) = \{\mathbb{U}\text{-invariant } \rho^{\text{op}}\text{-closed } \Psi^{\text{op}}\text{-filters in } A\} \\ f &\mapsto f^{-1}(1) \\ 1 - \rho(\phi, -) &\leftrightarrow \phi. \end{aligned}$$

PROOF. For ease of notation, we will prove that dually, for a  $\mathbb{U}$ - $\Psi$ -suplattice  $A$ ,

$$\begin{aligned} \mathbb{U}\Psi\text{Sup}(A, \mathbb{I})^{\text{op}} &\cong \overline{\Phi}^{\mathbb{U}}(A) = \{\mathbb{U}\text{-invariant } \rho\text{-closed } \Psi\text{-ideals in } A\} \\ f &\mapsto f^{-1}(0) \\ \rho(-, \phi) &\leftrightarrow \phi. \end{aligned}$$

It is immediate from the definitions that for a  $\mathbb{U}$ -equivariant  $\Psi$ -join-preserving  $f : A \rightarrow \mathbb{I}$ ,  $f^{-1}(0) \subseteq A$  is  $\mathbb{U}$ -invariant  $\rho$ -closed lower, and also that a  $\rho$ -closed lower  $\phi \subseteq A$  is equal to  $\rho(-, \phi)^{-1}(0)$ .

We now check that for a  $\mathbb{U}$ -invariant  $\Psi$ -ideal  $\phi \subseteq A$ ,  $\rho(-, \phi) : A \rightarrow \mathbb{I}$  is  $\mathbb{U}$ -equivariant  $\Psi$ -join-preserving (it is clearly monotone). For  $\psi \in \Psi(A)$ ,

$$\begin{aligned} \rho(\bigvee \psi, \phi) &= \bigwedge_{b \in \phi} \bigvee_{a \in \psi} \rho(a, b) \quad \text{by the dual of Lemma 4.7(g)} \\ &= \bigvee_{a \in \psi} \bigwedge_{b \in \phi} \rho(a, b) \quad \text{because } \Phi \subseteq \Psi^* \text{ (Remark 3.3)} \\ &= \bigvee_{a \in \psi} \rho(a, \phi); \end{aligned}$$

thus  $\rho(-, \phi)$  preserves  $\Psi$ -joins. To check  $\mathbb{U}$ -equivariance: let  $u \in \mathbb{U}$  and  $a \in A$ . We have

$$\begin{aligned} \rho(u(a), \phi) &= \bigwedge_{b \in \phi} \rho(u(a), b) = \bigwedge \{r \in \mathbb{I} \mid u(a) \leq_r b \in \phi\}, \\ u(\rho(a, \phi)) &= u(\bigwedge_{b \in \phi} \rho(a, b)) = \bigwedge_{b \in \phi} u(\rho(a, b)) = \bigwedge \{u(r) \mid a \leq_r b \in \phi\}. \end{aligned}$$

For each  $a \leq_r b \in \phi$ , find

$$u((-) \dot{+} r) \dot{\div} u(r) \leq v \in \mathbb{U},$$

whence  $u(a) \leq_{u(r)} v(b) \in \phi$  by Lemma 4.7(e); this proves  $u(\rho(a, \phi)) \geq \rho(u(a), \phi)$ . Conversely, for  $u(a) \leq_r b \in \phi$  with  $r < 1$ , let  $u^\times$  be the right adjoint of  $u$ , and similarly to before, find

$$u^\times((-) \dot{+} r) \dot{\div} u^\times(r) \leq v \in \mathbb{U},$$

whence  $a \leq_{u \times (r)} v(b) \in \phi$  by Lemma 4.7(f), whence  $u(\rho(a, \phi)) \leq r$ ; so  $\rho(u(a), \phi) \geq u(\rho(a, \phi))$ .

Finally, we check that for  $\mathbb{U}$ -equivariant monotone  $f : A \rightarrow \mathbb{I}$ , we have  $f = \rho(-, f^{-1}(0))$ . We have  $\leq$  since  $f$  is 1-Lipschitz. Conversely, for  $a \in A$  with  $f(a) < 1$ , find  $(-) \dot{+} f(a) \leq u \in \mathbb{U}$  with  $u(f(a)) = 0$ ; then  $a \leq_{f(a)} u(a)$  by Lemma 4.7(e), so  $\rho(a, f^{-1}(0)) \leq \rho(a, u(a)) \leq f(a)$ .  $\blacksquare$

The  $\mathbb{U}$ -poset  $\mathbb{I}$  obeys the following additional axioms, which must thus also hold in the dual of a  $\Phi$ -continuous lattice:

4.11. DEFINITION. We call a  $\mathbb{U}$ -poset  $A$  **Archimedean** if it obeys

$$\forall r > 0 (a \leq_r b) \implies a \leq b.$$

We call  $A$  (**Cauchy-**)**complete** if it is Archimedean and also complete in the metric  $d$ .

4.12. DEFINITION. We call a  $\mathbb{U}$ -poset  $A$  **unstackable** if for any  $0 < r < 1$  and  $u, v \in \mathbb{U}$  restricting to order-isomorphisms  $u : [0, r] \cong [0, 1]$  and  $v : [r, 1] \cong [0, 1]$ , we have

$$u(a) \leq u(b) \text{ and } v(a) \leq v(b) \implies a \leq b.$$

We call  $A$  **stackable** if it is unstackable and for  $r, u, v$  as above and  $a, b \in A$  such that  $v'(b) \leq u'(a)$  for all  $u', v' \in \mathbb{U}$ , there is a (unique, by unstackability)  $c \in A$  with  $u(c) = a$  and  $v(c) = b$ .

Intuitively, stackability means that, thinking of  $A$  as the dual of a  $\Phi$ -continuous lattice  $X$ , we may specify  $A \ni a : X \rightarrow \mathbb{I}$  via its restrictions to its sublevel and superlevel sets  $a^{-1}([0, r]), a^{-1}([r, 1])$ .

4.13. REMARK. As in Remark 4.4, it is enough to take some particular  $u, v$  above. Also, it is enough to take some particular  $r$  (e.g.,  $1/2$ ), since we may move  $r$  around via an order-isomorphism  $\mathbb{I} \cong \mathbb{I}$ .

4.14. LEMMA. If  $A$  is (un)stackable, then more generally, for  $0 = r_0 < r_1 < \dots < r_n = 1$  and  $u_1, \dots, u_n \in \mathbb{U}$  restricting to  $u_i : [r_{i-1}, r_i] \cong [0, 1]$ , for  $a_1, \dots, a_n \in A$  such that  $v'(a_{i+1}) \leq u'(a_i)$  for all  $u', v' \in \mathbb{U}$ , there is (at most one, depending monotonically on  $(a_1, \dots, a_n)$ )  $a \in A$  with  $u_i(a) = a_i$ .

PROOF. By a straightforward induction on  $n$ .  $\blacksquare$

4.15. LEMMA. If  $A$  is unstackable, then more generally, for  $0 \leq r = r_0 < r_1 < \dots < r_n = 1$  and  $u_1, \dots, u_n \in \mathbb{U}$  with  $u_i : [r_{i-1}, r_i] \cong [0, 1]$ , so that  $u_i((-) \dot{+} r) \in \mathbb{U}$ , for any  $a, b \in A$ , we have

$$u_1(a) \leq u_1(b \dot{+} r) \text{ and } \dots \text{ and } u_n(a) \leq u_n(b \dot{+} r) \implies a \leq_r b.$$

PROOF. By Remark 4.4, it suffices to check that for  $w \in \mathbb{U}$  with  $w : [r, 1] \cong [0, 1]$ , we have  $w(a) \leq w(b \dot{+} r)$ ; this follows from applying the preceding lemma to  $u_i \circ w^{-1} : [w(r_{i-1}), w(r_i)] \cong [0, 1]$ .  $\blacksquare$

## 5. The duality

Let  $\mathbf{CStU}\Psi^{\text{op}}\mathbf{Inf} \subseteq \mathbf{U}\Psi^{\text{op}}\mathbf{Inf}$  denote the full subcategory of complete stackable  $\mathbf{U}$ - $\Psi^{\text{op}}$ -inflattices. Since the  $\Phi$ -continuous lattice and  $\mathbf{U}$ - $\Psi^{\text{op}}$ -inflattice structures on  $\mathbb{I}$  commute, we have a dual adjunction

$$(5.1) \quad \Phi\mathbf{CtsLat}^{\text{op}} \begin{array}{c} \xrightarrow{\Phi\mathbf{CtsLat}(-, \mathbb{I})} \\ \xleftarrow{\mathbf{U}\Psi^{\text{op}}\mathbf{Inf}(-, \mathbb{I})} \end{array} \mathbf{CStU}\Psi^{\text{op}}\mathbf{Inf} \subseteq \mathbf{U}\Psi^{\text{op}}\mathbf{Inf}.$$

5.2. **THEOREM.** *For every  $\Phi$ -continuous lattice  $X$ , the evaluation map*

$$\begin{aligned} \eta : X &\longrightarrow \mathbf{U}\Psi^{\text{op}}\mathbf{Inf}(\Phi\mathbf{CtsLat}(X, \mathbb{I}), \mathbb{I}) \\ x &\longmapsto (f \mapsto f(x)), \end{aligned}$$

*which is the (co)unit on the left side of the above adjunction, is an order-isomorphism.*

**PROOF.** Via Propositions 2.7 and 4.10,  $\eta$  corresponds to the map

$$\begin{aligned} \tilde{\eta} : X &\longrightarrow \overline{\Phi}^{\mathbf{U}}(\ll^{\Phi}\mathbf{Sup}(\mathbb{I}, X)) \subseteq \mathcal{L}(\ll^{\Phi}\mathbf{Sup}(\mathbb{I}, X)) \\ x &\longmapsto \{f^+ \in \ll^{\Phi}\mathbf{Sup}(\mathbb{I}, X) \mid f^+(1) \leq x\} \end{aligned}$$

whose left adjoint is easily seen to be

$$\begin{aligned} \tilde{\eta}^+ : \mathcal{L}(\ll^{\Phi}\mathbf{Sup}(\mathbb{I}, X)) &\longrightarrow X \\ \phi &\longmapsto \bigvee_{f^+ \in \phi} f^+(1). \end{aligned}$$

That  $x \leq \tilde{\eta}^+(\tilde{\eta}(x))$  is Urysohn's lemma for  $\Phi$ -continuous lattices; see [G<sup>+</sup>03, IV-3.1, IV-3.32], [Joh82, VII 1.14, 3.2], [Xu95]. Since  $x = \bigvee \downarrow x$ , it suffices to show that for each  $y \ll x$  there is  $f^+ \in \ll^{\Phi}\mathbf{Sup}(\mathbb{I}, X)$  with  $y \leq f^+(1) \leq x$ . Let  $\mathbb{I}_2 \subseteq \mathbb{I}$  be the dyadic rationals, define  $g : \mathbb{I}_2 \rightarrow X$  by  $g(0) := y$ ,  $g(1) := x$ , and inductively using interpolation (Proposition 2.6(c)) so that  $r < s \implies g(r) \ll g(s)$ ; then  $f^+(r) := \bigvee g(\mathbb{I}_2 \cap [0, r))$  works.

Now let  $\phi \in \overline{\Phi}^{\mathbf{U}}(\ll^{\Phi}\mathbf{Sup}(\mathbb{I}, X))$ ; we must show  $\tilde{\eta}(\tilde{\eta}^+(\phi)) \subseteq \phi$ . Since  $\tilde{\eta}$  preserves  $\Phi$ -joins,

$$\tilde{\eta}(\tilde{\eta}^+(\phi)) = \bigvee_{f^+ \in \phi} \tilde{\eta}(f^+(1)).$$

For each  $f^+ \in \phi$  and  $g^+ \in \tilde{\eta}(f^+(1))$ , i.e.,  $g^+(1) \leq f^+(1)$ , we have  $1 \leq g(f^+(1))$ , thus there is  $g \circ f^+ \geq u \in \mathbf{U}$ , whence  $g \geq u \circ f$ , so  $g^+ \leq (u \circ f)^+ \in \phi$  since  $\phi$  is  $\mathbf{U}$ -invariant; thus  $\tilde{\eta}(f^+(1)) \subseteq \phi$ .  $\blacksquare$

5.3. **THEOREM.** *For every Archimedean unstackable  $\mathbf{U}$ - $\Psi^{\text{op}}$ -inflattice  $A$ , the evaluation map*

$$\begin{aligned} \iota : A &\longrightarrow \Phi\mathbf{CtsLat}(\mathbf{U}\Psi^{\text{op}}\mathbf{Inf}(A, \mathbb{I}), \mathbb{I}) \\ a &\longmapsto (f \mapsto f(a)) \end{aligned}$$

*is an embedding. If  $A$  is stackable, its image is dense; thus if  $A$  is also complete, it is an isomorphism.*

PROOF. Via Propositions 2.7 and 4.10,  $\iota$  corresponds to the map

$$\begin{aligned} \tilde{\iota} : A &\longrightarrow \llcorner^{\Phi} \mathbf{Sup}(\mathbb{I}, \overline{\Phi}^{\mathbb{U}}(A^{\text{op}}))^{\text{op}} \\ a &\longmapsto (r \mapsto \min\{\phi \in \overline{\Phi}^{\mathbb{U}}(A^{\text{op}}) \mid r \leq 1 - \rho(\phi, a)\}). \end{aligned}$$

We claim that in fact, for  $r > 0$ ,  $\tilde{\iota}(a)(r)$  is the  $\rho^{\text{op}}$ -closure  $\overline{U_r(a)}$  of

$$U_r(a) := \{u(a) \mid u \in \mathbb{U} \text{ and } u(r) = 1\}.$$

$\overline{U_r(a)}$  is a  $\mathbb{U}$ -invariant  $\Psi^{\text{op}}$ -filter by Lemma 4.9. Each  $u(a) \in U_r(a)$  is in each  $\phi \in \overline{\Phi}^{\mathbb{U}}(A^{\text{op}})$  with  $r \leq 1 - \rho(\phi, a)$ : if  $u(s) = 1$  for some  $s < r$ , we may let  $b \in \phi$  with  $b \leq_{1-s} a$  to get  $\phi \ni u(b \dot{-} (1-s)) \leq u(a)$ , while if there is no such  $s$ , we may write  $u$  as a limit of  $u_0, u_1, \dots$  for which there are such  $s$ , then use that  $\phi$  is closed. And  $r \leq 1 - \rho(\overline{U_r(a)}, a)$ : letting  $(-)\dot{+}(1-r) \geq u \in \mathbb{U}$  with  $u(r) = 1$ , we have  $U_r(a) \ni u(a) \leq_{1-r} a$  by Lemma 4.7(e). This proves the claim.

Now to show that  $\tilde{\iota}$  is an order-embedding: let  $\tilde{\iota}(a) \geq \tilde{\iota}(b) : \mathbb{I} \rightarrow \overline{\Phi}^{\mathbb{U}}(A^{\text{op}})$ , i.e.,  $\overline{U_r(a)} \supseteq U_r(b)$  for every  $r > 0$ ; since  $A$  is Archimedean, it suffices to show  $a \leq_{2/n} b$  for all  $n \geq 3$ . For  $i = 1, \dots, n$ , let

$$(*) \quad v_i \in \mathbb{U}, \quad v_i : [(i-1)/n, i/n] \cong [0, 1].$$

Then  $v_i(b) \in U_{i/n}(b)$ , so there is  $u_i \in \mathbb{U}$  with  $u_i(i/n) = 1$  such that

$$u_i(a) \leq_{1/n} v_i(b).$$

Let  $u', v' \in \mathbb{U}$  with  $u'((-)\dot{+}1/n) \leq v'$ ; then for  $2 \leq i \leq n-1$ , we have  $v_{i+1}(a) \leq u'(u_i(a)) \leq v'(v_i(b)) \leq v_{i+1}(b \dot{+} 2/n)$  since  $v_{i+1}(i/n) = 0$ ,  $u'(u_i(i/n)) = 1$ ,  $v'(v_i((i-1)/n)) = 0$ , and  $v_{i+1}((i-1)/n \dot{+} 2/n) = 1$ . Thus since  $A$  is unstackable, by Lemma 4.15 we have  $a \leq_{2/n} b$ .

Finally, suppose  $A$  is stackable, and let  $f^+ \in \llcorner^{\Phi} \mathbf{Sup}(\mathbb{I}, \overline{\Phi}^{\mathbb{U}}(A^{\text{op}}))$ , left adjoint to  $f$ ; we will find, for every  $n \geq 2$ , some  $a \in A$  with  $d(\iota(a), f) \leq 2/n$ . For  $i = 1, \dots, n$ , we have  $f^+((i-1)/n) \llcorner f^+(i/n) = \bigvee_{a \in f^+(i/n)} \overline{U_1(a)} = \bigvee_{a \in f^+(i/n)} \bigvee_{r < 1} \overline{U_r(a)}$  (again by Lemma 4.9), whence

$$f^+((i-1)/n) \subseteq \overline{U_{r_i}(a_i)}$$

for some  $a_i \in f^+(i/n)$  and  $r_i < 1$ . Let  $u_i \in \mathbb{U}$  with  $u_i(r_i) = 0$ , and let  $v_i$  as in (\*). Then for  $u' \in \mathbb{U}$ ,

$$f^+((i-1)/n) \subseteq \uparrow u'(u_i(a_i)) \subseteq \overline{U_1(u_i(a_i))},$$

since for  $b \in f^+((i-1)/n) \subseteq \overline{U_{r_i}(a_i)}$ , for every  $s > 0$ , there is  $u'' \in \mathbb{U}$  with  $u''(r_i) = 1$ , whence  $u' \circ u_i \leq u''$ , such that  $u'(u_i(a_i)) \leq u''(a_i) \leq_s b$ , whence  $u'(u_i(a_i)) \leq b$  since  $A$  is Archimedean. In particular, this holds for  $b = v'(u_{i-1}(a_{i-1}))$  for every  $v' \in \mathbb{U}$ , so by Lemma 4.14, there is  $a \in A$  with

$$v_i(a) = u_i(a_i)$$

for each  $i$ . Then

$$U_{i/n}(a) = U_1(v_i(a)) = U_1(u_i(a_i)),$$

since every  $u \in \mathbb{U}$  with  $u(i/n) = 1$  is  $\geq u' \circ v_i$  for some  $u' \in \mathbb{U}$ . We now show that  $d(f, \iota(a)) \leq 2/n$ , in terms of the left adjoints  $f^+, \tilde{\iota}(a)$ : for each  $t \in \mathbb{I}$ , letting  $1 \leq i \leq n$  with  $t \leq i/n \leq t \dot{+} 1/n$ ,

$$\begin{aligned} \tilde{\iota}(a)(t) &= \overline{U_t(a)} \subseteq \overline{U_{i/n}(a)} = \overline{U_1(u_i(a_i))} \subseteq f^+(i/n) \subseteq f^+(t \dot{+} 1/n), \\ f^+(t \dot{-} 1/n) &\subseteq f^+((i-1)/n) \subseteq \overline{U_1(u_i(a_i))} = \overline{U_{i/n}(a)} \subseteq \overline{U_{t+1/n}(a)} = \tilde{\iota}(a)(t \dot{+} 1/n). \quad \blacksquare \end{aligned}$$

**5.4. THEOREM.** *The dual adjunction (5.1) is a dual equivalence of categories between  $\Phi$ -continuous lattices and complete stackable  $\mathbb{U}$ - $\Psi^{\text{op}}$ -inflattices.*  $\blacksquare$

It is worth explicitly restating the duality for the two main examples of  $\Phi$ :

**5.5. COROLLARY.** *Hom into  $\mathbb{I}$  yields a dual equivalence of categories between completely distributive lattices and complete stackable  $\mathbb{U}$ -posets.*  $\blacksquare$

Let us say that a  **$\mathbb{U}$ -meet-semilattice** is a  $\mathbb{U}$ -poset with finite meets preserved by the  $\mathbb{U}$ -action.

**5.6. COROLLARY.** *Hom into  $\mathbb{I}$  yields a dual equivalence of categories between continuous lattices and complete stackable  $\mathbb{U}$ -meet-semilattices.*  $\blacksquare$

We end by showing that in the presence of meets, stackability admits a simpler formulation:

**5.7. DEFINITION.** *Let  $\widehat{\mathbb{U}} := \text{CtsLat}(\mathbb{I}, \mathbb{I}) \supseteq \mathbb{U}$  be the monoid of continuous lattice morphisms  $\mathbb{I} \rightarrow \mathbb{I}$ , i.e., continuous monotone maps preserving 1, but possibly not 0.*

*A  $\widehat{\mathbb{U}}$ -module is a (unital) meet-semilattice with a  $\widehat{\mathbb{U}}$ -action preserving finite meets on both sides.*

**5.8. PROPOSITION.** *The forgetful functor is an isomorphism of categories between complete  $\widehat{\mathbb{U}}$ -modules and complete stackable  $\mathbb{U}$ -meet-semilattices. The  $\leq_r$  relations in a  $\widehat{\mathbb{U}}$ -module are given by*

$$\rho(a, b) \leq r \iff a \leq_r b \iff a \leq b \dot{+} r.$$

**PROOF.** The characterization of  $\leq_r$  is proved as in Proposition 4.6.

Next, an Archimedean  $\widehat{\mathbb{U}}$ -module  $A$  is unstackable as a  $\mathbb{U}$ -poset: by Remark 4.13, it suffices to check that for  $0 < r < 1$ ,  $u := 1 \wedge (-)/r$ , and  $v := ((-) \dot{-} r)/(1 - r)$ , if  $u(a) \leq u(b)$  and  $v(a) \leq v(b)$ , then  $a \leq b$ . Let  $s > 0$ , and let  $r(-) \leq w \in \mathbb{U}$  with equality on  $[0, 1 - s]$ . Then  $1_{\mathbb{I}} \leq (w \circ u) \wedge (v^\times \circ v) \leq (-) \dot{+} rs$ , whence from  $u(a) \leq u(b)$  and  $v(a) \leq v(b)$  we have  $a \leq b \dot{+} rs$ , i.e.,  $a \leq_{rs} b$  by the above. Since  $A$  is Archimedean, it follows that  $a \leq b$ .

If moreover  $A$  is a complete  $\widehat{\mathbb{U}}$ -module, then it is stackable: for  $a, b \in A$  such that  $v'(b) \leq u'(a)$  for all  $u', v' \in \mathbb{U}$ , with the same  $s, u, v, w$  as above, letting  $c_s := w(a) \wedge v^\times(b)$ , we have  $u(c_s) = u(w(a)) \wedge u(v^\times(b)) = u(w(a))$  which is within distance  $s$  of  $a$  since

$1_{\mathbb{U}} \leq u \circ w \leq (-) \dot{+} s$ , and  $v(c_s) = v(w(a)) \wedge v(v^\times(b)) = v(v^\times(b)) = b$ . In particular, by unstackability (using Lemma 4.15 and uniform continuity of  $u$ ), the  $c_s$  form a Cauchy net as  $s \searrow 0$ , hence converge to some  $c$  such that  $u(c) = a$  and  $v(c) = b$ . Thus the forgetful functor restricts to the claimed subcategories.

The forgetful functor is full on Archimedean  $\widehat{\mathbb{U}}$ -modules: the action by  $w \in \widehat{\mathbb{U}} \setminus \mathbb{U}$  can be recovered from the  $\mathbb{U}$ -action, since  $w(a) = \top$  for  $w(0) = 1$ , while for  $0 < w(0) < 1$ , by unstackability,  $w(a)$  is the unique element such that  $u(w(a)) = \top$  and  $v(w(a)) = (v \circ w)(a)$  where  $u, v$  are as above for  $r := w(0)$ . Thus  $\mathbb{U}$ -equivariance implies  $\widehat{\mathbb{U}}$ -equivariance.

Conversely, in a complete stackable  $\mathbb{U}$ -meet-semilattice  $A$ , we may extend the  $\mathbb{U}$ -action to a  $\widehat{\mathbb{U}}$ -action by defining  $w(a)$  for  $0 < w(0) < 1$  to be the unique element as above.

The  $\mathbb{U}$ -action on an Archimedean stackable  $\mathbb{U}$ -poset  $A$  preserves binary meets in  $\mathbb{U}$ : for piecewise linear  $u, v \in \mathbb{U}$ , we may show  $(u \wedge v)(a) = u(a) \wedge v(a)$  by unstacking over a finite partition of  $[0, 1]$  on each piece of which  $u, v$  are comparable; for arbitrary  $u, v$ , take piecewise linear approximations.

Finally, on a complete stackable  $\mathbb{U}$ -meet-semilattice, the extended  $\widehat{\mathbb{U}}$ -action from above also preserves binary meets in  $\widehat{\mathbb{U}}$ , by a routine unstacking over  $0 < w(0) < 1$ . ■

5.9. COROLLARY. [of Corollary 5.6 and Proposition 5.8] *Hom into  $\mathbb{I}$  yields a dual equivalence of categories between continuous lattices and complete  $\widehat{\mathbb{U}}$ -modules.* ■

We end by noting that we currently do not know whether complete  $\widehat{\mathbb{U}}$ -modules can be equationally axiomatized, perhaps along the lines of [Abb19], thereby showing that  $\text{CtsLat}^{\text{op}}$  is a variety.

## References

- [Abb19] M. Abbadini, *The dual of compact ordered spaces is a variety*, Theory Appl. Categ. **34** (2019), no. 44, 1401–1439.
- [ABLR02] J. Adámek, F. Borceux, S. Lack, and J. Rosický, *A classification of accessible categories*, J. Pure Appl. Algebra **175** (2002), no. 1–3, 7–30.
- [AK88] M. H. Albert and G. M. Kelly, *The closure of a class of colimits*, J. Pure Appl. Algebra **51** (1988), no. 1–2, 1–17.
- [BE83] H.-J. Bandelt and M. Erné, *The category of  $Z$ -continuous posets*, J. Pure Appl. Algebra **30** (1983), no. 3, 219–226.
- [G<sup>+</sup>03] G. Gierz, K. H. Hofmann, K. Keimel, J. D. Lawson, M. Mislove, and D. S. Scott, *Continuous lattices and domains*, Encyclopedia of Mathematics and its Applications, vol. 93, Cambridge University Press, 2003.
- [HNN18] D. Hofmann, P. Neves, and P. Nora, *Generating the algebraic theory of  $C(X)$ : the case of partially ordered compact spaces*, Theory Appl. Categ. **33** (2018), no. 12, 276–295.

- [HMS74] K. H. Hofmann, M. Mislove, and A. Stralka, *The Pontryagin duality of compact 0-dimensional semilattices and its applications*, Lecture Notes in Mathematics, vol. 396. Springer-Verlag, Berlin-New York, 1974.
- [Joh82] P. T. Johnstone, *Stone spaces*, Cambridge Studies in Advanced Mathematics, vol. 3, Cambridge University Press, Cambridge, England, 1982.
- [KS05] G. M. Kelly and V. Schmitt, *Notes on enriched categories with colimits of some class*, Theory Appl. Categ. **14** (2005), no. 17, 399–423.
- [Kün09] H.-P. A. Künzi, *An introduction to quasi-uniform spaces*, in: *Beyond topology*, 239–304, Contemp. Math. vol. 486, Amer. Math. Soc., Providence, RI, 2009.
- [MR17] V. Marra and L. Reggio, *Stone duality above dimension zero: axiomatising the algebraic theory of  $C(X)$* , Adv. Math. **307** (2017), 253–287.
- [WWT78] J. B. Wright, E. G. Wagner, and J. W. Thatcher, *A uniform approach to inductive posets and inductive closure*, Theoret. Comput. Sci. **7** (1978), no. 1, 57–77.
- [Xu95] X. Xu, *Construction of homomorphisms of  $M$ -continuous lattices*, Trans. Amer. Math. Soc. **347** (1995), no. 8, 3167–3175.

*Department of Mathematics*  
*University of Michigan*  
*Ann Arbor, MI 48109, USA*  
Email: ruiyuan@umich.edu

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

**SUBSCRIPTION INFORMATION** Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to [tac@mta.ca](mailto:tac@mta.ca) including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

**INFORMATION FOR AUTHORS** L<sup>A</sup>T<sub>E</sub>X<sub>2</sub>ε is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

**MANAGING EDITOR.** Geoff Cruttwell, Mount Allison University: [gcruttwell@mta.ca](mailto:gcruttwell@mta.ca)

**T<sub>E</sub>XNICAL EDITOR.** Michael Barr, McGill University: [michael.barr@mcgill.ca](mailto:michael.barr@mcgill.ca)

**ASSISTANT T<sub>E</sub>X EDITOR.** Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: [gavin\\_seal@fastmail.fm](mailto:gavin_seal@fastmail.fm)

**TRANSMITTING EDITORS.**

Clemens Berger, Université de Nice-Sophia Antipolis: [cberger@math.unice.fr](mailto:cberger@math.unice.fr)

Julie Bergner, University of Virginia: [jeb2md@virginia.edu](mailto:jeb2md@virginia.edu)

Richard Blute, Université d' Ottawa: [rblute@uottawa.ca](mailto:rblute@uottawa.ca)

John Bourke, Masaryk University: [bourkej@math.muni.cz](mailto:bourkej@math.muni.cz)

Maria Manuel Clementino, Universidade de Coimbra: [mmc@mat.uc.pt](mailto:mmc@mat.uc.pt)

Valeria de Paiva, Nuance Communications Inc: [valeria.depaiva@gmail.com](mailto:valeria.depaiva@gmail.com)

Richard Garner, Macquarie University: [richard.garner@mq.edu.au](mailto:richard.garner@mq.edu.au)

Ezra Getzler, Northwestern University: [getzler@northwestern.edu](mailto:getzler@northwestern.edu)

Rune Haugseng, Norwegian University of Science and Technology: [rune.haug seng@ntnu.no](mailto:rune.haug seng@ntnu.no)

Dirk Hofmann, Universidade de Aveiro: [dirk@ua.pt](mailto:dirk@ua.pt)

Joachim Kock, Universitat Autònoma de Barcelona: [Joachim.Kock@uab.cat](mailto:Joachim.Kock@uab.cat)

Stephen Lack, Macquarie University: [steve.lack@mq.edu.au](mailto:steve.lack@mq.edu.au)

Tom Leinster, University of Edinburgh: [Tom.Leinster@ed.ac.uk](mailto:Tom.Leinster@ed.ac.uk)

Sandra Mantovani, Università degli Studi di Milano: [sandra.mantovani@unimi.it](mailto:sandra.mantovani@unimi.it)

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: [matias.menni@gmail.com](mailto:matias.menni@gmail.com)

Giuseppe Metere, Università degli Studi di Palermo: [giuseppe.metere@unipa.it](mailto:giuseppe.metere@unipa.it)

Kate Ponto, University of Kentucky: [kate.ponto@uky.edu](mailto:kate.ponto@uky.edu)

Robert Rosebrugh, Mount Allison University: [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca)

Jiri Rosický, Masaryk University: [rosicky@math.muni.cz](mailto:rosicky@math.muni.cz)

Giuseppe Rosolini, Università di Genova: [rosolini@unige.it](mailto:rosolini@unige.it)

Michael Shulman, University of San Diego: [shulman@sandiego.edu](mailto:shulman@sandiego.edu)

Alex Simpson, University of Ljubljana: [Alex.Simpson@fmf.uni-lj.si](mailto:Alex.Simpson@fmf.uni-lj.si)

James Stasheff, University of North Carolina: [jds@math.upenn.edu](mailto:jds@math.upenn.edu)

Tim Van der Linden, Université catholique de Louvain: [tim.vanderlinden@uclouvain.be](mailto:tim.vanderlinden@uclouvain.be)

Christina Vasilakopoulou, National Technical University of Athens: [cvasilak@math.ntua.gr](mailto:cvasilak@math.ntua.gr)