

THE CATEGORY OF NECKLACES IS REEDY MONOIDAL

VIOLETA BORGES MARQUES AND ARNE MERTENS

ABSTRACT. In the first part of this note we further the study of the interactions between Reedy and monoidal structures on a small category, building upon the work of Barwick. We define a *Reedy monoidal category* as a Reedy category \mathcal{R} which is monoidal such that for all symmetric monoidal model categories \mathbf{A} , the category $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})_{\text{Reedy}}$ is monoidal model when equipped with the Day convolution. In the second part, we study the category \mathcal{Nec} of necklaces, as defined by Baues and Dugger-Spivak. Making use of a combinatorial description present in Grady-Pavlov and Lowen-Mertens, we streamline some proofs from the literature, and finally show that \mathcal{Nec} is simple Reedy monoidal.

1. Introduction

Reedy categories were first defined by Kan in unpublished notes and provide an abstract setting to generalize [24, Lemma 1.2], where simplicial objects are inductively defined through the factorization of the canonical maps $\text{sk}_n X \rightarrow \text{cosk}_n X$. This inductive construction allows one to define a model structure, the Reedy model structure, on functor categories $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})$. As opposed to the projective and injective model structure [20], the existence result for the Reedy model structure relies on imposing strong conditions on the small category \mathcal{R} rather than on the model category \mathbf{A} . The Reedy model structure also has the advantage that both (trivial) fibrations and cofibrations are explicitly prescribed.

Necklaces first appeared in [5] (under the name “cellular strings”) in the study of the bar and cobar construction and their relation to loop spaces. The terminology “necklace” was introduced and popularized by Dugger and Spivak, in [12][11] to provide several homotopically equivalent models of the mapping spaces of a quasi-category. Since then, necklaces have found further applications in work by Rivera and others, in particular in the construction of models of path spaces [23][27]; an adaptation of necklaces, called closed necklaces, was used to construct models of free loop spaces in [25] and [26]. In [22] a strengthening of Baues’ original results was presented. Several generalizations that use necklaces followed, for example, for dendroidal ∞ -operads in [6], for cubical quasi-categories in [9], for cartesian enriched quasi-categories in [13] and not necessarily cartesian

The authors would like to thank Wendy Lowen for her helpful discussions and support during the writing of this paper. They are also grateful to Clemens Berger for drawing attention to the reference [5], and to Dmitry Kaledin for pointing out the work of Manuel Rivera.

Received by the editors 2024-01-08 and, in final form, 2024-01-17.

Transmitted by Julie Bergner. Published on 2024-01-23.

2020 Mathematics Subject Classification: 18M05, 18N40 (Primary), 05E45 (Secondary).

Key words and phrases: Reedy category, necklaces, monoidal model category.

© Violeta Borges Marques and Arne Mertens, 2024. Permission to copy for private use granted.

enriched in [19][21]. The category of necklaces has also been employed to construct a Segalification functor, providing a Segal space generated by a simplicial space, in [7], to treat concurrency problems in [8] and to study (fully) extended functorial field theories (FFT) in [15].

In the present paper, we consider the model category $\text{Fun}(\mathcal{R}^{op}, \mathbf{A})_{\text{Reedy}}$ for a Reedy category \mathcal{R} and a symmetric monoidal model category \mathbf{A} . This category is naturally endowed with the pointwise monoidal structure and the question of whether this makes the Reedy model structure monoidal model is treated in [14]. On the other hand, if \mathcal{R} is itself monoidal, we can also consider the Day convolution product [10] on $\text{Fun}(\mathcal{R}^{op}, \mathbf{A})$. We define \mathcal{R} to be *Reedy monoidal* (Definition 2.22) if $\text{Fun}(\mathcal{R}^{op}, \mathbf{A})_{\text{Reedy}}$ is monoidal model with respect to the Day convolution for every symmetric monoidal model category \mathbf{A} . Building on general model categorical results by Barwick [4], we also provide simple combinatorial conditions to ensure that a category that is both Reedy and monoidal is indeed Reedy monoidal (Theorem 2.23).

Subsequently, we construct a Reedy structure on the category \mathcal{Nec} of necklaces (Theorem 3.23). The main ingredients of this structure are (at least implicitly) present in [27][28], but to the best of our knowledge the full Reedy (monoidal) structure has not been presented explicitly in the literature yet. For this, we make use of a combinatorial description of \mathcal{Nec} put forward in [15][19] which also makes some existing results easier to prove. Finally, we show that \mathcal{Nec} is a simple Reedy monoidal category (Theorem 3.25) so that $\text{Fun}(\mathcal{Nec}^{op}, \mathbf{A})_{\text{Reedy}}$ is always monoidal model when equipped with the Day convolution.

1.1. MOTIVATION. In [29], Simpson developed a theory of Segal categories enriched in a cartesian model category. Given an appropriate cartesian model category \mathcal{M} , a model structure on the category of unital \mathcal{M} -precategories is established, where the weak equivalences are the global weak equivalences and the fibrant objects satisfy the Segal condition (Theorems IV.19.2.1 and IV.19.4.1). The results of the present paper are motivated by an ongoing project to construct a model for Segal categories enriched in non-cartesian monoidal model categories \mathbf{A} as well. A main example of interest is the projective model structure on chain complexes $\mathbf{A} = \text{Ch}(k)$ over a field k . The unital \mathbf{A} -precategories in this case are no longer given by simplicial objects $S\mathbf{A}$, but by *templcial objects* $S_{\otimes}\mathbf{A}$. These are certain strictly unital colax monoidal functors which were introduced in [19] as replacements for simplicial objects in the non-cartesian context, and inspired by earlier work of Leinster [18] and Bacard [1][3]. In the non-cartesian setting we propose then the following definition.

1.2. DEFINITION. Let \mathbf{A} be a monoidal model category. A *Segal \mathbf{A} -category* is a templcial \mathbf{A} -object $(X, S) \in S_{\otimes}\mathbf{A}$ such that the comultiplication maps $\mu_{i,j} : X_{i+j} \xrightarrow{\sim} X_i \otimes_S X_j$ are weak equivalences in \mathbf{A} for all $i, j \geq 0$.

A first step in this direction is the definition of a Reedy model structure on $S_{\otimes}\mathbf{A}$ as

follows. It is shown in [19] that there is an adjunction with fully faithful left adjoint

$$S_{\otimes} \mathbf{A} \begin{array}{c} \xrightarrow{(-)^{nec}} \\ \perp \\ \xleftarrow{(-)^{temp}} \end{array} \text{Fun}(\mathcal{N}ec^{op}, \mathbf{A})\text{-Cat}$$

In future work, we will show that under suitable yet moderate conditions, the monoidal model structure $\text{Fun}(\mathcal{N}ec^{op}, \mathbf{A})_{\text{Reedy}}$ induces a model structure on $\text{Fun}(\mathcal{N}ec^{op}, \mathbf{A})\text{-Cat}_S$, the category of necklicial categories with fixed object set S . In case \mathbf{A} is cartesian or $\mathbf{A} = \text{Ch}(k)$, the former can be further transferred to $S_{\otimes} \mathbf{A}_S$. Moreover, in case \mathbf{A} is cartesian, we recover the classical Reedy structure under the equivalence $S_{\otimes} \mathbf{A}_S \cong \text{Fun}(\Delta_S^{op}/S, \mathbf{A})$ [29, Proposition III.12.3.1].

2. Reedy monoidal categories

We start this section by recalling some concepts of the theory of Reedy categories following [4, §3]. While *loc. cit.* introduces these concepts with a model category theoretical approach, we opt to adopt the equivalent combinatorial characterizations as our primary definitions. For a treatment of the model category theory aspects, we defer to the latter part of this section.

In §2.1, we show that any morphism of Reedy categories that restricts to a discrete fibration between direct subcategories, is a right fibration (Proposition 2.10). As a consequence, the monoidal product of any direct divisible Reedy category in the sense of [3] is a right fibration. We conclude in §2.2 by introducing Reedy monoidal categories (Definition 2.22), along with sufficient conditions for checking this property (Theorem 2.23).

2.1. REEDY CATEGORIES AND RIGHT FIBRATIONS. For standard treatments of the theory of Reedy categories, we refer to [17] [16]. Let us fix a small category \mathcal{R} .

2.2. DEFINITION. [17, Definition 5.2.1] A *Reedy structure* on \mathcal{R} is a pair of wide *inverse* and *direct* subcategories $(\mathcal{R}^{\leftarrow}, \mathcal{R}^{\rightarrow})$ and a *degree function* $\text{deg} : \text{Ob}(\mathcal{R}) \rightarrow \lambda$ with λ an ordinal, such that the following conditions are satisfied:

1. Every morphism f in \mathcal{R} factors uniquely as $f = f^{\rightarrow} \circ f^{\leftarrow}$ with $f^{\rightarrow} \in \mathcal{R}^{\rightarrow}$ and $f^{\leftarrow} \in \mathcal{R}^{\leftarrow}$.
2. Every non-identity morphism in \mathcal{R}^{\leftarrow} lowers the degree and every non-identity morphism in $\mathcal{R}^{\rightarrow}$ raises the degree.

2.3. EXAMPLES.

1. The terminal category $\{*\}$ has a trivial Reedy structure.
2. The simplex category Δ is Reedy with the degree function $d : \text{Ob}(\Delta) \rightarrow \mathbb{N} : [n] \mapsto n$, and Δ^{\leftarrow} and Δ^{\rightarrow} containing the surjective and injective order morphisms respectively.

3. Suppose that $(\mathcal{R}, \mathcal{R}^{\leftarrow}, \mathcal{R}^{\rightarrow})$ and $(\mathcal{S}, \mathcal{S}^{\leftarrow}, \mathcal{S}^{\rightarrow})$ are Reedy categories, then so are $(\mathcal{R}^{\text{op}}, (\mathcal{R}^{\rightarrow})^{\text{op}}, (\mathcal{R}^{\leftarrow})^{\text{op}})$ and $(\mathcal{R} \times \mathcal{S}, \mathcal{R}^{\leftarrow} \times \mathcal{S}^{\leftarrow}, \mathcal{R}^{\rightarrow} \times \mathcal{S}^{\rightarrow})$.

2.4. DEFINITION. [17, Def 5.1.2] Let \mathcal{R} be a Reedy category and $\alpha \in \mathcal{R}$. The *latching category* at α is

$$\partial(\mathcal{R}^{\rightarrow}/\alpha) = \{f \in \mathcal{R}^{\rightarrow}/\alpha \mid f \neq \text{id}\}$$

and the *matching category* at α is

$$\partial(\alpha/\mathcal{R}^{\leftarrow}) = \{f \in \alpha/\mathcal{R}^{\leftarrow} \mid f \neq \text{id}\}.$$

2.5. REMARK. Note that there is a canonical isomorphism between the latching (resp. matching) category of \mathcal{R} at α and the matching (resp. latching) category of \mathcal{R}^{op} at α .

Now let us turn to functors between Reedy categories.

2.6. DEFINITION. [4, Definition 3.16.1] Let \mathcal{R} and \mathcal{S} be two Reedy categories. A functor $\mathcal{R} \xrightarrow{F} \mathcal{S}$ is a *morphism of Reedy categories* if $F(\mathcal{R}^{\rightarrow}) \subseteq \mathcal{S}^{\rightarrow}$ and $F(\mathcal{R}^{\leftarrow}) \subseteq \mathcal{S}^{\leftarrow}$.

2.7. DEFINITION. [4, Theorem 3.22] A morphism of Reedy categories $\mathcal{R} \xrightarrow{F} \mathcal{S}$ is called

- a *right fibration* if for every $\alpha \in \mathcal{R}$ and every $\beta \xrightarrow{f} F(\alpha) \in \beta/F$ the category $\partial((\beta/F)^{\rightarrow}/f)$ is empty or connected. The category \mathcal{R} is *right fibrant* if the functor $\mathcal{R} \rightarrow \{*\}$ is a right fibration.
- a *left fibration* if for every $\alpha \in \mathcal{R}$ and every $F(\alpha) \xrightarrow{f} \beta \in F/\beta$ the category $\partial(f/(F/\beta)^{\leftarrow})$ is empty or connected. The category \mathcal{R} is *left fibrant* if the functor $\mathcal{R} \rightarrow \{*\}$ is a left fibration.

2.8. REMARK. In the previous definition we make implicit use of the Reedy structure on β/F and F/β given in [4, Lemma 3.10].

2.9. REMARK. Observe that \mathcal{R} is left fibrant if and only if for any $\alpha \in \mathcal{R}$, the matching categories $\partial(\alpha/\mathcal{R}^{\leftarrow})$ are empty and connected. Note that if \mathcal{R} has a terminal object \star such that for all $\alpha \in \mathcal{R}$, the unique map $\alpha \rightarrow \star$ belongs to \mathcal{R}^{\leftarrow} , then \mathcal{R} is left fibrant.

The conditions of the above definition are purely combinatorial and very tractable. As we are going to make use of it later, let us make the latching category involved in the definition of right fibration more explicit. For fixed $\alpha \in \mathcal{R}$ and $f : \beta \rightarrow F(\alpha)$, its objects are triples $(\alpha' \in \mathcal{R}, f' : \beta \rightarrow F(\alpha'), g : \alpha' \rightarrow \alpha)$ with $g \neq \text{id}$ in $\mathcal{R}^{\rightarrow}$ such that $f = F(g) \circ f'$ which we represent by the diagram

$$\begin{array}{ccc} & & F(\alpha') \\ & \nearrow^{f'} & \searrow^{F(g)} \\ \beta & \xrightarrow{f} & F(\alpha) \end{array}$$

and the morphisms $(\alpha', f', g') \rightarrow (\alpha'', f'', g'')$ are morphisms $h : \alpha' \rightarrow \alpha''$ in \mathcal{R}^\rightarrow such that $F(h) \circ f' = f''$ and $g'' \circ h = g'$.

Recall that a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between categories is called a *discrete fibration* if for all $C \in \mathcal{C}$ and all $f : D \rightarrow F(C)$ in \mathcal{D} , there is a unique $g : \bar{C} \rightarrow C$ in \mathcal{C} such that $F(g) = f$.

2.10. PROPOSITION. *Let $F : \mathcal{R} \rightarrow \mathcal{S}$ be a morphism of Reedy categories. If the restriction $F^\rightarrow : \mathcal{R}^\rightarrow \rightarrow \mathcal{S}^\rightarrow$ is a discrete fibration of categories, then $F : \mathcal{R} \rightarrow \mathcal{S}$ is a right fibration of Reedy categories.*

PROOF. Let us fix $\alpha \in \mathcal{R}$ and $f : \beta \rightarrow F(\alpha)$ in \mathcal{S} . Since F^\rightarrow is a discrete fibration and \mathcal{S} is a Reedy category, we have a unique factorization of f :

$$\begin{array}{ccc} & & F(\bar{\alpha}) \\ & \nearrow^{f^\leftarrow} & \swarrow^{F(g^\rightarrow)} \\ \beta & \xrightarrow{f} & F(\alpha) \end{array} \quad (1)$$

with f^\leftarrow in \mathcal{S}^\leftarrow and $g^\rightarrow : \bar{\alpha} \rightarrow \alpha$ in \mathcal{R}^\rightarrow . We show that this is the initial object of the undercategory $((\beta/F)^\rightarrow/f)$. Consider an arbitrary object (represented in a slightly different manner):

$$\begin{array}{ccc} \beta & \xrightarrow{f} & F(\alpha) \\ & \searrow^{f'} & \nearrow^{F(g')} \\ & & F(\alpha') \end{array} \quad (2)$$

We take the $(\mathcal{S}^\leftarrow, \mathcal{S}^\rightarrow)$ -factorization of f' :

$$\begin{array}{ccc} & & \bar{\alpha}' \\ & \nearrow^{f'^\leftarrow} & \downarrow^{f'^\rightarrow} \\ \beta & \xrightarrow{f'} & F(\alpha) \\ & \searrow^{f'} & \nearrow^{F(g')} \\ & & F(\alpha') \end{array}$$

but now by uniqueness of factorization we know $f'^\leftarrow = f^\leftarrow$, $\bar{\alpha}' = F(\bar{\alpha})$ and $F(g') \circ f'^\rightarrow = F(g^\rightarrow)$. Again since F^\rightarrow is a discrete fibration, we can write $f'^\rightarrow = g'^\rightarrow$ for some $g'^\rightarrow : \bar{\alpha} \rightarrow \alpha'$ in \mathcal{R}^\rightarrow with $g' \circ g'^\rightarrow = f^\rightarrow$. Thus we obtain the uniquely defined map

$$\begin{array}{ccc} & & F(\bar{\alpha}) \\ & \nearrow^{f'^\leftarrow} & \swarrow^{F(f'^\rightarrow)} \\ \beta & \xrightarrow{f'} & F(\alpha) \\ & \searrow^{f'} & \nearrow^{F(g)} \\ & & F(\alpha') \end{array}$$

Finally, if $f^\rightarrow \neq \text{id}$, the initial object (1) lies in the latching category $\partial((\beta/F)^\rightarrow/f)$, whereby it is connected. If $f^\rightarrow = \text{id}$, then it follows for an arbitrary object (2) that $F(g) = \text{id}$ as well. Again since F is a discrete fibration, this implies that $g = \text{id}$. Thus (2) does not belong to $\partial((\beta/F)^\rightarrow/f)$, whereby it is empty. \blacksquare

Let us end this subsection by establishing some terminology on how monoidal and Reedy structures interact, inspired by [3].

2.11. DEFINITION. Let \mathcal{R} be a Reedy category with a monoidal structure (\vee, I) .

1. The monoidal structure (\vee, I) is *compatible* if the functor \vee is a morphism of Reedy categories $\mathcal{R} \times \mathcal{R} \rightarrow \mathcal{R}$.
2. The category \mathcal{R} is *direct divisible with respect to* \vee if \vee^\rightarrow is a discrete fibration, i.e., for any map $f : X \rightarrow Y_1 \vee Y_2 \in \mathcal{R}^\rightarrow$, there exist a unique pair of maps in \mathcal{R}^\rightarrow $f_1 : X_1 \rightarrow Y_1$ and $f_2 : X_2 \rightarrow Y_2$ such that $f = f_1 \vee f_2$.
3. (\mathcal{R}, \vee, I) is *simple* if for any $\alpha, \beta \in \mathcal{R}$, $\deg(\alpha \vee \beta) = \deg \alpha + \deg \beta$.

2.12. EXAMPLE. Consider the subcategory $\Delta_f \subseteq \Delta$ containing all morphisms $f : [m] \rightarrow [n]$ such that $f(0) = 0$ and $f(m) = n$. It is monoidal with $[n] + [m] = [n + m]$ and Reedy with the restriction of the Reedy structure of Δ . The monoidal structure is compatible, making $(\Delta_f, +, [0])$ is simple. However, Δ_f is not direct divisible with respect to $+$.

2.13. REMARK. What we call a Reedy category with compatible monoidal structure is exactly a one-object *locally Reedy 2-category* from [2], and similarly for a simple Reedy category. Bacard defines direct divisibility only for simple Reedy categories, by requiring \vee^\rightarrow to be a Grothendieck fibration. However in the simple case, this is equivalent to \vee^\rightarrow being a discrete fibration.

2.14. REEDY MONOIDAL MODEL STRUCTURES. In this subsection we highlight the usefulness and inspiration of the definitions of the previous subsection in the theory of model categories. Standard references for model category theory are [17] and [16].

2.15. DEFINITION. [17, Def 5.2.2] Let \mathcal{R} be a Reedy category, \mathbf{A} a bicomplete category, $\alpha \in \mathcal{R}$ and $X \in \text{Fun}(\mathcal{R}, \mathbf{A})$. The *latching object* of X at α is

$$L_\alpha X = \text{colim}_{\beta \rightarrow \alpha \in \partial(\mathcal{R}^\rightarrow / \alpha)} X(\beta)$$

and the *matching object* of X at α is

$$M_\alpha X = \lim_{\alpha \rightarrow \beta \in \partial(\alpha / \mathcal{R}^\leftarrow)} X(\beta)$$

Firstly we present the very well known fact that for any model category \mathbf{A} , the functor category $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})$ can be endowed with a model structure. Note that the latching and matching objects and categories are now referring to \mathcal{R}^{op} (see Example 2.3.3).

2.16. THEOREM. [17, Theorem 5.2.5] *Let \mathcal{R} be a Reedy category and \mathbf{A} a model category. Then there exists a model structure on $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})$, denoted $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})_{\text{Reedy}}$, such that*

1. *a map $X \xrightarrow{f} Y$ is a weak equivalence if and only if $X_\alpha \xrightarrow{f_\alpha} Y_\alpha$ is a weak equivalence for all $\alpha \in \mathcal{R}$.*
2. *a map $X \xrightarrow{f} Y$ is a (trivial) cofibration if and only if $X_\alpha \amalg_{L_\alpha X} L_\alpha Y \rightarrow Y_\alpha$ is a (trivial) cofibration for all $\alpha \in \mathcal{R}$.*
3. *a map $X \xrightarrow{f} Y$ is a (trivial) fibration if and only if $X_\alpha \rightarrow M_\alpha X \times_{M_\alpha Y} Y_\alpha$ is a (trivial) fibration for all $\alpha \in \mathcal{R}$.*

2.17. PROPOSITION. [4, Definition 3.16.3] *A morphism of Reedy categories $\mathcal{R} \xrightarrow{F} \mathcal{S}$ is a right fibration if and only if for any model category \mathbf{A} the adjunction*

$$\text{Fun}(\mathcal{S}^{\text{op}}, \mathbf{A})_{\text{Reedy}} \xleftarrow[F^*]{F_!} \text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})_{\text{Reedy}}$$

is a Quillen adjunction. It is a left fibration if and only if for any model category \mathbf{A} the adjunction

$$\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})_{\text{Reedy}} \xleftarrow[F_*]{F^*} \text{Fun}(\mathcal{S}^{\text{op}}, \mathbf{A})_{\text{Reedy}}$$

is a Quillen adjunction.

2.18. COROLLARY. *A Reedy category \mathcal{R} is left fibrant if and only if \mathcal{R}^{op} has cofibrant constants, that is for any cofibrant $X \in \mathbf{A}$, the constant functor ΔX is cofibrant in $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})_{\text{Reedy}}$.*

PROOF. Immediate from Proposition 2.17. ■

Our particular interest in right fibrations comes from the following theorem.

2.19. THEOREM. [4, Corollary 3.50] *Suppose \mathbf{A} is a symmetric monoidal model category and \mathcal{R} is a Reedy category equipped with a monoidal product*

$$\vee : \mathcal{R} \times \mathcal{R} \rightarrow \mathcal{R}$$

that is a right fibration of Reedy categories. Then the Day convolution product \otimes_{Day} is a Quillen bifunctor with respect to the Reedy model structure.

Note that this theorem only concerns the monoidal product. However, in a monoidal model category also the monoidal unit is required to be homotopically well-behaved.

2.20. DEFINITION. [17, Definition 4.2.6] *A monoidal model category is a monoidal closed category $(\mathbf{A}, \otimes, \mathbb{I})$ with a model structure on \mathbf{A} , such that the following conditions hold.*

1. *The monoidal product $\otimes : \mathbf{A} \times \mathbf{A} \rightarrow \mathbf{A}$ is a Quillen bifunctor.*
2. *Let $q : Q\mathbb{I} \xrightarrow{\sim} \mathbb{I}$ be the cofibrant replacement of the unit \mathbb{I} . Then the natural maps $q \otimes X : Q\mathbb{I} \otimes X \rightarrow \mathbb{I} \otimes X$ and $X \otimes q : X \otimes Q\mathbb{I} \rightarrow X \otimes \mathbb{I}$ are weak equivalences for all cofibrant X .*

2.21. **REMARK.** Observe that by standard model category theoretical arguments, condition (b) is equivalent to the statement: for *any* cofibrant replacement $\bar{q} : \bar{\mathbb{I}} \xrightarrow{\sim} \mathbb{I}$ of the monoidal unit \mathbb{I} and any cofibrant object X , $X \otimes \bar{q}$ and $\bar{q} \otimes X$ are weak equivalences.

In the spirit of Proposition 2.17 and [16, Definition 15.10.1], we establish the following definition.

2.22. **DEFINITION.** Let \mathcal{R} be a Reedy category with a compatible monoidal structure (\vee, I) . The category \mathcal{R} is *Reedy monoidal* if for any symmetric monoidal model category \mathbf{A} , $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})_{\text{Reedy}}$ is a monoidal model category when equipped with Day convolution.

2.23. **THEOREM.** *Let \mathcal{R} be a left fibrant Reedy category with a compatible monoidal structure (\vee, I) . If I is a terminal object and \mathcal{R} is direct divisible with respect to \vee , then \mathcal{R} is Reedy monoidal.*

PROOF. By Proposition 2.10 and Theorem 2.19 we obtain condition (a) of Definition 2.20. Let $(\mathbf{A}, \otimes, \mathbb{I})$ be a symmetric monoidal model category. Since I is terminal, the Day convolution unit of $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})$ is the constant diagram $\Delta\mathbb{I}$. Let $q : Q\mathbb{I} \rightarrow \mathbb{I}$ be the cofibrant replacement of \mathbb{I} in \mathbf{A} . Then by Corollary 2.18, $\Delta(Q\mathbb{I})$ is cofibrant and moreover $\Delta(Q\mathbb{I}) \rightarrow \Delta\mathbb{I}$ is a levelwise weak equivalence, and thus a weak equivalence in $\text{Fun}(\mathcal{R}^{\text{op}}, \mathbf{A})_{\text{Reedy}}$. Finally, observe that $(X \otimes_{\text{Day}} \Delta q)_\alpha \cong X_\alpha \otimes q$ for all $\alpha \in \mathcal{R}$ and $X \in \mathbf{A}$ and thus the final result follows by the assumption that \mathbf{A} is monoidal model and by [17, Remark 5.1.7]. \blacksquare

3. The category of necklaces

Let us now turn our attention to the category \mathcal{Nec} of necklaces, as considered in [5][12]. In §3.1, we give a self-contained account of a number of relevant factorization classes of morphisms, making use of a combinatorial description of necklaces put forth in [15][19]. In §3.2, we go on to show that the (epi, mono)-factorization can be made into a Reedy monoidal structure (see Definition 2.22) with an appropriate degree function (Theorem 3.25). This makes use of the notion of dimension of a necklace, which plays a fundamental role in [28]. The interaction between the classes of epimorphisms and monomorphisms and the dimension, which constitutes an important part of our proof, is implicit in loc. cit as we explain in Remarks 3.14 and 3.21, and a decomposition of \mathcal{Nec} by degree is used in [27].

3.1. **A COMBINATORIAL DESCRIPTION OF NECKLACES.** Let us denote by $\text{SSet}_{\star, \star}$ the category of bipointed simplicial sets, i.e. the under category $\partial\Delta^1/\text{SSet}$. This is a monoidal category when equipped with the wedge product

$$(X, x_0, x_1) \vee (Y, y_0, y_1) = \left(X \coprod_{x_1=y_0} Y, x_0, y_1 \right)$$

and with the 0-simplex $(\Delta^0, 0, 0)$ as monoidal unit. We consider the standard n -simplices Δ^n as bipointed at 0 and n . To simplify notation, and as it will not bring about any confusion, we will simply denote $(\Delta^n, 0, n)$ by Δ^n .

3.2. DEFINITION. The *category of necklaces* \mathcal{Nec} is the full subcategory of $\mathbf{SSet}_{*,*}$ spanned by objects X of shape

$$X = \Delta^{n_1} \vee \dots \vee \Delta^{n_k}$$

with $k \geq 0$ and $n_i > 0$, called *necklaces*. The simplices Δ^{n_i} are the *beads* of the necklace X . If $k = 0$, the corresponding necklace is denoted Δ^0 . We call the number of beads k the *bead length* of X , denoted $\ell(X)$; and the sum $n_1 + \dots + n_k$ the *spine length* of X , denoted $\|X\|$.

Independently in [15, Proposition 3.4.2] and [19, Proposition 3.4], the following alternative combinatorial description of necklaces was given. Recall the category Δ_f from Example 2.12.

3.3. PROPOSITION. *The category of necklaces \mathcal{Nec} is equivalent to the category defined as follows:*

- *the objects are pairs (T, p) with $p \geq 0$ and $\{0, p\} \subseteq T \subseteq [p]$;*
- *the morphisms $(T, p) \rightarrow (S, q)$ are morphisms $f : [p] \rightarrow [q] \in \Delta_f$ such that $S \subseteq f(T)$;*
- *composition and identities are given as in Δ_f .*

Moreover, under this equivalence, the wedge \vee corresponds to

$$(T, p) \vee (S, q) = (T \cup (p + S), p + q)$$

where $p + S = \{p + s \mid s \in S\}$.

Under this equivalence of categories, we can identify a “combinatorial” necklace $(\{0 = t_0 < t_1 < \dots < t_k = p\}, p)$ with the “geometric” necklace $\Delta^{t_1-t_0} \vee \dots \vee \Delta^{t_k-t_{k-1}}$. Note that in particular, $(\{0 < n\}, n)$ corresponds to the n -simplex Δ^n , while $([p], p)$ corresponds to a sequence of edges, that we call *spines*. In the rest of this section we establish a correspondence between the combinatorial and geometric descriptions of several natural concepts.

3.4. DEFINITION. [19, Def. 3.5] Let $f : (T, p) \rightarrow (S, q)$ be a map of necklaces. The map f is *inert* if $p = q$ and $f = \text{id}_{[p]}$. It is *active* if $f(T) = S$.

Consider an arbitrary necklace map $f : (T, p) \rightarrow (S, q)$. Then it factors uniquely as an active map followed by an inert map:

$$(T, p) \xrightarrow{f^{\text{ac}}} (f(T), q) \xrightarrow{f^{\text{in}}} (S, q) \tag{3}$$

3.5. LEMMA. *A map of necklaces $X = (T, p) \xrightarrow{f} Y = (S, q)$ is inert if and only if it is the wedge $\iota_1 \vee \cdots \vee \iota_k$ of the inclusions*

$$\iota_i : \Delta^{n_i^1} \vee \cdots \vee \Delta^{n_i^{l_i}} \hookrightarrow \Delta^{n_i^1 + \cdots + n_i^{l_i}}$$

PROOF. By hypothesis $p = q$ so we are mainly concerned with the inclusion $S \subseteq T$. Denote $S = \{0 = s_0 < s_1 < \cdots < s_k = q\}$ and write

$$T = \{0 < \cdots < s_i = t_i^0 < t_i^1 < \cdots < t_i^{l_i} = s_{i+1} < \cdots < p\}$$

Then we can set $n_i^j = t_i^j - t_i^{j-1}$ and thus $n_i^1 + \cdots + n_i^{l_i} = s_i - s_{i-1}$. ■

3.6. LEMMA. *A map of necklaces $X = (T, p) \xrightarrow{f} Y = (S, q)$ is active if and only if it is the wedge $f_1 \vee \cdots \vee f_k$ of necklace maps $f_i : \Delta^{n_i} \rightarrow \Delta^{m_i}$ induced by morphisms $[n_i] \rightarrow [m_i]$ in Δ_f .*

PROOF. Simply write $n_i = t_i - t_{i-1}$ and $f_i = f|_{\{t_{i-1} < \cdots < t_i\}}$. ■

Again consider an arbitrary necklace map $f : (T, p) \rightarrow (S, q)$. We can factor $[p] \rightarrow [q]$ as a surjective map followed by an injective map, $[p] \xrightarrow{f_1} [r] \xrightarrow{f_2} [q]$. We thus get a new way of uniquely factoring necklace maps

$$(T, p) \xrightarrow{f_1} (f_1(T), r) \xrightarrow{f_2} (S, q) \quad (4)$$

as $f_1(T) \subseteq f(T)$ and $S \subseteq f(T) = f_2(f_1(T))$. We note that the first map is active and surjective on vertices and the second is injective on vertices.

3.7. LEMMA. *A map of necklaces $X = (T, p) \xrightarrow{f} Y = (S, q)$ is a monomorphism if and only if $[p] \rightarrow [q]$ is injective.*

PROOF. Under the identification provided by Proposition 3.3, $[p]$ and $[q]$ are the sets of vertices of X and Y respectively, and so the necessity is immediate.

We now show the sufficiency. Factor $f = f^{\text{in}} \circ f^{\text{ac}}$. Notice that an inert map is the wedge of inclusions, and thus a monomorphism. Now f^{ac} is the wedge of simplicial maps, which are completely determined by the values at their vertices. If $[p] \rightarrow [q]$ is injective, so will the restriction to the beads be. Thus f^{ac} is a wedge of monomorphisms and thus a monomorphism. This shows f is a monomorphism as desired. ■

3.8. LEMMA. *A map of necklaces $X = (T, p) \xrightarrow{f} Y = (S, q)$ is an epimorphism if and only if it is active and $[p] \rightarrow [q]$ is surjective.*

PROOF. Again factor f as $f = f^{\text{in}} \circ f^{\text{ac}}$ with f^{ac} active and f^{in} inert. As f is an epimorphism, so is f^{in} , but as inert maps are injective, $f^{\text{in}} = \text{id}$ and thus f is active. Again, as $[p] \rightarrow [q]$ is the map on vertices, it will necessarily be surjective.

Assume now that f is active and that $[p] \rightarrow [q]$ is surjective. We can write f as a wedge of epimorphisms between beads, and those are epimorphic. ■

Recall that as $\mathbb{S}\text{Set}_{*,*}$ is a topos, the (epi, mono) orthogonal factorization system through the image exists. Via the two lemmas above we deduce the following proposition.

3.9. PROPOSITION. *The factorization given in (4) is the restriction of the (epi, mono)-factorization of $\mathbb{S}\text{Set}_{*,*}$ to $\mathcal{N}ec$.*

As a corollary, we obtain a different proof the following result from [12].

3.10. COROLLARY. [12, Lemma 3.5(7)] *Let $X \xrightarrow{f} Y$ be a map of necklaces. Then the image of f is a necklace.*

	Combinatorial	Geometric
$f : X \rightarrow Y$	$f : (T, p) \rightarrow (S, q)$	$f : \Delta^{n_1} \vee \dots \vee \Delta^{n_k} \rightarrow \Delta^{m_1} \vee \dots \vee \Delta^{m_l}$
f is active	$S = f(T)$	$k = l, f = f_1 \vee \dots \vee f_k, f_i : \Delta^{n_i} \rightarrow \Delta^{m_i}$
f is inert	$p = q, [p] \xrightarrow{\text{id}} [q]$	$f = \iota_1 \vee \dots \vee \iota_l, m_i = n_{j_{i-1}+1} + \dots + n_{j_i}$ $\iota_i : \Delta^{n_{j_i+1}} \vee \dots \vee \Delta^{n_{j_{i+1}}} \hookrightarrow \Delta^{m_i}$
f is epi	$[p] \rightarrow [q]$ surjective, $S = f(T)$	f_n is surjective, $\forall n \geq 0$
f is mono	$[p] \rightarrow [q]$ injective	f_n is injective, $\forall n \geq 0$

Table 1: Combinatorial and geometric characterizations of the four factorization classes.

We now introduce a further decomposition of the epimorphisms, that will come in handy in the next section.

3.11. DEFINITION. Let $f = f_1 \vee \dots \vee f_k$ be an active surjective necklace map with $f_i : \Delta^{n_i} \rightarrow \Delta^{m_i}$. The map f is *bead reducing* if $m_i \geq 1$ for all $1 \leq i \leq k$ and *spine collapsing* if for all $1 \leq i \leq k$, $f_i = \text{id}_{\Delta^{n_i}}$ or $f_i : \Delta^1 \rightarrow \Delta^0$.

Observe that if $(T, p) \xrightarrow{f} (f(T), q)$ is an active surjective map with $T = \{0 = t_0 < \dots < t_k = p\}$, then we have for all $0 < i \leq k$

$$0 \leq f(t_i) - f(t_{i-1}) \leq t_i - t_{i-1}$$

3.12. LEMMA. *A map of necklaces $X = (T, p) \xrightarrow{f} (f(T), q)$ is bead reducing if for all $0 < i \leq k$, $0 < f(t_i) - f(t_{i-1})$.*

PROOF. Follows immediately as $m_i = f(t_i) - f(t_{i-1})$. ■

3.13. LEMMA. *A map of necklaces $X = (T, p) \xrightarrow{f} (f(T), q)$ is spine collapsing if and only if for all $0 < i \leq k$, $f(t_i) - f(t_{i-1}) = t_i - t_{i-1}$ or $t_i - t_{i-1} = 1$ and $f(t_i) - f(t_{i-1}) = 0$.*

PROOF. The first possibility corresponds to $f_i = \text{id}$ and the second to $f_i : \Delta^1 \rightarrow \Delta^0$. ■

It clearly follows from the previous lemmas that any active surjective necklace map $(T, p) \rightarrow (f(T), q)$ can be factored as a bead reducing map followed by a spine collapsing map. We thus obtain a (unique) factorization of any necklace map as follows

$$\bullet \xrightarrow{\text{bead reducing}} \bullet \xrightarrow{\text{spine collapsing}} \bullet \xrightarrow{\text{monomorphism}} \bullet \quad (5)$$

3.14. **REMARK.** In [28, Proposition 3.1], a finer factorization of necklace maps is described, in terms of maps of types (i), (ii) and (iii). It is readily seen that a map is a monomorphism (resp. bead reducing, resp. spine collapsing) precisely when it is a compositions of maps of type (i) (resp. (ii), resp. (iii)).

3.15. **THE REEDY MONOIDAL STRUCTURE ON NECKLACES.** In this section, in order to show that the (epi, mono)-factorization extends to a Reedy monoidal structure on \mathcal{Nec} , we make use of the following fundamental notion.

3.16. **DEFINITION.** [27, §2], [28, §4] Let $X \in \mathcal{Nec}$. Its *dimension* is

$$\dim(X) = \|X\| - \ell(X)$$

	Combinatorial	Geometric
X	(T, p)	$\Delta^{n_1} \vee \dots \vee \Delta^{n_k}$
$\ell(X)$	$ T - 1$	k
$\ X\ $	p	$n_1 + \dots + n_k$
$\dim X$	$p - T + 1$	$n_1 + \dots + n_k - k$

Table 2: Combinatorial and geometric characterizations of numerical invariants of necklaces.

3.17. **REMARK.** It is readily seen that for any $X, Y \in \mathcal{Nec}$, we have $\|X \vee Y\| = \|X\| + \|Y\|$ and $\ell(X \vee Y) = \ell(X) + \ell(Y)$ and consequently $\dim(X \vee Y) = \dim X + \dim Y$.

3.18. **LEMMA.** *Let $f : X \rightarrow Y$ be a non-identity monomorphism. Then $\dim X < \dim Y$.*

PROOF. Factor f as $f = f^{\text{in}} \circ f^{\text{ac}}$. As $f \neq \text{id}$ then either $f^{\text{ac}} \neq \text{id}$ or $f^{\text{in}} \neq \text{id}$. In the first case, $[p] \rightarrow [q]$ is a strict injection, thus $q > p$, and $|T| = |f(T)|$, so $\dim X < \dim Z$. In the second case, we know that $S \subsetneq f(T)$ so $|f(T)| > |S|$ and finally $\dim Z < \dim Y$. ■

3.19. **LEMMA.** *Let $f : X \rightarrow Y$ be a non-identity bead reducing map. Then $\dim X < \dim Y$.*

PROOF. Write $f : X = (T, p) \rightarrow Y = (f(T), q)$. We begin by noting that as $0 < f(t_i) - f(t_{i-1})$ for all i , then $|T| = |f(T)|$. As $f \neq \text{id}$, then $p > q$ and the result follows. ■

3.20. **LEMMA.** *Let $f : X \rightarrow Y$ be a non-identity spine collapsing necklace map. Then $\dim X = \dim Y$ and $\ell(X) > \ell(Y)$.*

PROOF. The equality follows from the additivity of dimension and the fact that $\dim \Delta^1 = 0$. As for the length, writing $f = f^1 \vee \dots \vee f^k$, and as $f \neq \text{id}$, there exists at least one $1 \leq i \leq k$ such that $f^i : [1] \rightarrow [0]$ and thus $\ell(X) > \ell(Y)$. ■

3.21. **REMARK.** In the light of Remark 3.14, Lemmas 3.18, 3.19 and 3.20 can also be deduced from the proof of [28, Prop.4.2].

$f : X \xrightarrow{\neq} Y$	spine length	bead length	dimension
bead reducing	$\ X\ > \ Y\ $	$\ell(X) = \ell(Y)$	$\dim(X) > \dim(Y)$
spine collapsing	$\ X\ > \ Y\ $	$\ell(X) > \ell(Y)$	$\dim(X) = \dim(Y)$
active injective	$\ X\ < \ Y\ $	$\ell(X) = \ell(Y)$	$\dim(X) < \dim(Y)$
inert	$\ X\ = \ Y\ $	$\ell(X) > \ell(Y)$	$\dim(X) < \dim(Y)$

Table 3: Interaction of spine length, bead length and dimension with respect to several classes of necklace maps.

3.22. **DEFINITION.** The *degree function* is $\deg : \text{Ob}(\mathcal{Nec}) \rightarrow \mathbb{N} \times \mathbb{N}$ given by $\deg X = (\dim X, \ell(X))$

3.23. **THEOREM.** *The category \mathcal{Nec} equipped with $\mathcal{Nec}^{\leftarrow} = \{\text{epimorphisms}\}$, $\mathcal{Nec}^{\rightarrow} = \{\text{monomorphisms}\}$ and \deg is a Reedy category.*

PROOF. We already showed that $(\mathcal{Nec}^{\leftarrow}, \mathcal{Nec}^{\rightarrow})$ provides unique factorizations. Equip $\mathbb{N} \times \mathbb{N}$ with the lexicographical order, so that it may be identified with the ordinal ω^2 . Lemmas 3.19 and 3.20 show that every non-identity morphism in $\mathcal{Nec}^{\leftarrow}$ lowers degree and Lemma 3.18 shows that every non-identity morphism in $\mathcal{Nec}^{\rightarrow}$ raises degree. ■

3.24. **REMARK.** A decomposition of \mathcal{Nec} by degree is present in [27, §2]. By inspection of Table 3, one observes that other possible degree functions define Reedy structures with the (epi, mono)-factorization system (e.g. $\deg X = (\|X\|, \dim X)$). Note that any choice of degree function compatible with the factorization will give rise to an isomorphic Reedy structure.

3.25. **THEOREM.** *The category \mathcal{Nec} is simple Reedy monoidal.*

PROOF. We make use of Theorem 2.23. Using Remark 2.9 and noting that any map $X \rightarrow \Delta^0$ is an epimorphism, we conclude that \mathcal{Nec} is left fibrant. Consider a monomorphism $f : X \hookrightarrow Y_1 \vee Y_2$ that we write in combinatorial fashion

$$(T, p) \xrightarrow{f} (S_1, q_1) \vee (S_2, q_2) = (S_1 \cup (q_1 + S_2), q_1 + q_2) \quad (6)$$

We know that $q_1 \in S_1 \cup (S_2 + q_1) \subseteq f(T)$ and as $f : [p] \rightarrow [q_1 + q_2]$ is injective, there is a unique $r \in T$ such that $f(r) = q_1$. We can thus write $T_1 = (T \cap [r], r)$, $T_2 = ((T \cup \{r, \dots, p\}) - r, p - r)$ and $f_i = f|_{X_i}$. We have then unique f_1 and f_2 such that $f = f_1 \vee f_2$. We conclude that \mathcal{Nec} is direct divisible with respect to \vee . Finally, to show \mathcal{Nec} is simple we observe that pointwise sum in $\mathbb{N} \times \mathbb{N}$ corresponds to ordinal sum in ω^2 and that both dimension and bead length are additive with respect to \vee . ■

3.26. COROLLARY. *Let $(\mathbf{A}, \otimes, \mathbb{I})$ be a symmetric monoidal model category. Then $(\text{Fun}(\mathcal{N}ec^{op}, \mathbf{A})_{\text{Reedy}}, \otimes_{\text{Day}}, \Delta\mathbb{I})$ is a monoidal model category.*

References

- [1] H. V. Bacard, Segal enriched categories I, Version 1, Sept. 20 2010, arXiv: 1009.3673v1 [math.CT]
- [2] H. V. Bacard, Lax Diagrams and Enrichment, Version 1 June 16 2012, arXiv: 1206.3704v1 [math.CT]
- [3] H. V. Bacard, Colax Reedy diagrams, Version 1, July 27 2013, arXiv: 1307. 7215v1 [math.CT]
- [4] C. Barwick, On left and right model categories and left and right Bousfield localizations, In: Homology Homotopy Appl. 12.2 (2010), pp. 245–320, issn: 1532-0073,1532-0081.
- [5] H. J. Baues, Geometry of loop spaces and the cobar construction, In: Mem. Amer. Math. Soc. 25.230 (1980), pp. ix+171, issn: 0065-9266,1947-6221.
- [6] P. Bonventre and L. A. Pereira, Rigidification of dendroidal infinity-operads, In: Homology Homotopy Appl. 23.2 (2021), pp. 349–372, issn: 1532-0073,1532- 0081.
- [7] S. Barkan and J. Steinebrunner, Segalification and the Boardmann-Vogt tensor product, Version 1, 20 January 2023, arXiv: 2301.08650 [math.AT]
- [8] P. Bubenik, Simplicial models for concurrency, In: Proceedings of the Workshop on Geometric and Topological Methods in Computer Science (GETCO), Vol. 283, Electron. Notes Theor. Comput. Sci, Elsevier Sci. B. V., Amsterdam, 2012, pp. 3–12
- [9] P.-L. Curien, M. Livernet, and G. Saadia, Rigidification of cubical quasi-categories, Version 1, November 24 2022, arXiv: 2211.13679 [math.AT]
- [10] B. Day, On closed categories of functors, In: Reports of the Midwest Category Seminar, IV. Lecture Notes in Mathematics, Vol. 137. Springer, Berlin, 1970, pp. 1–38
- [11] D. Dugger and D. I. Spivak, Mapping spaces in quasi-categories, In: Algebr. Geom. Topol. 11.1 (2011), pp. 263–325
- [12] D. Dugger and D. I. Spivak, Rigidification of quasi-categories, In: Algebr. Geom. Topol. 11.1 (2011), pp. 225–261
- [13] H. Gindi, Rigidification and the Coherent Nerve for Enriched Quasicategories, Version 2, 29 September 2019, arXiv: 1810.10075 [math.CT]
- [14] M. Ghazel and F. Kadhi, Reedy diagrams in V-model categories, In: Appl. Categ. Structures 27.5 (2019), pp. 549–566. issn: 0927-2852,1572-9095.
- [15] D. Grady and D. Pavlov, Extended field theories are local and have classifying spaces, Version 4. Sept. 17 2023, arXiv: 2011.01208v4 [math.AT]
- [16] P. S. Hirschhorn, Model categories and their localizations, Vol. 99, Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2003, pp. xvi+457
- [17] M. Hovey, Model categories, Vol. 63, Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 1999, pp. xii+209
- [18] T. Leinster, Homotopy algebras for operads, Version 1, Feb. 22, 2000, arXiv: math/0002180v1 [math.QA]

- [19] W. Lowen and A. Mertens, Enriched quasi-categories and the templicial homotopy coherent nerve, Version 1. Feb. 5, 2023, arXiv: 2302.02484v1 [math.CT]
- [20] J. Lurie, Higher topos theory, Vol. 170, Annals of Mathematics Studies, Princeton University Press, Princeton, NJ, 2009, pp. xviii+925
- [21] A. Mertens, On nerves of enriched categories, In preparation.
- [22] A. M. Medina-Mardones and M. Rivera, Adams' cobar construction as a monoidal E_∞ -coalgebra model of the based loop space, Version 3, Jan. 14 2023. arXiv: 2108.02790 [math.AT]
- [23] E. Minichiello, M. Rivera, and M. Zeinalian, Categorical models for path spaces, In: Adv. Math. 415 (2023), Paper No. 108898, 48. issn: 0001-8708,1090- 2082.
- [24] C. L. Reedy, Homotopy theory of model categories, unpublished notes (1974), Available at the Hopf Topology Archive as <ftp://hopf.math.purdue.edu/pub/Reedy/reedy.dvi>
- [25] M. Rivera, An algebraic model for the free loop space, Version 5, Nov. 21 2023, arXiv: 2210.10096 [math.AT]
- [26] M. Rivera and S. Saneblidze, A combinatorial model for the free loop fibration, In: Bull. Lond. Math. Soc. 50.6 (2018), pp. 1085–1101. issn: 0024- 6093,1469-2120
- [27] M. Rivera and S. Saneblidze, A combinatorial model for the path fibration, In: J. Homotopy Relat. Struct. 14.2 (2019), pp. 393–410. issn: 2193-8407,1512- 2891.
- [28] M. Rivera and M. Zeinalian, Cubical rigidification, the cobar construction and the based loop space, In: Algebr. Geom. Topol. 18.7 (2018), pp. 3789– 3820. issn: 1472-2747.
- [29] C. Simpson, Homotopy theory of higher categories, Vol. 19, New Mathematical Monographs, Cambridge University Press, Cambridge, 2012.

*Department Wiskunde, Universiteit Antwerpen
Middelheimlaan 1, 2020 Antwerp, Belgium*

*Department Wiskunde, Universiteit Antwerpen
Middelheimlaan 1, 2020 Antwerp, Belgium*

Email: violeta.borgesmarques@uantwerpen.be
arne.mertens@uantwerpen.be

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS L^AT_EX₂ε is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Geoff Cruttwell, Mount Allison University: gcruttwell@mta.ca

T_EXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT T_EX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

John Bourke, Masaryk University: bourkej@math.muni.cz

Maria Manuel Clementino, Universidade de Coimbra: mmc@mat.uc.pt

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Rune Haugseng, Norwegian University of Science and Technology: rune.haug seng@ntnu.no

Dirk Hofmann, Universidade de Aveiro: dirk@ua.pt

Joachim Kock, Universitat Autònoma de Barcelona: Joachim.Kock@uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Sandra Mantovani, Università degli Studi di Milano: sandra.mantovani@unimi.it

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Giuseppe Metere, Università degli Studi di Palermo: giuseppe.metere@unipa.it

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

Jiri Rosický, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@unige.it

Michael Shulman, University of San Diego: shulman@sandiego.edu

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be

Christina Vasilakopoulou, National Technical University of Athens: cvasilak@math.ntua.gr