

A COHERENCE THEOREM FOR PSEUDO SYMMETRIC MULTIFUNCTORS

DIEGO MANCO

ABSTRACT. Donald Yau defined the notion of pseudo symmetric **Cat**-enriched multifunctor between **Cat**-enriched multicategories and proved that Mandell’s inverse K -theory multifunctor is pseudo symmetric. We prove a coherence theorem for pseudo symmetric **Cat**-enriched multifunctors. As an application we prove that pseudo symmetric **Cat**-enriched multifunctors, and in particular Mandell’s inverse K -theory, preserve Σ -free E_n -algebras ($n = 1, 2, \dots, \infty$), at the cost of changing the parameterizing Σ -free E_n -operad \mathcal{O} for the Σ -free E_n -operad $\mathcal{O} \times E\Sigma_*$.

1. Introduction

Permutative categories are symmetric monoidal categories that are strictly associative and unital. Let **Perm** be the category of permutative categories. By a construction of May [May, 1974], we can define algebraic K -theory as a functor from **Perm** to spectra. Elmendorf and Mandell [Elmendorf and Mandell, 2006] introduced multicategories in homotopy theory to study the multiplicative properties of this functor. They gave **Perm** the structure of a multicategory and showed that the K -theory construction can be extended to a symmetric multifunctor landing in spectra. This implies that K -theory preserves certain multiplicative structures—for example, the K -theory of a bipermutative category is an E_∞ -ring spectrum.

Following work of Thomason [Thomason, 1995], Mandell [Mandell, 2010] introduced inverse K -theory \mathcal{P} , a functor from Γ -categories (modelling connective spectra) to **Perm** that provides a homotopy inverse to K -theory. Elmendorf [Elmendorf, 2021] had initially extended \mathcal{P} to a symmetric multifunctor, but some mistakes were found as it is described in [Yau, 2024] Explanation 10.3.13. Johnson and Yau [Johnson and Yau, 2022] finally extended \mathcal{P} to a **Cat**-enriched multifunctor, but one that is not symmetric: it is not compatible with the permutation of elements in the domains of multicategory mapping spaces. To account for this Yau [Yau, 2024] introduced pseudo-symmetric multifunctors, where there is a compatibility only up to coherent natural isomorphisms, and he proved that Mandell’s inverse K -theory multifunctor \mathcal{P} is pseudo symmetric in his sense.

In this article we establish a 2-adjunction that lets us rigidify pseudo symmetric mul-

Received by the editors 2023-11-20 and, in final form, 2024-10-18.

Transmitted by Julie Bergner. Published on 2024-10-29.

2020 Mathematics Subject Classification: Primary 18M65, 19D23; Secondary 55P47, 55P43..

Key words and phrases: Multicategories, K -theory.

© Diego Manco, 2024. Permission to copy for private use granted.

tifunctors and write them as symmetric multifunctors at the cost of fattening up their domain in a specific way. As an application we get a new result in multiplicative K -theory: pseudo symmetric multifunctors, and in particular Mandell's inverse K -theory, preserve E_n -algebras for $n = 1, 2, \dots, \infty$ at the cost of changing the parameterizing E_n -operad. For example, they send commutative monoids to E_∞ -algebras.

Let us go back and provide more details of the above panorama. Segal's infinite loop space machine [Segal, 1974] allows the construction of spectra from symmetric monoidal categories. May's construction [May, 1974] provides an alternative way of building spectra from permutative categories. Both K -theory constructions turn out to be equivalent [May and Thomason, 1978], with **Perm** being equivalent to the category of symmetric monoidal categories by a theorem of Isbell [Isbell, 1969]. The question about what kind of structure to impose on a permutative category so that its K -theory is an E_∞ -ring spectrum was answered independently by Elmendorf and Mandell [Elmendorf and Mandell, 2006] and May [May, 2009], the former using the theory of multicategories. To study multiplicative K -theory, one would like the domain of the K -theory construction **Perm** to have a symmetric monoidal structure and K -theory to be a monoidal functor. That way, K -theory would preserve multiplicative structures in **Perm**. However, **Perm** lacks a natural symmetric monoidal structure, although it admits one in a 2-categorical sense [Gurski, Johnson, and Osorno].

Multicategories, also known as colored operads, generalize symmetric monoidal categories by supplying a setup for working with multi-input maps, thus providing an alternative way of encoding multiplicative structures even in the absence of symmetric monoidal structures. In a sense, multicategories allow us to talk about multilinear maps without making any reference to tensor products. Multiplicative structures can then be encoded in a multicategory via the actions of operads and similar gadgets. Elmendorf and Mandell [Elmendorf and Mandell, 2006] gave **Perm** the structure of a multicategory and extended algebraic K -theory to a symmetric multifunctor landing in symmetric spectra. This implies that K -theory preserves multiplicative structures. This is how they proved that the K -theory of a bipermutative category is an E_∞ -ring spectrum. Multiplicative K -theory has also been defined as a symmetric multifunctor from the multicategory of Waldhausen categories **Wald** to spectra [Blumberg and Mandell, 2011], with **Wald** providing another example of a multicategory that doesn't arise from a symmetric monoidal structure [Zkharevich, 2018].

Spectra arising from the Segal-May construction are all connective, and by a theorem of Thomason [Thomason, 1995] the K -theory construction is surjective on homotopy types. Mandell's inverse K -theory functor $\mathcal{P}: \Gamma\text{-Cat} \rightarrow \mathbf{Perm}$ witnesses this by providing a homotopy stable inverse to K -theory. Here Γ -categories model connective spectra by [Thomason, 1980, Cisinski, 1999, Bousfield and Friedlander, 78]. The question of whether Mandell's inverse K -theory functor could be extended to a multifunctor was first attacked by Elmendorf [Elmendorf, 2021]. After mistakes were found (see [Yau, 2024] Ex-

planation 10.3.13) Johnson and Yau settled the matter and proved that Mandell’s inverse K -theory is equivalent to a **Cat**-enriched multifunctor $\mathcal{P}: \Gamma\text{-Cat} \rightarrow \mathbf{Perm}$, albeit one that is not symmetric. The **Cat**-enriched multifunctor \mathcal{P} doesn’t preserve the action of the symmetric group on the hom objects of the multicategories by permutation of inputs. So, Johnson and Yau’s results can only be used to prove that \mathcal{P} preserves multiplicative structures that don’t involve symmetry, like associative monoids [Johnson and Yau, 2022]. This obstruction led Yau [Yau, 2024] to define pseudo symmetric multifunctors. These are non-symmetric **Cat**-enriched multifunctors that preserve the action of the symmetric group of multicategory mapping spaces only up to coherent natural isomorphisms. One of the main results of [Yau, 2024] is that \mathcal{P} is in fact pseudo symmetric.

Our main result can be interpreted as a coherence result for pseudo symmetric multifunctors. If $F: \mathcal{M} \rightarrow \mathcal{N}$ is a pseudo symmetric multifunctor between **Cat**-enriched multicategories, we prove that the natural isomorphisms attesting the pseudo symmetry of F assemble together to give a symmetric **Cat**-enriched multifunctor $\phi(F): \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$ satisfying a universal property, where $E\Sigma_*$ is the categorical Barratt-Eccles operad defined in Example 2.5. We can also think about our result as a rigidification result. We can rigidify F and turn it into a symmetric **Cat**-enriched multifunctor $\phi(F)$, at the cost of changing its domain.

1.1. THEOREM. (Theorem 3.3) Let \mathcal{M} be a **Cat**-enriched multicategory. There is a pseudo symmetric multifunctor $\eta_{\mathcal{M}}: \mathcal{M} \rightarrow \mathcal{M} \times E\Sigma_*$ such that for every **Cat**-enriched multicategory \mathcal{N} and pseudo symmetric multifunctor $F: \mathcal{M} \rightarrow \mathcal{N}$, there exists a unique symmetric **Cat**-enriched multifunctor $\phi(F): \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$ such that the following diagram commutes:

$$\begin{array}{ccc}
 & \mathcal{M} \times E\Sigma_* & \\
 \eta_{\mathcal{M}} \nearrow & & \searrow \phi(F) \\
 \mathcal{M} & \xrightarrow{F} & \mathcal{N}.
 \end{array}$$

That is, $F = \phi(F) \circ \eta_{\mathcal{M}}$ as pseudo symmetric multifunctors.

Thus, if \mathcal{O} is an operad in **Cat**, pseudo symmetric algebras in a **Cat**-enriched multicategory \mathcal{M} over \mathcal{O} , i.e., pseudo symmetric multifunctors $\mathcal{O} \rightarrow \mathcal{M}$, are symmetric algebras in \mathcal{M} over $\mathcal{O} \times E\Sigma_*$, i.e., symmetric multifunctors $\mathcal{O} \times E\Sigma_* \rightarrow \mathcal{M}$. The following result, which appears as Corollary 4.7, holds since multiplying by $E\Sigma_*$ sends the commutative operad $\{*\}$ to the E_∞ -operad $E\Sigma_*$ and Σ -free E_n -operads in **Cat**, like the ones defined in [Berger, 1996] and [Fiedorowicz et al., 2003], to E_n -operads.

1.2. COROLLARY. (Corollary 4.7) Let $F: \mathcal{M} \rightarrow \mathcal{N}$ be **Cat**-enriched pseudo symmetric multifunctor. Then,

1. F sends commutative monoids to E_∞ -algebras.

2. F sends symmetric E_n -algebras over a Σ -free E_n **Cat**-operad \mathcal{O} to symmetric E_n -algebras over the free E_n **Cat**-operad $\mathcal{O} \times E\Sigma_*$, for $n = 1, 2, \dots, \infty$.

In this sense, pseudo symmetric **Cat**-enriched multifunctors preserve symmetric E_n -algebras parameterized by free E_n -operads at the cost of changing the parameterizing operad. We notice that requiring that Σ_* acts freely on the E_n -operad \mathcal{O} doesn't impose a serious restriction since there are E_n **Cat**-operads ($n = 1, 2, \dots, \infty$) like the ones defined in [Berger, 1996] and [Fiedorowicz et al., 2003] that are Σ -free. This corollary extends our understanding of the behavior of inverse K -theory since it implies that the inverse K -theory multifunctor \mathcal{P} sends commutative monoids to E_∞ -algebras and sends E_n -algebras ($n = 1, 2, \dots$) parameterized by free Σ -operads to E_n -algebras. Since \mathcal{P} provides a stable inverse to K -theory, and K -theory is a symmetric multifunctor, this implies that every symmetric E_n -algebra parameterized by a Σ -free **Cat**-operad in Γ -categories is stably equivalent to the K -theory of an symmetric E_n -algebra in permutative categories for $n = 1, 2, \dots, \infty$. This shows how Theorem 1.1 can be used to grasp the behavior of pseudo symmetric multifunctors on structures parameterized by symmetric operads in general.

In [Yau, 2024] Yau defines the 2-category **Cat-Multicat** having **Cat**-enriched multicategories as 0-cells, symmetric multifunctors as 1-cells and multinatural transformations as 2-cells. He also defines the 2-category **Cat-Multicat^{ps}** with 0-cells **Cat**-enriched multicategories, 1-cells pseudo symmetric multifunctors, and 2-cells pseudo symmetric **Cat**-multinatural transformations. Every symmetric **Cat**-enriched multifunctor (respectively multinatural transformation) is canonically a pseudo symmetric multifunctor (respectively multinatural transformation), so there is a 2-functorial inclusion $j: \mathbf{Cat-Multicat} \rightarrow \mathbf{Cat-Multicat}^{\text{ps}}$. Taking into account these 2-categorical structures we can improve our previous result by providing a left adjoint ψ to j , which, at the 0-cell level, sends a multicategory \mathcal{M} to $\psi(\mathcal{M}) = \mathcal{M} \times E\Sigma_*$.

1.3. THEOREM. (Corollary 3.5 and Theorem 3.7) The inclusion $j: \mathbf{Cat-Multicat} \rightarrow \mathbf{Cat-Multicat}^{\text{ps}}$ admits a left 2-adjoint $\psi: \mathbf{Cat-Multicat}^{\text{ps}} \rightarrow \mathbf{Cat-Multicat}$ with $\psi(\mathcal{M}) = \mathcal{M} \times E\Sigma_*$ for \mathcal{M} a **Cat**-multicategory. In particular, for **Cat**-multicategories \mathcal{M} and \mathcal{N} we have an isomorphism of categories

$$\mathbf{Cat-Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N}) \cong \mathbf{Cat-Multicat}(\mathcal{M} \times E\Sigma_*, \mathcal{N}).$$

An important consequence of this theorem is that we can give a very simple and compact description of the 2-category **Cat-Multicat^{ps}** solely in terms of symmetric **Cat**-multifunctors and **Cat**-multinatural transformations, which we do in Definition 3.8.

The equivariant infinite loop space machine \mathbb{K}_G from [Guillou, May, Merling, and Orno, 2023] is also suspected to be pseudo symmetric, so our result might help understand the preservation of multiplicative structures in the equivariant context as well. A multifunctorial equivariant Algebraic K -theory construction was provided by Yau [Yau, 2024].

Our result could also be used to prove that the two equivariant K -theory constructions from [Guillou, May, Merling, and Osorno, 2023] and [Yau, 2024] are equivalent.

Outline. In Section 2 we recall the definition of the 2-categories **Cat-Multicat** and **Cat-Multicat**^{ps}. In Section 3 we prove Theorems 1.1 and 1.3. We also extract a new and compact description of the 2-category **Cat-Multicat**^{ps}. In Section 4 we obtain the desired consequences for Mandell’s inverse K -theory functor \mathcal{P} included in Corollary 1.2.

Acknowledgements. The author would like to thank Angélica Osorno, without whom this article wouldn’t exist, for her mentorship and guidance during all stages of this project. The author would like to thank Dan Dugger for providing helpful advice and feedback as well as Donald Yau for his helpful comments, questions, email exchanges, and encouragement. The author would also like to thank the referee for their very careful reading and helpful suggestions. The author was supported by the Fulbright-COLCIENCIAS scholarship awarded by the Fulbright Colombia Commission and COLCIENCIAS, now a part of the Ministry of Science, Technology, and Innovation of the Colombian government. The author was also supported by the University of Oregon through the Anderson Mathematics PhD Student Research Award.

2. Symmetric and pseudo symmetric Multifunctors

We begin by reviewing the definition of multicategory enriched in a symmetric monoidal category. In the following definition $(C, 1, \oplus, \lambda, \rho, \xi)$ is a symmetric monoidal category with $\oplus: C \times C \rightarrow C$ the monoidal product, 1 the monoidal unit, λ the left unit isomorphism, ρ the right unit isomorphism and ξ the symmetry. In this paper we will consider only categories enriched over **Cat** with the monoidal structure given by products, but we use a general monoidal category in the definition to make explicit the fact that this definition doesn’t make use of the 2-categorical structure of **Cat**.

2.1. **REMARK.** We will also use the following notation: if $\sigma \in \Sigma_n$ and $\tau_i \in \Sigma_{k_i}$ for $1 \leq i \leq n$, $\sigma \langle \tau_1, \dots, \tau_n \rangle \in \Sigma_{k_1 + \dots + k_n}$ is the permutation that permutes n blocks of lengths k_1, \dots, k_n according to σ and each block of length k_i according to τ_i .

2.2. **DEFINITION.** *If C is a symmetric monoidal category, a C -multicategory $(\mathcal{M}, \gamma, 1)$ consists of the following data.*

- A class of objects $Ob(\mathcal{M})$.
- For every $n \geq 0$, $\langle a \rangle = \langle a_i \rangle_{i=1}^n \in Ob(\mathcal{M})^n$ and $b \in Ob(\mathcal{M})$, an object in C denoted by

$$\mathcal{M}(\langle a \rangle; b) = \mathcal{M}(a_1, \dots, a_n; b).$$

We will write $\langle a \rangle$ instead of $\langle a_i \rangle_{i=1}^n$ when n is clear from the context or irrelevant. [In the case $C = \mathbf{Cat}$, an object f of $\mathcal{M}(\langle a \rangle; b)$ will be called an n -ary 1-cell with

input $\langle a \rangle$ and output b and will be denoted as $f: \langle a \rangle \rightarrow b$. Similarly, we will call $\alpha: f \rightarrow g$ in $\mathcal{M}(\langle a \rangle; b)(f, g)$ an n -ary 2-cell.]

- For each $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$, and $\sigma \in \Sigma_n$, a C -isomorphism

$$\mathcal{M}(\langle a \rangle; b) \xrightarrow[\cong]{\sigma} \mathcal{M}(\langle a \rangle \sigma; b)$$

called the right σ action or the symmetric group action. Here

$$\langle a \rangle \sigma = \langle a_1, \dots, a_n \rangle \sigma = \langle a_{\sigma(1)}, \dots, a_{\sigma(n)} \rangle.$$

[In the case $C = \mathbf{Cat}$ we write $f\sigma$ for the image of an n -ary 1-cell $f: \langle a \rangle \rightarrow b$ in \mathcal{M} and similarly for 2-cells.]

- For each object $a \in \text{Ob}(\mathcal{M})$, a morphism

$$1 \xrightarrow{1_a} \mathcal{M}(a; a)$$

called the a -unit. In the case $C = \mathbf{Cat}$ we notice that if $a \in \text{Ob}(\mathcal{M})$, $1_a: a \rightarrow a$ is a 1-ary 1-cell while if $f: \langle a \rangle \rightarrow b$ is an n -ary 1-cell, then $1_f: f \rightarrow f$ is an n -ary 2-cell in $\mathcal{M}(\langle a \rangle; b)(f, f)$ so our notation is unambiguous.

- For every $c \in \text{Ob}(\mathcal{M})$, $n \geq 0$, $\langle b \rangle = \langle b_j \rangle_{j=1}^n \in \text{Ob}(\mathcal{M})^n$, $k_j \geq 0$ for $1 \leq j \leq n$, and $\langle a_j \rangle = \langle a_{j,i} \rangle_{i=1}^{k_j} \in \text{Ob}(\mathcal{M})^{k_j}$ for $1 \leq j \leq n$, a morphism in C ,

$$\mathcal{M}(\langle b \rangle; c) \otimes \bigotimes_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) \xrightarrow{\gamma} \mathcal{M}(\langle a \rangle; c),$$

where we adopt the convention that $\langle a \rangle \in \text{Ob}(\mathcal{M})^k$, where $k = \sum_{i=1}^n k_j$, denotes the concatenation of the varying a_j 's for $j = 1, \dots, n$. We write this as

$$\langle a \rangle = \langle a_1, \dots, a_n \rangle = \langle \langle a_j \rangle \rangle_{j=1}^n = \langle a_{1,1}, \dots, a_{1,k_1}, a_{2,1}, \dots, a_{n-1,k_{n-1}}, a_{n,1}, \dots, a_{n,k_n} \rangle.$$

The previous data are required to satisfy the following axioms.

- **Symmetric group action:** For every $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})$, $b \in \text{Ob}(\mathcal{M})$, and σ, τ in Σ_n the following diagram commutes in C :

$$\begin{array}{ccc} \mathcal{M}(\langle a \rangle; b) & \xrightarrow{\sigma} & \mathcal{M}(\langle a \rangle \sigma; b) \\ & \searrow \sigma\tau & \downarrow \tau \\ & & \mathcal{M}(\langle a \rangle \sigma\tau; b). \end{array}$$

We also require the identity permutation $id_n \in \Sigma_n$ to act as the identity morphism on $\mathcal{M}(\langle a \rangle; b)$.

- **Associativity:** For every $d \in Ob(\mathcal{M})$, $n \geq 1$, $\langle c \rangle = \langle c_j \rangle_{j=1}^n \in Ob(\mathcal{M})^n$, $k_j \geq 0$ for $1 \leq j \leq n$ with $k_j \geq 1$ for at least one j , $\langle b_j \rangle = \langle b_{j,i} \rangle_{i=1}^{k_j} \in Ob(\mathcal{M})^{k_j}$ for $1 \leq j \leq n$, $l_{i,j} \geq 0$ for $1 \leq j \leq n$ and $1 \leq i \leq k_j$, and $\langle a_{j,i} \rangle = \langle a_{j,i,p} \rangle_{p=1}^{l_{i,j}} \in Ob(\mathcal{M})^{l_{i,j}}$ for $1 \leq j \leq n$ and $1 \leq i \leq k_j$, the following associativity diagram commutes in C :

$$\begin{array}{ccc}
 \mathcal{M}(\langle c \rangle; d) \otimes \left(\bigotimes_{j=1}^n \mathcal{M}(\langle b_j \rangle; c_j) \right) \otimes \bigotimes_{j=1}^n \left(\bigotimes_{i=1}^{k_j} \mathcal{M}(\langle a_{j,i} \rangle; b_{j,i}) \right) & & \\
 \downarrow \cong & \searrow \gamma \otimes 1 & \\
 & & \mathcal{M}(\langle b \rangle; c) \otimes \bigotimes_{j=1}^n \left(\bigotimes_{i=1}^{k_j} \mathcal{M}(\langle a_{j,i} \rangle; b_{j,i}) \right) \\
 & & \downarrow \gamma \\
 \mathcal{M}(\langle c \rangle; d) \otimes \bigotimes_{j=1}^n \left(\mathcal{M}(\langle b_j \rangle; c_j) \otimes \bigotimes_{i=1}^{k_j} \mathcal{M}(\langle a_{j,i} \rangle; b_{j,i}) \right) & & \\
 \downarrow 1 \otimes \bigotimes_{j=1}^n \gamma & & \\
 \mathcal{M}(\langle c \rangle; d) \otimes \bigotimes_{j=1}^n \mathcal{M}(\langle a_j \rangle; c_j) & \xrightarrow{\gamma} & \mathcal{M}(\langle a \rangle; b).
 \end{array} \tag{1}$$

- **Unity:** Suppose $b \in Ob(\mathcal{M})$ and $\langle a \rangle = \langle a_j \rangle_{j=1}^n \in Ob(\mathcal{M})$, then the following right unity diagram commutes in C :

$$\begin{array}{ccc}
 \mathcal{M}(\langle a \rangle; b) \otimes \bigotimes_{j=1}^n 1 & & \\
 \downarrow id \otimes \bigotimes_{j=1}^n 1_{a_j} & \searrow \cong & \\
 \mathcal{M}(\langle a \rangle; b) \otimes \bigotimes_{j=1}^n \mathcal{M}(a_j; a_j) & \xrightarrow{\gamma} & \mathcal{M}(\langle a \rangle; b).
 \end{array}$$

With $b, \langle a \rangle$ as before, we also demand that the following left unity diagram commutes in C .

$$\begin{array}{ccc}
 1 \otimes \mathcal{M}(\langle a \rangle; b) & & \\
 \downarrow 1_b \otimes id & \searrow \lambda & \\
 \mathcal{M}(b; b) \otimes \mathcal{M}(\langle a \rangle; b) & \xrightarrow{\gamma} & \mathcal{M}(\langle a \rangle; b).
 \end{array}$$

- **Top equivariance:** For every $c \in \text{Ob}(\mathcal{M})$, $n \geq 1$, $\langle b \rangle = \langle b_j \rangle_{j=1}^n \in \text{Ob}(\mathcal{M})^n$, $k_j \geq 0$ for $1 \leq j \leq n$, $\langle a_j \rangle = \langle a_{j,i} \rangle_{i=1}^{k_j} \in \text{Ob}(\mathcal{M})^{k_j}$ for $1 \leq j \leq n$, and $\sigma \in \Sigma_n$, the following diagram commutes:

$$\begin{array}{ccc}
 \mathcal{M}(\langle b \rangle; c) \otimes \bigotimes_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) & \xrightarrow{\sigma \otimes \sigma^{-1}} & \mathcal{M}(\langle b \rangle \sigma; c) \otimes \bigotimes_{j=1}^n \mathcal{M}(\langle a_{\sigma(j)} \rangle; b_{\sigma(j)}) \\
 \gamma \downarrow & & \downarrow \gamma \\
 \mathcal{M}(\langle a_1 \rangle, \dots, \langle a_n \rangle; c) & \xrightarrow{\sigma \langle id_{k_{\sigma(1)}}, \dots, id_{k_{\sigma(n)}} \rangle} & \mathcal{M}(\langle a_{\sigma(1)} \rangle, \dots, \langle a_{\sigma(n)} \rangle; c).
 \end{array} \tag{2}$$

Here σ^{-1} is the unique isomorphism in C , given by the coherence theorem for symmetric monoidal categories, that permutes the factors $\mathcal{M}(\langle a_j \rangle, b_j)$ according to σ^{-1} .

- **Bottom equivariance:** For $\langle a_j \rangle, \langle b \rangle$ and c as in Top equivariance 2, the following diagram commutes:

$$\begin{array}{ccc}
 \mathcal{M}(\langle b \rangle; c) \otimes \bigotimes_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) & \xrightarrow{id \otimes \bigotimes_{j=1}^n \tau_j} & \mathcal{M}(\langle b \rangle, c) \otimes \bigotimes_{j=1}^n \mathcal{M}(\langle a_j \rangle \tau_j; b_j) \\
 \gamma \downarrow & & \downarrow \gamma \\
 \mathcal{M}(\langle a_1 \rangle, \dots, \langle a_n \rangle; c) & \xrightarrow{id_n \langle \tau_1, \dots, \tau_n \rangle} & \mathcal{M}(\langle a_1 \rangle \tau_1, \dots, \langle a_n \rangle \tau_n; c).
 \end{array} \tag{3}$$

This concludes the definition of a C -multicategory.

2.3. REMARK. A C -operad is a C -multicategory with one object. If \mathcal{O} is a C -operad, its n -ary operations will be denoted by $\mathcal{O}_n \in \text{Ob}(C)$. A non symmetric C -multicategory (C -operad) is defined in the same way as a C -multicategory (C -operad) excluding the data of the symmetric group action as well as the symmetric group, top and bottom equivariance coherence axioms. We will only be concerned with symmetric multicategories and operads. C -multicategories are often referred to as colored operads, with the objects of the C -multicategory being referred to as colors and C -operads having just one color.

2.4. EXAMPLE. As examples of **Set**-operads, where **Set** has the monoidal structure induced by products in **Set**, we have the commutative operad $\text{Comm} = \{*\}$ with $\text{Comm}_n = \{*\}$. Another example is the associative operad $\text{Ass} = \Sigma_*$ with $\text{Ass}_n = \Sigma_n$, with the right action of the symmetric product given by right multiplication and γ defined in the following way. If $n \geq 1$ and k_1, \dots, k_n natural numbers with $k = \sum_{i=1}^n k_i$, we define $\gamma: \Sigma_n \times (\prod_{i=1}^n \Sigma_{k_i}) \rightarrow \Sigma_k$ given for $\sigma \in \Sigma_n$ and $\langle \tau_1, \dots, \tau_n \rangle \in \prod_{i=1}^n \Sigma_{k_i}$ by

$$\gamma(\sigma, \langle \rho_i \rangle_{i=1}^n) = \sigma \langle \rho_i \rangle_{i=1}^n = \sigma \langle \rho_1, \dots, \rho_n \rangle,$$

as in Remark 2.1. When n is clear from the context we will write $\sigma \langle \rho_i \rangle = \sigma \langle \rho_i \rangle_{i=1}^n$.

2.5. **EXAMPLE.** We will consider **Cat**-multicategories where the monoidal structure in **Cat** is given by products. One source of examples is the forgetful functor $\text{Ob}: \mathbf{Cat} \rightarrow \mathbf{Set}$ which forgets the morphism structure and remembers only the object set. Its right adjoint $E: \mathbf{Set} \rightarrow \mathbf{Cat}$ is the functor that takes a set A to EA , the category with objects $\text{Ob}(EA) = A$, and with a unique isomorphism between each pair of objects. E sends a morphism $f: A \rightarrow B$ of sets to the functor $Ef: EA \rightarrow EB$, the only functor such that $f = \text{Ob}(Ef)$. E preserves products, and thus, if \mathcal{O} is a **Set**-operad, $E\mathcal{O}$ is a **Cat**-operad. Similarly, if \mathcal{M} is a **Set**-multicategory, $E\mathcal{M}$ is a **Cat**-multicategory with the same collection of objects as \mathcal{M} .

We will call $E\text{Comm} = \{*\}$ the commutative **Cat**-operad. The Barratt-Eccles operad is the **Cat**-operad $E\Sigma_* = EAss$.

2.6. **EXAMPLE.** Another source of examples for multicategories are symmetric monoidal categories, and thus also permutative categories. Each symmetric monoidal category C has an associated **Set**-multicategory $\text{End}(C)$, whose objects agree with the objects of C and such that for $\langle a \rangle \in \text{Ob}(C)^n$ and $b \in \text{Ob}(C)$,

$$\text{End}(C)(\langle a \rangle; b) = C(a_1 \otimes \cdots \otimes a_n, b).$$

Here we take $a_1 \otimes \cdots \otimes a_n$ with the leftmost parenthesization. Any fixed parenthesization would work. An empty string of objects is interpreted as the monoidal unit $1 \in \text{Ob}(C)$.

Next, we define 1-cells between C -multicategories that preserve the action of the symmetric group. These are called symmetric C -multifunctors.

2.7. **DEFINITION.** A symmetric C -multifunctor $F: \mathcal{M} \rightarrow \mathcal{N}$ between C -multicategories \mathcal{M} and \mathcal{N} consists of the following data.

- An object assignment $F: \text{Ob}(\mathcal{M}) \rightarrow \text{Ob}(\mathcal{N})$.
- For each $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$ and $b \in \text{Ob}(\mathcal{M})$ a C morphism

$$\mathcal{M}(\langle a \rangle; b) \xrightarrow{F} \mathcal{N}(\langle Fa \rangle; Fb).$$

These data are required to preserve units, composition, and the action of the symmetric group.

- **Units:** For each object $a \in \text{Ob}(\mathcal{M})$, $F(1_a) = 1_{Fa}$, i.e., the following diagram commutes in C :

$$\begin{array}{ccc}
 & \mathcal{M}(a, a) & \\
 1_a \nearrow & & \searrow F \\
 1 & \xrightarrow{1_{Fa}} & \mathcal{N}(Fa, Fa).
 \end{array}$$

- **Composition:** For every $c \in \text{Ob}(\mathcal{M})$, $n \geq 0$, $\langle b \rangle = \langle b_j \rangle_{j=1}^n \in \text{Ob}(\mathcal{M})^n$, $k_j \geq 0$ for $1 \leq j \leq n$, and $\langle a_j \rangle = \langle a_{j,i} \rangle_{i=1}^{k_j} \in \text{Ob}(\mathcal{M})^{k_j}$ for $1 \leq j \leq n$ and $1 \leq i \leq k_j$, the following diagram commutes in \mathcal{C} :

$$\begin{array}{ccc}
 \mathcal{M}(\langle b \rangle; c) \otimes \bigotimes_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) & \xrightarrow{F \otimes \bigotimes_{j=1}^n F} & \mathcal{N}(\langle Fb \rangle; Fc) \otimes \bigotimes_{j=1}^n \mathcal{N}(\langle Fa_j \rangle; Fb_j) \\
 \downarrow \gamma & & \downarrow \gamma \\
 \mathcal{M}(\langle a \rangle; c) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle; Fc).
 \end{array} \tag{4}$$

- **Symmetric Group Action:** For each $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$ and $b \in \text{Ob}(\mathcal{M})$ the following diagram commutes in \mathcal{C} :

$$\begin{array}{ccc}
 \mathcal{M}(\langle a \rangle; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle; Fb) \\
 \cong \downarrow \sigma & & \cong \downarrow \sigma \\
 \mathcal{M}(\langle a \rangle \sigma; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle \sigma; Fb).
 \end{array}$$

2.8. DEFINITION. Let \mathcal{O} be a \mathcal{C} -operad and a \mathcal{M} be a \mathcal{C} -multicategory. A symmetric algebra in \mathcal{M} over \mathcal{O} is a symmetric multifunctor $\mathcal{O} \rightarrow \mathcal{M}$.

Symmetric algebras are usually called algebras, but we add the adjective symmetric to distinguish them from pseudo symmetric algebras, which will be defined later.

2.9. EXAMPLE. Since their introduction by May [May, 1972], operads have been used to characterize certain categories as the categories of symmetric algebras over a certain operad. For example, symmetric algebras over $Comm$ in **Set** are commutative monoids. Symmetric algebras over Σ_* in **Set** are associative monoids. Symmetric algebras over the Barrat-Eccles operad $E\Sigma_*$ in **Cat** are precisely permutative categories [May, 1974].

Next we define composition of \mathcal{C} -multifunctors.

2.10. DEFINITION. We define the horizontal composition of \mathcal{C} -multifunctors in the following way.

- Let $F: \mathcal{M} \rightarrow \mathcal{N}$, and $G: \mathcal{N} \rightarrow \mathcal{Q}$ be \mathcal{C} -multifunctors, we define the \mathcal{C} -multifunctor $GF: \mathcal{M} \rightarrow \mathcal{Q}$ on objects as the composition

$$\text{Ob}(\mathcal{M}) \xrightarrow{F} \text{Ob}(\mathcal{N}) \xrightarrow{G} \text{Ob}(\mathcal{Q}),$$

and its component functors for $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$ as the composite

$$\mathcal{M}(\langle a \rangle; b) \xrightarrow{F} \mathcal{N}(\langle Fa \rangle; Fb) \xrightarrow{G} \mathcal{Q}(\langle GFa \rangle; GFb).$$

- The identity C -multifunctor $1_{\mathcal{M}}: \mathcal{M} \rightarrow \mathcal{M}$ is defined as the identity assignment on objects with the identity functors as component functors.

Next we define 2-cells between C -multifunctors. These will be the 2-cells of a 2-category with 0-cells C -multicategories and 1-cells C -multifunctors.

2.11. DEFINITION. ([Yau, 2024], Def. 3.2.5) For (symmetric) C -multifunctors $F, G: \mathcal{M} \rightarrow \mathcal{N}$, we define a C -multinatural transformation $\theta: F \Rightarrow G$ as the data of a component morphism $\theta_a: 1 \rightarrow \mathcal{N}(Fa, Ga)$ in C for each $a \in \text{Ob}(\mathcal{M})$ subject to the commutativity of the following diagram in C for each $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$ and $b \in \text{Ob}(\mathcal{M})$,

$$\begin{array}{ccc} & 1 \otimes \mathcal{M}(\langle a \rangle; b) & \xrightarrow{\theta_b \otimes F} & \mathcal{N}(Fb; Gb) \otimes \mathcal{N}(\langle Fa \rangle; Fb) \\ & \nearrow \cong & & \searrow \gamma \\ \mathcal{M}(\langle a \rangle, b) & & & \mathcal{N}(\langle Fa \rangle; Gb). \\ & \searrow \cong & & \nearrow \gamma \\ & \mathcal{M}(\langle a \rangle; b) \otimes \bigotimes_{j=1}^n 1 & \xrightarrow{G \otimes \theta_{a_j}} & \mathcal{N}(\langle Ga \rangle; Gb) \otimes \bigotimes_{j=1}^n \mathcal{N}(Fa_j; Ga_j) \end{array}$$

We define the identity multinatural transformation $1_F: F \rightarrow F$ as having component $(1_F)_a = 1_{Fa}$ for a an object of \mathcal{M} .

2.12. REMARK. When $C = \mathbf{Cat}$, and given $F, G: \mathcal{M} \rightarrow \mathcal{N}$ \mathbf{Cat} -multifunctors and the data of a 1-ary 1-cell $\theta_a: Fa \rightarrow Ga$ for each $a \in \text{Ob}(\mathcal{M})$, the commutativity of the diagram in the previous definition means that for every $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$ and each 1-cell $f: \langle a \rangle \rightarrow b$,

$$\gamma(Gf; \langle \theta_{a_j} \rangle) = \gamma(\theta_b; Ff) \tag{5}$$

holds in $\mathcal{N}(\langle Fa \rangle; Gb)$ and that, for every 2-cell $\alpha: f \rightarrow g$ in $\mathcal{M}(\langle a \rangle; b)(f, g)$,

$$\gamma(G\alpha; \langle 1_{\theta_{a_j}} \rangle) = \gamma(1_{\theta_b}; F\alpha) \tag{6}$$

in $\mathcal{N}(\langle Fa \rangle; Gb)$. We can express (5) diagrammatically as the commutativity of the square

$$\begin{array}{ccc} \langle Fa \rangle & \xrightarrow{\langle \theta_{a_j} \rangle} & \langle Ga \rangle \\ Ff \downarrow & & \downarrow Gf \\ Fb & \xrightarrow{\theta_b} & Gb, \end{array}$$

where the composition of adjacent 1-cells is done through γ and a square represents an equality between composite 1-cells. In the same fashion, and using (5), we can express (6) as the equality of multicategorical pasting diagrams

$$\begin{array}{ccc}
 \langle Fa \rangle \xrightarrow{\langle \theta_{a_j} \rangle} \langle Ga \rangle & & \langle Fa \rangle \xrightarrow{\langle \theta_{a_j} \rangle} \langle Ga \rangle \\
 Ff \left(\begin{array}{c} \xrightarrow{F\alpha} \\ \xrightarrow{Fg} \end{array} \right) & \downarrow Gg & = & Ff \downarrow & Gf \left(\begin{array}{c} \xrightarrow{G\alpha} \\ \xrightarrow{Gg} \end{array} \right) \\
 Fb \xrightarrow{\theta_b} Gb & & & Fb \xrightarrow{\theta_b} Gb.
 \end{array}$$

Here the concatenation of adjacent 2-cells is done through γ , and an arrow labeled with the 1-cell h is interpreted as the 2-cell $1_h : h \rightarrow h$. For example, the left hand side diagram represents $\gamma(1_{\theta_b}, F\alpha)$ while the right hand side represents $\gamma(G\alpha, \langle \theta_{\alpha_j} \rangle)$. The empty squares represent equalities between composite 1-cells.

Next, we define horizontal and vertical compositions of C -multinatural transformations.

2.13. DEFINITION. ([Yau, 2024], Def. 3.2.7)

Suppose given $\theta : F \Rightarrow G, \zeta : G \Rightarrow H$ C -multinatural transformations with $F, G, H : \mathcal{M} \rightarrow \mathcal{N}$ C -multifunctors. The *vertical composition* $\zeta\theta : F \Rightarrow H$ is defined as having as component at each $a \in \text{Ob}(\mathcal{M})$ $(\zeta\theta)_a$, the composite

$$1 \xrightarrow{\cong} 1 \otimes 1 \xrightarrow{\zeta_a \otimes \theta_a} \mathcal{N}(Ga; Ha) \otimes \mathcal{N}(Fa; Ga) \xrightarrow{\gamma} \mathcal{N}(Fa; Ha).$$

Suppose that $\theta : F \Rightarrow G$ and $\zeta : F' \Rightarrow G'$ are C -multinatural transformations with $F, G : \mathcal{M} \rightarrow \mathcal{N}$ and $F', G' : \mathcal{N} \rightarrow \mathcal{Q}$ C -multifunctors. The *horizontal composition* $\zeta * \theta : F'F \Rightarrow G'G$ is defined as the C -multinatural transformation with component at each $a \in \text{Ob}(\mathcal{M})$, given by the composite

$$\begin{array}{ccc}
 1 & \xrightarrow{(\zeta * \theta)_a} & \mathcal{Q}(F'Fa; G'Ga) \\
 \downarrow \cong & & \uparrow \gamma \\
 1 \otimes 1 & \xrightarrow{\zeta_{Ga} \otimes \theta_a} \mathcal{Q}(F'Ga; G'Ga) \otimes \mathcal{N}(Fa; Ga) \xrightarrow{1 \otimes F'} & \mathcal{Q}(F'Ga; G'Ga) \otimes \mathcal{Q}(F'Fa; F'Ga).
 \end{array}$$

2.14. REMARK. When $C = \mathbf{Cat}$ and given $\theta : F \Rightarrow G, \zeta : G \Rightarrow H$ \mathbf{Cat} -multinatural transformations with $F, G, H : \mathcal{M} \rightarrow \mathcal{N}$ C -multifunctors and $a \in \text{Ob}(\mathcal{M})$,

$$(\zeta\theta)_a = \gamma(\zeta_a, \theta_a.) \tag{7}$$

On the other hand, if $\theta : F \Rightarrow G$ and $\zeta : F' \Rightarrow G'$ are \mathbf{Cat} -multinatural transformations with $F, G : \mathcal{M} \rightarrow \mathcal{N}$ and $F', G' : \mathcal{N} \rightarrow \mathcal{Q}$ \mathbf{Cat} -multifunctors,

$$(\zeta * \theta)_a = \gamma(\zeta_{Ga}, F'\theta_a). \tag{8}$$

Yau proves in [Yau, 2024] that Definitions 2.2, 2.7, 2.10, and 2.13 assemble together to give the 2-category $C\text{-Multicat}$, with 0-cells consisting of C -multicategories, 1-cells

symmetric C -multifunctors, and 2-cells C -multinatural transformations. There is a non symmetric variant where we drop the requirement that the C -multifunctors preserve the symmetric group action, as well as dropping the coherence axioms related to the symmetric group action, but we won't refer to this 2-category again.

For the rest of the article we fix our symmetric monoidal category C to be **Cat**, with the symmetric monoidal structure induced by products. In this context we can define a pseudo symmetric variant of this 2-category, namely **Cat-Multicat^{ps}** using the 2-categorical structure of **Cat**. The objects of **Cat-Multicat^{ps}** are still **Cat**-multicategories, but the 1-cells are pseudo symmetric **Cat**-multifunctors: **Cat**-multifunctors where we only require that they preserve the symmetric group action up to coherent isomorphisms.

2.15. DEFINITION. ([Yau, 2024] Def. 4.1.1) Suppose that \mathcal{M}, \mathcal{N} are **Cat**-multicategories. A *pseudo symmetric **Cat**-multifunctor* $F: \mathcal{M} \rightarrow \mathcal{N}$ consists of the following data:

- A function on object sets $F: \text{Ob}(\mathcal{M}) \rightarrow \text{Ob}(\mathcal{N})$.
- For each $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$ and $b \in \text{Ob}(\mathcal{M})$, a component functor

$$\mathcal{M}(\langle a \rangle; b) \xrightarrow{F} \mathcal{N}(\langle Fa \rangle; Fb).$$

- For each $\sigma \in \Sigma_n$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$, a natural isomorphism $F_{\sigma, \langle a \rangle, b}$

$$\begin{array}{ccc} \mathcal{M}(\langle a \rangle; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle; Fb) \\ \sigma \downarrow & \cong \nearrow_{F_{\sigma, \langle a \rangle, b}} & \downarrow \sigma \\ \mathcal{M}(\langle a \rangle \sigma; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle \sigma; Fb). \end{array}$$

When $\langle a \rangle$ and b are clear from the context we write simply F_σ , and if $f \in \text{Ob}(\mathcal{M}(\langle a \rangle, b))$ we will denote by $F_{\sigma, \langle a \rangle, b, f} = F_{\sigma; f}: F(f\sigma) \rightarrow F(f)\sigma$ the 2-cell in $\mathcal{N}(\langle Fa \rangle \sigma; Fb)$ corresponding to the component of F_σ at f . Naturality for F_σ means that given $\alpha: f \rightarrow g$ a 2-cell in $\mathcal{M}(\langle a \rangle; b)(f, g)$, the following diagram commutes in $\mathcal{N}(\langle Fa \rangle \sigma; b)$:

$$\begin{array}{ccc} F(f\sigma) & \xrightarrow{F_{\sigma; f}} & F(f)\sigma \\ F(\alpha\sigma) \downarrow & & \downarrow (F\alpha)\sigma \\ F(g\sigma) & \xrightarrow{F_{\sigma; g}} & F(g)\sigma. \end{array} \tag{9}$$

These data are subject to the same axioms of unit and composition preservation (4) as a symmetric **Cat**-multifunctor, but we replace the symmetric group action preservation axiom by the following four axioms.

- **Unit permutation:** Let $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$ and $b \in \text{Ob}(\mathcal{M})$, then

$$F_{\text{id}_n, \langle a \rangle, b} = 1_F. \tag{10}$$

- **Product permutation:** This axiom expresses the coherence of the natural isomorphisms F_σ , for varying σ , with respect to the symmetric group action. Let $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$ and $\sigma, \tau \in \Sigma_n$. Then, the following equality of pasting diagrams holds.

$$\begin{array}{ccc}
 \mathcal{M}(\langle a \rangle; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle; Fb) & & \mathcal{M}(\langle a \rangle; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle; Fb) \\
 \sigma \downarrow & \nearrow F_\sigma & \downarrow \sigma & & \downarrow \sigma\tau & \nearrow F_{\sigma\tau} & \downarrow \sigma\tau \\
 \mathcal{M}(\langle a \rangle\sigma; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle\sigma; Fb) & = & \mathcal{M}(\langle a \rangle\sigma\tau; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle\sigma\tau; Fb) \\
 \tau \downarrow & \nearrow F_\tau & \downarrow \tau & & & & \\
 \mathcal{M}(\langle a \rangle\sigma\tau; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle\sigma\tau; Fb) & & \mathcal{M}(\langle a \rangle\sigma\tau; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle\sigma\tau; Fb)
 \end{array}$$

Thus, for every 1-cell $f \in \text{Ob}(\mathcal{M}(\langle a \rangle; b))$, the following diagram of 2-cells commutes in $\mathcal{N}(\langle Fa \rangle; Fb)$:

$$\begin{array}{ccc}
 & F(f\sigma)\tau & \\
 F_{\tau; f\sigma} \nearrow & & \searrow (F_{\sigma; f})\tau \\
 F(f\sigma\tau) & \xrightarrow{F_{\sigma\tau; f}} & F(f)\sigma\tau.
 \end{array} \tag{11}$$

- **Top equivariance:** For every $c \in \text{Ob}(\mathcal{M})$, $n \geq 0$, $\langle b \rangle = \langle b_j \rangle_{j=1}^n \in \text{Ob}(\mathcal{M})^n$, $k_j \geq 0$ for $1 \leq j \leq n$, and $\langle a_j \rangle = \langle a_{j,i} \rangle_{i=1}^{k_j} \in \text{Ob}(\mathcal{M})^{k_j}$ for $1 \leq j \leq n$, and $\sigma \in \Sigma_n$, the following two pasting diagrams are equal.

$$\begin{array}{ccc}
 \mathcal{M}(\langle b \rangle; c) \times \prod_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) & \xrightarrow{F \times \prod_j F} & \mathcal{N}(\langle Fb \rangle; Fc) \times \prod_{j=1}^n \mathcal{N}(\langle Fa_j \rangle; Fb_j) \\
 \downarrow \gamma & & \downarrow \gamma \\
 \mathcal{M}(\langle \langle a_j \rangle \rangle_{j=1}^n; c) & \xrightarrow{F} & \mathcal{N}(\langle \langle Fa_j \rangle \rangle_{j=1}^n; Fc) \\
 \sigma \langle \text{id}_{k_{\sigma(j)}} \rangle \downarrow & \nearrow F_{\sigma \langle \text{id}_{k_{\sigma(j)}} \rangle} & \downarrow \sigma \langle \text{id}_{k_{\sigma(j)}} \rangle \\
 \mathcal{M}(\langle \langle a_{\sigma(j)} \rangle \rangle_{j=1}^n; c) & \xrightarrow{F} & \mathcal{N}(\langle \langle Fa_{\sigma(j)} \rangle \rangle_{j=1}^n; Fc) \\
 & \parallel & \\
 \mathcal{M}(\langle b \rangle; c) \times \prod_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) & \xrightarrow{F \times \prod_j F} & \mathcal{N}(\langle Fb \rangle; Fc) \times \prod_{j=1}^n \mathcal{N}(\langle Fa_j \rangle; Fb_j) \\
 \sigma \times \sigma^{-1} \downarrow & \nearrow F_{\sigma \times 1} & \downarrow \sigma \times \sigma^{-1} \\
 \mathcal{M}(\langle b \rangle; c) \times \prod_{j=1}^n \mathcal{M}(\langle a_{\sigma(j)} \rangle; b_{\sigma(j)}) & \xrightarrow{F \times \prod_j F} & \mathcal{N}(\langle Fb \rangle; Fc) \times \prod_{j=1}^n \mathcal{N}(\langle Fa_{\sigma(j)} \rangle; Fb_{\sigma(j)}) \\
 \downarrow \gamma & & \downarrow \gamma \\
 \mathcal{M}(\langle \langle a_{\sigma(j)} \rangle \rangle_{j=1}^n; c) & \xrightarrow{F} & \mathcal{N}(\langle \langle Fa_{\sigma(j)} \rangle \rangle_{j=1}^n; Fc)
 \end{array}$$

Here $\sigma \langle \text{id}_{k_{\sigma(j)}} \rangle = \sigma \langle \text{id}_{k_{\sigma(1)}}, \dots, \text{id}_{k_{\sigma(n)}} \rangle$. This means that for 1-cells $f \in \text{Ob}(\mathcal{M}(\langle b \rangle; c))$ and $g_j \in \text{Ob}(\mathcal{M}(\langle a_j \rangle; b_j))$ for $1 \leq j \leq n$,

$$F_{\sigma \langle \text{id}_{k_{\sigma(j)}} \rangle; \gamma(f; \langle g_j \rangle)} = \gamma \left(F_{\sigma; f}; \langle 1_{Fg_{\sigma(j)}} \rangle_{j=1}^n \right). \tag{12}$$

The domains and codomains of these pasting diagrams are equal by top equivariance in \mathcal{M} and \mathcal{N} , and the fact that F preserves γ implies the commutativity of the empty rectangles, see [Yau, 2024].

- Bottom Equivariance:** For every $c \in \text{Ob}(\mathcal{M})$, $n \geq 0$, $\langle b \rangle = \langle b_j \rangle_{j=1}^n \in \text{Ob}(M)^n$, $k_j \geq 0$ for $1 \leq j \leq n$, and $\langle a_j \rangle = \langle a_{j,i} \rangle_{i=1}^{k_j} \in \text{Ob}(\mathcal{M})^{k_j}$ for $1 \leq j \leq n$ and $1 \leq i \leq k_j$, and $\tau_j \in \Sigma_{k_j}$, the following two pasting diagrams are equal.

$$\begin{array}{ccc}
 \mathcal{M}(\langle b \rangle; c) \times \prod_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) & \xrightarrow{F \times \prod_j F} & \mathcal{N}(\langle Fb \rangle; Fc) \times \prod_{j=1}^n (\langle Fa_j \rangle; Fb_j) \\
 \downarrow \gamma & & \downarrow \gamma \\
 \mathcal{M}(\langle \langle a_j \rangle \rangle_{j=1}^n; c) & \xrightarrow{F} & \mathcal{N}(\langle \langle Fa_j \rangle \rangle_{j=1}^n; Fc) \\
 \text{id}_n \langle \tau_j \rangle \downarrow & \nearrow F_{\text{id}_n \langle \tau_i \rangle} & \downarrow \text{id}_n \langle \tau_j \rangle \\
 \mathcal{M}(\langle \langle a_j \rangle \tau_j \rangle_{j=1}^n; c) & \xrightarrow{F} & \mathcal{N}(\langle \langle Fa_j \rangle \tau_j \rangle_{j=1}^n; Fc) \\
 & \parallel & \\
 \mathcal{M}(\langle b \rangle; c) \times \prod_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) & \xrightarrow{F \times \prod_j F} & \mathcal{N}(\langle Fb \rangle; Fc) \times \prod_{j=1}^n \mathcal{N}(\langle Fa_j \rangle; Fb_j) \\
 \text{id} \times \prod_j \tau_j \downarrow & \nearrow 1 \times \prod_j F_{\tau_j} & \downarrow \text{id} \times \prod_j \tau_j \\
 \mathcal{M}(\langle b \rangle; c) \times \prod_{j=1}^n \mathcal{M}(\langle a_j \rangle \tau_j; b_j) & \xrightarrow{F \times \prod_j F} & \mathcal{N}(\langle Fb \rangle; Fc) \times \prod_{j=1}^n \mathcal{N}(\langle \langle Fa_j \rangle \tau_j \rangle; Fb_j) \\
 \downarrow \gamma & & \downarrow \gamma \\
 \mathcal{M}(\langle \langle a_j \rangle \tau_j \rangle_{j=1}^n; c) & \xrightarrow{F} & \mathcal{N}(\langle \langle Fa_j \rangle \tau_j \rangle_{j=1}^n; Fc)
 \end{array}$$

This means that for 1-cells $f: \langle b \rangle \rightarrow c$ and $g_j: \langle a_j \rangle \rightarrow b_j$ for $1 \leq j \leq n$,

$$F_{\text{id}_n \langle \tau_j \rangle; \gamma(f; \langle g_j \rangle)} = \gamma(1_{Ff}; \langle F_{\tau_j; g_j} \rangle) \tag{13}$$

as 2-cells in $\mathcal{N}(\langle \langle Fa_j \rangle \tau_j \rangle; Fc)$. The domain and codomain of these pasting diagrams are equal by bottom equivariance for \mathcal{M} and \mathcal{N} , and the preservation of γ by F guarantees that the empty squares commute, see [Yau, 2024].

Next we describe the horizontal composition of 1-cells in the 2-category **Cat-Multicat^{ps}**.

2.16. DEFINITION. ([Yau, 2024] Def. 4.1.1) Let $F: \mathcal{M} \rightarrow \mathcal{N}$, and $G: \mathcal{N} \rightarrow \mathcal{Q}$ be pseudo symmetric **Cat**-multifunctors. We define the pseudo symmetric functor $GF: \mathcal{M} \rightarrow \mathcal{Q}$. On objects GF is the composite function $GF: \text{Ob}(\mathcal{M}) \rightarrow \text{Ob}(\mathcal{Q})$. The composite component functor is given for $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, and $b \in \text{Ob}(\mathcal{M})$ by the pasting

$$\mathcal{M}(\langle a \rangle; b) \xrightarrow{F} \mathcal{N}(\langle Fa \rangle; b) \xrightarrow{G} \mathcal{Q}(\langle GFa \rangle; GFb).$$

The symmetry isomorphisms are given for each $\sigma \in \Sigma_n$, $\langle a \rangle \in \text{Ob}(\mathcal{M})$, and $b \in \text{Ob}(\mathcal{M})$ by

$$\begin{array}{ccccc}
 \mathcal{M}(\langle a \rangle; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle; Fb) & \xrightarrow{G} & \mathcal{Q}(\langle GFa \rangle; GFb) \\
 \sigma \downarrow & \nearrow F_{\sigma, \langle a \rangle, b} & \downarrow \sigma & \nearrow G_{\sigma, \langle Fa \rangle, Fb} & \downarrow \sigma \\
 \mathcal{M}(\langle a \rangle \sigma; b) & \xrightarrow{F} & \mathcal{N}(\langle Fa \rangle \sigma; fb) & \xrightarrow{G} & \mathcal{Q}(\langle GFa \rangle \sigma; GFb).
 \end{array}$$

That is, for each 1-cell $f: \langle a \rangle \rightarrow b$, the f component of GF_σ is given by the composite

$$\begin{array}{ccc}
 & G((Ff)\sigma) & \\
 G(F_{\sigma;f}) \nearrow & & \searrow G_{\sigma;Ff} \\
 GF(f\sigma) & \xrightarrow{(GF)_{\sigma;f}} & (GFf)\sigma.
 \end{array} \tag{14}$$

Next we define the 2-cells of the category **Cat-Multicat**^{PS}.

2.17. DEFINITION. ([Yau, 2024] Def. 4.2.1) Suppose that $F, G: \mathcal{M} \rightarrow \mathcal{N}$ are pseudo symmetric **Cat**-multifunctors. A pseudo symmetric **Cat**-multinatural transformation $\theta: F \Rightarrow G$ is the data of a component 1-cell $\theta_a : Fa \rightarrow Ga$ for each $a \in \text{Ob}(\mathcal{M})$ subject to axioms (5), (6) and the following extra axiom. For each $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$, object $f \in \text{Ob}(\mathcal{M}(\langle a \rangle; b))$, and permutation $\sigma \in \Sigma_n$, the following arrow equality holds in the category $\mathcal{N}(\langle Fa \rangle \sigma; b)$,

$$\gamma(1_{\theta_b}; F_{\sigma;f}) = \gamma(G_{\sigma;f}; \langle 1_{\theta_{a_{\sigma(j)}}} \rangle). \tag{15}$$

This can also be expressed diagrammatically as the equality of multicategorical pasting diagrams

$$\begin{array}{ccc}
 \langle Fa \rangle \sigma \xrightarrow{\langle \theta_{a_{\sigma(j)}} \rangle} G \langle a \rangle \sigma & & \langle Fa \rangle \sigma \xrightarrow{\langle \theta_{a_{\sigma(j)}} \rangle} \langle Ga \rangle \sigma \\
 F(f\sigma) \left(\begin{array}{c} \xrightarrow{F_{\sigma;f}} \\ \xrightarrow{(Ff)\sigma} \end{array} \right) \downarrow (Gf)\sigma & = & F(f\sigma) \downarrow \left(\begin{array}{c} \xrightarrow{G_{\sigma;f}} \\ \xrightarrow{(Gf)\sigma} \end{array} \right) \\
 Fb \xrightarrow{\theta_b} Gb & & Fb \xrightarrow{\theta_b} Gb,
 \end{array}$$

where the diagrams are interpreted as in Remark 2.12, the squares commuting by (5) and top and bottom equivariance for \mathcal{N} , see [Yau, 2024].

We define the vertical and horizontal composition of pseudo symmetric **Cat**-multinatural transformations in the same way that we did for symmetric ones, through diagrams (7) and (8).

It is a theorem of Yau [Yau, 2024] that the data we have just defined gives the structure of a 2-category, namely **Cat-Multicat**^{PS}. Definition 3.8 says that we can describe this 2-category solely in terms of symmetric **Cat**-multifunctors and symmetric **Cat**-multinatural transformations.

3. Coherence

To prove our first result we use finite products in the category **Cat-Multicat**. Having just the 1-categorical structure in mind, the products in **Cat-Multicat** are given in

the following way. If \mathcal{M} and \mathcal{N} are two **Cat**-multicategories, then $\mathcal{M} \times \mathcal{N}$ has objects $\text{Ob}(\mathcal{M} \times \mathcal{N}) = \text{Ob}(\mathcal{M}) \times \text{Ob}(\mathcal{N})$. Now, for $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $\langle c \rangle \in \text{Ob}(\mathcal{N})^n$, $b \in \text{Ob}(\mathcal{M})$, and $d \in \text{Ob}(\mathcal{N})$, we define

$$\mathcal{M} \times \mathcal{N}(\langle (a, c) \rangle; (b, d)) = \mathcal{M}(\langle a \rangle; b) \times \mathcal{N}(\langle c \rangle; d).$$

The composition γ of $\mathcal{M} \times \mathcal{N}$, as well as the Σ_* action and the multicategorical units, are defined componentwise. Next we define the pseudo symmetric multifunctor $\eta_{\mathcal{M}}$ appearing in the statement of 1.1.

3.1. DEFINITION. *Let \mathcal{M} be a **Cat**-multicategory. We define the pseudo symmetric **Cat**-multifunctor $\eta_{\mathcal{M}}: \mathcal{M} \rightarrow \mathcal{M} \times E\Sigma_*$ which, when there is no room for confusion, we will denote η . For an object $a \in \text{Ob}(\mathcal{M})$ as $\eta(a) = (a, *)$. We will abuse notation and denote the object $(a, *)$ of $\mathcal{M} \times E\Sigma_*$ as a .*

For $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$ and $b \in \text{Ob}(\mathcal{M})$ we need to define a functor $\eta: \mathcal{M}(\langle a \rangle; b) \rightarrow \mathcal{M}(\langle a \rangle; b) \times E\Sigma_n$. For a 1-cell $f: \langle a \rangle \rightarrow b$, we define

$$\eta(f) = (f, id_n) \in \text{Ob}(\mathcal{M}(a; b) \times E\Sigma_n).$$

Similarly, for a 2-cell $\alpha: f \rightarrow g$ in $\mathcal{M}(\langle a \rangle; b)$,

$$\eta(\alpha) = (\alpha, 1_{id_n}) \in \mathcal{M}(\langle a \rangle; b) \times E\Sigma_n((f, id_n), (g, id_n)).$$

Next, we define the components of the pseudo symmetry isomorphisms. For $\sigma, \tau \in \Sigma_n$ we will denote from here on by E_{σ}^{τ} the unique arrow $\sigma \rightarrow \tau$ in $E\Sigma_n$. For $\sigma \in \Sigma_n$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, and $b \in \text{Ob}(\mathcal{M})$ we need to define a natural isomorphism $\eta_{\sigma, \langle a \rangle, b}: (\eta \circ \sigma) \rightarrow (\sigma \circ \eta)$ that fits in the following diagram

$$\begin{array}{ccc} \mathcal{M}(\langle a \rangle; b) & \xrightarrow{\eta} & \mathcal{M}(\langle a \rangle; b) \times E\Sigma_n \\ \sigma \downarrow & \xrightarrow[\cong]{\eta_{\sigma, \langle a \rangle, b}} & \downarrow \sigma \times \sigma \\ \mathcal{M}(\langle a \rangle \sigma, b) & \xrightarrow{\eta} & \mathcal{M}(\langle a \rangle \sigma; b) \times E\Sigma_n. \end{array}$$

The isomorphism $\eta_{\sigma, \langle a \rangle, b}$ is defined for every 1-cell $f: \langle a \rangle \rightarrow b$ as the 2-cell

$$\eta_{\sigma; f} = (1_{f\sigma}, E_{id}^{\sigma}): (f\sigma, id_n) \rightarrow (f\sigma, \sigma).$$

3.2. LEMMA. *Let \mathcal{M} be a **Cat**-multicategory, then $\eta_{\mathcal{M}}: \mathcal{M} \rightarrow \mathcal{M} \times E\Sigma_*$ is pseudo symmetric.*

PROOF. We start from a non symmetric multifunctor $\eta: \mathcal{M} \rightarrow \mathcal{M} \times E\Sigma_*$ that is the identity on the first coordinate and the multicategorical unit in the second coordinate. As a non symmetric multifunctor, η preserves units and γ composition. We need to show that η is a pseudo symmetric **Cat**-multifunctor. The naturality of $\eta_{\sigma; f}$ follows from the commutativity of the following diagram for any 2-cell $\alpha: f \rightarrow g$:

$$\begin{array}{ccc}
 (f\sigma, \text{id}_n) & \xrightarrow{(1_{f\sigma}, E_{\text{id}_n}^\sigma)} & (f\sigma, \sigma) \\
 (\alpha\sigma, 1_{\text{id}_n}) \downarrow & & \downarrow (\alpha\sigma, 1_\sigma) \\
 (g\sigma, \text{id}_n) & \xrightarrow{(1_{g\sigma}, E_{\text{id}_n}^\sigma)} & (g\sigma, \sigma).
 \end{array}$$

Next we focus on the coherence axioms. The unit permutation axiom (10) holds since, for all $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$, and $f: \langle a \rangle \longrightarrow b$,

$$\eta_{\text{id}_n;f} = (1_{f\text{id}_n}, E_{\text{id}_n}^{\text{id}_n}) = (1_f, 1_{\text{id}_n}) = 1_{(f, \text{id}_n)} = 1_{\eta(f)}.$$

Let $\langle a \rangle, b$ and f be as before, the product permutation axiom (11) holds again by definition. Indeed, for $\tau, \sigma \in \Sigma_n$, we have

$$\eta_{\sigma\tau;f} = (1_{f\sigma\tau}, E_{\text{id}}^{\sigma\tau}) = (1_{f\sigma\tau}, E_\tau^{\sigma\tau}) \circ (1_{f\sigma\tau}, E_{\text{id}_n}^\tau) = (\eta_{\sigma;f\tau}) \circ \eta_{\tau;f\sigma}.$$

For Top Equivariance (12), suppose that $c \in \text{Ob}(\mathcal{M})$, $n \geq 1$, $\langle b \rangle = \langle b_j \rangle_{j=1}^n \in \text{Ob}(\mathcal{M})^n$, $k_j \geq 0$ for $1 \leq j \leq n$, $\langle a_j \rangle = \langle a_{j,i} \rangle_{i=1}^{k_j} \in \text{Ob}(\mathcal{M})^{k_j}$ for $1 \leq j \leq n$, $\sigma \in \Sigma_n$, $f \in \text{Ob}(\mathcal{M}(\langle b \rangle; c))$, and $g_j \in \text{Ob}(\mathcal{M}(\langle a_j \rangle; b_j))$. We have that

$$\begin{aligned}
 \gamma(\eta_{\sigma;f}; \langle 1_{i(g_{\sigma(j)})} \rangle) &= \gamma((1_{f\sigma}, E_{\text{id}}^\sigma); \langle (1_{g_{\sigma(j)}}, 1_{\text{id}_{k_{\sigma(j)}}}) \rangle) \\
 &= \left(\gamma(1_{f\sigma}; 1_{g_{\sigma(j)}}), \gamma \left(E_{\text{id}}^\sigma; E_{\text{id}_{k_{\sigma(j)}}}^{\text{id}_{k_{\sigma(j)}}} \right) \right) \\
 &= \left(1_{\gamma(f; \langle g_{\sigma(j)} \rangle)}, E_{\text{id}_{\langle \text{id}_{k_{\sigma(j)}} \rangle}}^{\sigma(\text{id}_{k_{\sigma(j)}})} \right) \\
 &= \left(1_{\gamma(f; \langle g_j \rangle) \sigma(\text{id}_{k_{\sigma(j)}})}, E_{\text{id}_k}^{\sigma(\text{id}_{k_{\sigma(j)}})} \right) \\
 &= \eta_{\sigma(\text{id}_{k_{\sigma(j)}}); \gamma(f; \langle g_j \rangle)}.
 \end{aligned}$$

For Bottom Equivariance, let $c, n, \langle b \rangle, k_j$ for $1 \leq j \leq n$, $\langle a_j \rangle$ for $1 \leq j \leq n$, f and g_j be as above and let $\tau_j \in \Sigma_{k_j}$ for $1 \leq j \leq n$. We also let $k = \sum_{j=1}^n k_j$. Bottom Equivariance (13) for i is

$$\begin{aligned}
 \gamma(1_{if}; \langle \eta_{\tau_j; g_j} \rangle) &= \gamma \left((1_f, 1_{\text{id}_n}); \langle (1_{g_j\tau_j}, E_{\text{id}_{k_j}}^{\tau_j}) \rangle \right) \\
 &= \left(\gamma(1_f; 1_{g_j\tau_j}), 1_{\text{id}_n} \langle E_{\text{id}_{k_j}}^{\tau_j} \rangle \right) \\
 &= \left(1_{\gamma(f; \langle g_j\tau_j \rangle)}, E_{\text{id}_k}^{\text{id}_n \langle \tau_j \rangle} \right) \\
 &= \left(1_{\gamma(f; \langle g_j \rangle) \text{id}_n \langle \tau_j \rangle}, E_{\text{id}_k}^{\text{id}_n \langle \tau_j \rangle} \right) \\
 &= \eta_{\text{id} \langle \tau_j \rangle; \gamma(f; \langle g_j \rangle)}.
 \end{aligned}$$

Thus, we conclude that $\eta: \mathcal{M} \longrightarrow \mathcal{M} \times E\Sigma_*$ is a pseudo symmetric **Cat**-multifunctor. ■

Recall that $j: \mathbf{Cat}\text{-Multicat} \rightarrow \mathbf{Cat}\text{-Multicat}^{\text{PS}}$ denotes the inclusion functor. We are ready to present a proof of 1.1.

3.3. THEOREM. *Let \mathcal{M} and \mathcal{N} be \mathbf{Cat} -multicategories and $F: \mathcal{M} \rightarrow \mathcal{N}$ a pseudo symmetric \mathbf{Cat} -multifunctor. There exists a unique symmetric \mathbf{Cat} -multifunctor $\phi(F): \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$ such that the following diagram commutes:*

$$\begin{array}{ccc}
 & \mathcal{M} \times E\Sigma_* & \\
 \eta_{\mathcal{M}} \nearrow & & \searrow j\phi(F) \\
 \mathcal{M} & \xrightarrow{F} & \mathcal{N}
 \end{array}$$

That is, $F = j\phi(F) \circ \eta_{\mathcal{M}}$ in $\mathbf{Cat}\text{-Multicat}^{\text{PS}}$.

PROOF PROOF OF THEOREM 1.1. For uniqueness, suppose that $\phi(F): \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$ is a symmetric \mathbf{Cat} -multifunctor satisfying $F = (j\phi(F)) \circ \eta$. We will abuse notation and write $j\phi(F) = \phi(F)$. We will prove there is a unique way of defining $\phi(F)$. At the level of the objects of the multicategory we must have $\phi(F)(a, *) = \phi(F) \circ \eta(a) = F(a)$ for each $a \in \text{Ob}(\mathcal{M})$. Next, we show that there is a unique way of defining each component functor of $\phi(F)$. For this let $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$, and consider the functor $\phi(F): \mathcal{M}(\langle a \rangle; b) \times E\Sigma_n \rightarrow \mathcal{N}(\langle Fa \rangle; Fb)$. If $f: \langle a \rangle \rightarrow b$ is a 1-cell and $\sigma \in \Sigma_n$, we must have that

$$\begin{aligned}
 \phi(F)(f, \sigma) &= \phi(F)((f\sigma^{-1}, \text{id}_n)\sigma) \\
 &= \phi(F)((f\sigma^{-1}, \text{id}_n))\sigma \\
 &= \phi(F) \circ \eta(f\sigma^{-1})\sigma \\
 &= F(f\sigma^{-1})\sigma,
 \end{aligned} \tag{16}$$

where in the second equality we used that $\phi(F)$ is symmetric. So the values of the component functors of $\phi(F)$ on n -ary 1-cells are uniquely determined by F . In exactly the same fashion, for $\langle a \rangle, b$ and σ as before, $f, g: \langle a \rangle \rightarrow b$, and $\alpha: f \rightarrow g$ a 2-cell,

$$\phi(F)(\alpha, 1_{\sigma}) = F(\alpha\sigma^{-1})\sigma. \tag{17}$$

Finally, if f, σ are as before and $\tau \in \Sigma_n$, we get that

$$\begin{aligned}
 \phi(F)(1_f, E_{\sigma}^{\tau}) &= \phi(F)(1_{f\sigma^{-1}}\sigma, E_{\text{id}}^{\tau\sigma^{-1}}\sigma) \\
 &= \phi(F)((1_{f\sigma^{-1}}, E_{\text{id}}^{\tau\sigma^{-1}}))\sigma \\
 &= \phi(F)(\eta_{\tau\sigma^{-1}; f\tau^{-1}})\sigma \\
 &= (\phi(F) \circ \eta_{\tau\sigma^{-1}; f\tau^{-1}})\sigma \\
 &= (F_{\tau\sigma^{-1}; f\tau^{-1}})\sigma.
 \end{aligned} \tag{18}$$

We have used the definition of composition of pseudo symmetric \mathbf{Cat} -multifunctors (14) where we see $\phi(F)$ trivially as a pseudo symmetric functor. For $\langle a \rangle, b, f, g, \alpha, \sigma$, and τ as

before, we can write the morphism $(\alpha: f \rightarrow g, E_\sigma^\tau)$ in $\mathcal{M}(\langle a \rangle; b) \times \Sigma_n$ as $(1_y, E_\sigma^\tau) \circ (f, 1_\sigma)$. Since both $\phi(F)(1_y, E_\sigma^\tau)$ and $\phi(F)(f, 1_\sigma)$ are uniquely determined by F , we conclude that the component functors of $\phi(F)$ are uniquely determined. We have proven the uniqueness of $\phi(F)$.

Next we prove the existence of $\phi(F)$. By uniqueness, we have no choice but to define $\phi(F)(b, *) = Fb$ for any $b \in \text{Ob}(M)$. Likewise, for $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$ and $b \in \text{Ob}(\mathcal{M})$, uniqueness forces the definition of the component functor $\phi(F): \mathcal{M}(\langle a \rangle; b) \times \Sigma_n \rightarrow \mathcal{N}(\langle Fa \rangle; b)$. For $f: \langle a \rangle \rightarrow b$, a 1-cell in $\mathcal{M}(\langle a \rangle; b)$ and $\sigma \in \Sigma_n$ we define

$$\phi(F)(f, \sigma) = F(f\sigma^{-1})\sigma \tag{19}$$

as in (16). For a 2-cell $\alpha: f \rightarrow g$ in $\mathcal{M}(\langle a \rangle; b)(f, g)$, we define

$$\phi(F)(\alpha, 1_\sigma) = F(\alpha\sigma^{-1})\sigma \tag{20}$$

as in (17). For $\tau \in \Sigma_n$ we define

$$\phi(F)(1_f, E_\sigma^\tau) = (F_{\tau\sigma^{-1}; f\tau^{-1}})\sigma \tag{21}$$

as in (18).

We still have to prove that $\phi(F) : \mathcal{M}(\langle a \rangle; b) \times \Sigma_n$ is well defined and extend our definition to all 2-cells. Notice that for a 1-cell $f: \langle a \rangle \rightarrow b$ our definition is ambiguous for the identity arrow $(1_f, 1_\sigma)$ since both (20) and (21) apply. However, $\phi(F)$ is well defined in this case since F is a functor componentwise and so, it preserves identities. Explicitly,

$$F(1_f\sigma^{-1})\sigma = F(1_{f\sigma^{-1}})\sigma = 1_{F(f\sigma^{-1})}\sigma = 1_{F(f\sigma^{-1})\sigma},$$

and

$$(F_{\sigma\sigma^{-1}, f\sigma^{-1}})\sigma = F_{\text{id}_n, f\sigma^{-1}}\sigma = 1_{F(f\sigma^{-1})}\sigma = 1_{F(f\sigma^{-1})\sigma}.$$

So, our definition is so far unambiguous and $\phi(F)$ preserves identities. We go on to extend the definition of $\phi(F)$ to the rest of the arrows. For $\alpha: f \rightarrow g$ 2-cell in $\mathcal{M}(\langle a \rangle, b)$ and σ, τ in Σ_n , we define $\phi(F)(\alpha, E_\sigma^\tau): F(f\sigma^{-1})\sigma \rightarrow F(g\tau^{-1})\tau$ by

$$\begin{aligned} \phi(F)(\alpha, E_\sigma^\tau) &= \phi(F)(1_g, E_\sigma^\tau) \circ \phi(F)(\alpha, 1_\sigma) \\ &= \phi(F)(\alpha, 1_\tau) \circ \phi(F)(1_f, E_\sigma^\tau). \end{aligned} \tag{22}$$

The last equality together with the preservation of identities already proven implies that our definition is unambiguous. This equality holds since,

$$\begin{aligned} \phi(F)(1_g, E_\sigma^\tau) \circ \phi(F)(\alpha, 1_\sigma) &= (F_{\tau\sigma^{-1}; g\tau^{-1}})\sigma \circ F(\alpha\sigma^{-1})\sigma \\ &= (F_{\tau\sigma^{-1}; g\tau^{-1}} \circ F(\alpha\sigma^{-1}))\sigma \\ &= (F(\alpha\tau^{-1})\tau\sigma^{-1} \circ F_{\tau\sigma^{-1}; f\tau^{-1}})\sigma \\ &= F(\alpha\tau^{-1})\tau \circ (F_{\tau\sigma^{-1}; f\tau^{-1}})\sigma \\ &= \phi(F)(\alpha, 1_\tau) \circ \phi(F)(1_f, E_\sigma^\tau). \end{aligned}$$

The third equality holds since it is precisely the commutativity of the following diagram:

$$\begin{array}{ccc}
 F(f\tau^{-1}\tau\sigma^{-1}) & \xrightarrow{F_{\tau\sigma^{-1};f\tau^{-1}}} & F(f\tau^{-1})\tau\sigma^{-1} \\
 F_{(\alpha\tau^{-1}\tau\sigma^{-1})}\downarrow & & \downarrow_{(F\alpha\tau^{-1})\tau\sigma^{-1}} \\
 F(g\tau^{-1}\tau\sigma^{-1}) & \xrightarrow{F_{\tau\sigma^{-1};g\tau^{-1}}} & F(g\tau^{-1})\tau\sigma^{-1}.
 \end{array} \tag{23}$$

This diagram commutes since it is an instance of the pseudo symmetry naturality coherence axiom for F , (9). Next, we check that the defined assignments give a functor $\phi(F): \mathcal{M}(\langle a \rangle; b) \times E\Sigma_n \rightarrow \mathcal{N}(\langle Fa \rangle; b)$. The fact that $\phi(F)$ preserves identities was already proven. We prove functoriality in the second variable first. For $f: \langle a \rangle \rightarrow b$ 1-cell, σ, τ , and ρ in Σ_n ,

$$\begin{aligned}
 \phi(F)(1_f, E_\tau^\rho) \circ \phi(F)(1_f, E_\sigma^\tau) &= (F_{\rho\tau^{-1};f\rho^{-1}\tau}) \circ (F_{\tau\sigma^{-1};f\tau^{-1}\sigma}) \\
 &= ((F_{\rho\tau^{-1};f\rho^{-1}})\tau\sigma^{-1} \circ F_{\tau\sigma^{-1};f\tau^{-1}}) \sigma \\
 &= (F_{\rho\sigma^{-1};f\rho^{-1}}) \sigma \\
 &= \phi(F)(1_f, E_\sigma^\rho).
 \end{aligned} \tag{24}$$

Here the third equality holds by (11), which implies the commutativity of the following diagram:

$$\begin{array}{ccc}
 & F(f\rho^{-1}\rho\tau^{-1})\tau\sigma^{-1} & \\
 F_{\tau\sigma^{-1};f\rho^{-1}\rho\tau^{-1}} \nearrow & & \searrow_{(F_{\rho\tau^{-1};f\rho^{-1}})\tau\sigma^{-1}} \\
 F(f\rho^{-1}\rho\tau^{-1}\tau\sigma^{-1}) & \xrightarrow{F_{\rho\tau^{-1}\tau\sigma^{-1};f\rho^{-1}}} & F(f\rho^{-1})\rho\tau^{-1}\tau\sigma^{-1}.
 \end{array} \tag{25}$$

On the other hand, if $\alpha: f \rightarrow g$ and $\beta: g \rightarrow h$ are 2-cells in $\mathcal{M}(\langle a \rangle; b)$, and $\sigma \in \Sigma_n$ we have that

$$\phi(F)(\beta, 1_\sigma) \circ \phi(F)(\alpha, 1_\sigma) = \phi(F)(\beta\alpha, 1_\sigma). \tag{26}$$

The functoriality of $\phi(F)$ follows from a straightforward argument by (24), and (26) together with the exchange property (22).

The next step is to prove that the component functors give rise to a symmetric **Cat**-multifunctor $\phi(F): \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$. First, notice that $\phi(F)$ preserves units since, for $a \in \text{Ob}(\mathcal{M})$ $\phi(F)(1_a, \text{id}_1) = F(1_a \text{id}_1^{-1}) \text{id}_1 = F(1_a) = 1_{Fa}$, since F itself preserves units. Next we prove that $\phi(F)$ preserves the Σ_n -action. For $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$, and $\sigma \in \Sigma_n$, we show that the following diagram commutes in **Cat** :

$$\begin{array}{ccc}
 \mathcal{M}(\langle a \rangle; b) \times E\Sigma_n & \xrightarrow{\phi(F)} & \mathcal{N}(\langle Fa \rangle; Fb) \\
 \sigma \downarrow & & \downarrow \sigma \\
 \mathcal{M}(\langle a_j \rangle \sigma; b) \times E\Sigma_n & \xrightarrow{\phi(F)} & \mathcal{N}(\langle Fa \rangle \sigma; Fb).
 \end{array}$$

For this we don't need any of the pseudo symmetry axioms for F . For 1-cells $(f: \langle a \rangle \rightarrow b, \tau)$ of $\mathcal{M}(\langle a \rangle; b) \times E\Sigma_n$,

$$\begin{aligned} \phi(F)(f, \tau)\sigma &= (F(f\tau^{-1})\tau)\sigma \\ &= F(f\tau^{-1})\tau\sigma \\ &= F(f\sigma(\tau\sigma)^{-1})\tau\sigma \\ &= \phi(F)((f\sigma, \tau\sigma)) \\ &= \phi(F)((f, \tau)\sigma). \end{aligned}$$

A similar calculation works for 2-cells of the form $(\alpha: f \rightarrow g, 1_\tau)$ in $\mathcal{M}(\langle a \rangle; b) \times E\Sigma_n$. For morphisms of the form $(1_f, E_\tau^\rho)$ in $\mathcal{M}(\langle a \rangle; b) \times E\Sigma_n$,

$$\begin{aligned} (\phi(F)(1_f, E_\tau^\rho))\sigma &= (F_{\rho\tau^{-1}; f\rho^{-1}\tau})\sigma \\ &= F_{\rho\tau^{-1}; f\rho^{-1}}(\tau\sigma) \\ &= F_{\rho\sigma(\tau\sigma)^{-1}; f\sigma(\rho\sigma)^{-1}}(\tau\sigma) \\ &= \phi(F)(1_{f\sigma}, E_{\tau\sigma}^{\rho\sigma}) \\ &= \phi(F)((1_f, E_\tau^\rho)\sigma). \end{aligned}$$

By functoriality of $\phi(F)$ and σ we conclude that $\phi(F)$ preserves the action of the symmetric group.

The only step we are missing to finish proving that $\phi(F)$ defines a **Cat**-multifunctor is the preservation of γ . Let $c \in \text{Ob}(\mathcal{M})$, $n \geq 0$, $\langle b \rangle \in \text{Ob}(\mathcal{M})^n$, $k_j \geq 0$ for $1 \leq j \leq n$, $\langle a_j \rangle = \langle a_{j,i} \rangle_{i=1}^{k_j}$ for $1 \leq j \leq n$. Set $k = \sum_{j=1}^n k_j$. As usual $\langle a \rangle = \langle a_j \rangle = \langle \langle a_{j,i} \rangle_{i=1}^{k_j} \rangle_{j=1}^n$ denotes the concatenation of the a_j 's. We will prove that the following square is commutative:

$$\begin{array}{ccc} \mathcal{M}(\langle b \rangle; c) \times E\Sigma_n \times \prod_{j=1}^n \mathcal{M}(\langle a_j \rangle; b_j) \times E\Sigma_{k_j} & \xrightarrow{\phi(F) \times \prod \phi(F)} & \mathcal{N}(\langle Fb \rangle; Fc) \times \prod_{j=1}^n \mathcal{N}(\langle Fa_j \rangle; Fb_j) \\ \downarrow \gamma & & \downarrow \gamma \\ \mathcal{M}(\langle a \rangle; c) \times E(\Sigma_k) & \xrightarrow{\phi(F)} & \mathcal{N}(\langle Fa \rangle; Fc). \end{array} \tag{27}$$

The commutativity of this diagram at the level of 1-cells will follow from top and bottom equivariance for \mathcal{M} and Σ_* , as well as the fact that F preserves γ . Let $f: \langle b \rangle \rightarrow c$, $\sigma \in \Sigma_n$, and $g_j: \langle a_j \rangle \rightarrow b_j$ and $\tau_j \in \Sigma_{k_j}$ for $1 \leq j \leq n$. We have that

$$\begin{aligned} \gamma(\phi(F)(f, \sigma), \langle \phi(F)(g_j, \tau_j) \rangle) &= \gamma(F(f\sigma^{-1})\sigma, \langle F(g_j\tau_j^{-1})\tau_j \rangle) \\ &= \gamma\left(\left(F(f\sigma^{-1}), \left\langle F\left(g_{\sigma^{-1}(j)}\tau_{\sigma^{-1}(j)}^{-1}\right)\right\rangle\right)\right) \sigma\langle \tau_j \rangle \\ &= F\left(\gamma\left(f\sigma^{-1}, \left\langle g_{\sigma^{-1}(j)}\tau_{\sigma^{-1}(j)}^{-1}\right\rangle\right)\right) \sigma\langle \tau_j \rangle \\ &= F\left(\gamma(f, \langle g_j \rangle)(\sigma\langle \tau_j \rangle)^{-1}\right) \sigma\langle \tau_j \rangle \\ &= \phi(F)(\gamma(f, \langle g_j \rangle), \sigma\langle \tau_j \rangle) \\ &= \phi(F)(\gamma((f, \sigma), \langle g_j, \tau_j \rangle)). \end{aligned}$$

We have proven that our diagram is commutative at the level of 1-cells. For the morphisms we will consider again morphisms that change the first variable only and morphisms that change the second variable only separately.

For 2-cells that change the first variable only, the commutativity of our diagram follows in the same way as it did for 1-cells. We consider two cases for 2-cells that change the second variable. For 2-cells of the form $((1_f, E_\sigma^\tau), \langle 1_{g_j}, 1_{\rho_j} \rangle)$ where $f: \langle b \rangle \rightarrow c$, $\sigma, \tau \in \Sigma_n$, and $g_j \in \text{Ob}(\mathcal{M}(\langle a_j \rangle; b_j))$ and $\rho_j \in \Sigma_{k_j}$ for $1 \leq j \leq n$, we have that

$$\begin{aligned} & \gamma \left(\phi(F)(1_f, E_\sigma^\tau) \left\langle \phi(F)(1_{g_j}, 1_{\rho_j}) \right\rangle \right) \\ &= \gamma \left((F_{\tau\sigma^{-1}; f\tau^{-1}})\sigma, \left\langle 1_{F(g_j\rho_j^{-1})\rho_j} \right\rangle \right) \\ &= \gamma \left(F_{\tau\sigma^{-1}; f\tau^{-1}}, \left\langle 1_{F(g_{\sigma^{-1}(j)}\rho_{\sigma^{-1}(j)}^{-1})} \right\rangle \right) \sigma \langle \rho_j \rangle \\ &= F_{\tau\sigma^{-1} \langle \text{id}_{k_{\sigma^{-1}(j)}} \rangle; \gamma(f\tau^{-1} \langle g_{\tau^{-1}(j)}\rho_{\tau^{-1}(j)}^{-1} \rangle)} \sigma \langle \rho_j \rangle \\ &= F_{\tau \langle \rho_j \rangle (\sigma \langle \rho_j \rangle)^{-1}; \gamma(f, \langle g_j \rangle) (\tau \langle \rho_j \rangle)^{-1}} \sigma \langle \rho_j \rangle \\ &= \phi(F)(1_{\gamma(f, \langle g_j \rangle)}, E_{\sigma \langle \rho_j \rangle}^\tau) \\ &= \phi(F)(\gamma(1_f, \langle 1_{g_j} \rangle), \gamma(E_\sigma^\tau, \langle 1_{\rho_j} \rangle)). \end{aligned}$$

The above equalities follow from our definitions, top and bottom equivariance in \mathcal{M}, \mathcal{N} , and $E\Sigma_*$ except the third equality which follows from top equivariance for F (12). Next, let's consider two cells of the form $((1_f, 1_\sigma), \langle 1_{g_j}, E_{\rho_j}^{\nu_j} \rangle)$ where $f: \langle b \rangle \rightarrow c$, $\sigma \in \Sigma_n$, and $g_j \in \text{Ob}(\mathcal{M}(\langle a_j \rangle; b_j))$ and $\rho_j, \nu_j \in \Sigma_{k_j}$ for $1 \leq j \leq n$. We get that

$$\begin{aligned} & \gamma \left(\phi(F)(1_f, 1_\sigma), \phi(F) \left\langle (1_{g_j}, E_{\rho_j}^{\nu_j}) \right\rangle \right) \\ &= \gamma \left(1_{F(f\sigma^{-1})\sigma}, \left(F_{\nu_j\rho_j^{-1}; g_j\nu_j^{-1}} \right) \rho_j \right) \\ &= \gamma \left(1_{F(f\sigma^{-1})}, \left\langle F_{\nu_{\sigma^{-1}(j)}\rho_{\sigma^{-1}(j)}^{-1}; g_{\sigma^{-1}(j)}\nu_{\sigma^{-1}(j)}^{-1}} \right\rangle \right) \sigma \langle \rho_j \rangle \\ &= F_{\text{id}_n \langle \nu_{\sigma^{-1}(j)}\rho_{\sigma^{-1}(j)}^{-1} \rangle; \gamma(f\sigma^{-1}, \langle g_{\sigma^{-1}(j)}\nu_{\sigma^{-1}(j)}^{-1} \rangle)} \sigma \langle \rho_j \rangle \\ &= F_{\sigma \langle \nu_j \rangle (\sigma \langle \rho_j \rangle)^{-1}; \gamma(f, \langle g_j \rangle) (\rho \langle \nu_j \rangle)^{-1}} \sigma \langle \rho_j \rangle \\ &= \phi(F) \left(1_{\gamma(f, \langle g_j \rangle)}, \left\langle E_{\sigma \langle \rho_j \rangle}^{\sigma \langle \nu_j \rangle} \right\rangle \right) \\ &= \phi(F) \left(\gamma \left((1_f, 1_\sigma), \left\langle (1_{g_j}, E_{\rho_j}^{\nu_j}) \right\rangle \right) \right). \end{aligned}$$

The third equality above follows from the bottom equivariance axiom for F (13) and the rest by our definitions as well as top and bottom equivariance for \mathcal{M}, \mathcal{N} , and $E\Sigma_*$.

By functoriality of γ and $\phi(F)$, and since every morphism in the source category can be written as a composite of arrows for which we already proved the commutativity of (27), we can conclude that the square (27) is commutative.

We are almost done, we just have to prove that our definition of $\phi(F)$ gives us $F = \phi(F) \circ \eta$ in **Cat-Multicat**^{ps}. This is clear for objects of the multicategory \mathcal{M} . For each $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in \text{Ob}(\mathcal{M})$, and $f: \langle a \rangle \rightarrow b$,

$$\phi(F) \circ \eta(f) = \phi(F)(f, \text{id}_n) = F(f \text{id}_n^{-1}) \text{id}_n = F(f).$$


Similarly for $\alpha: f \rightarrow g$ a 2-cell in $\mathcal{M}(\langle a \rangle; b)$. Finally, we just need to prove that $(\phi(F) \circ \eta)_{\sigma, \langle a_i \rangle, b} = F_{\sigma, \langle a_i \rangle, b}$ for any $\sigma \in \Sigma_n$. Let $f: \langle a \rangle \rightarrow b$ be a 1-cell. Since $\phi(F)$ is symmetric,

$$(\phi(F)\eta)_{\sigma;f} = \phi(F)(\eta_{\sigma;f}) = \phi(F)(1_{f\sigma}, E_{\text{id}^\sigma}) = F_{\sigma(\text{id})^{-1};f\sigma\sigma^{-1}} = F_{\sigma;f},$$

where we have used the notation introduced just before (9). We have proven that $j\phi(F) \circ \eta = F$. This finishes our proof. ■

Similarly, pseudo symmetric **Cat**-multinatural transformations between F and G correspond to symmetric **Cat**-multinatural transformations between $\phi(F)$ and $\phi(G)$.

3.4. LEMMA. *Let \mathcal{M}, \mathcal{N} be **Cat**-multicategories with $F, G: \mathcal{M} \rightarrow \mathcal{N}$ pseudo symmetric **Cat**-multifunctors and $\theta: F \rightarrow G$ a pseudo symmetric **Cat**-multinatural transformation. There exists a unique symmetric **Cat**-multinatural transformation $\phi(\theta): \phi(F) \rightarrow \phi(G)$ such that $\phi(\theta) * 1_{\eta_{\mathcal{M}}} = \theta$ in **Cat-Multicat**^{ps}. That is, the following pasting diagram equality holds in **Cat-Multicat**^{ps} :*



PROOF. We prove uniqueness first. Suppose $\phi(\theta)$ is a symmetric **Cat**-multinatural transformation $\phi(\theta): \phi(F) \rightarrow \phi(G)$ such that $\phi(\theta) * 1_\eta = \theta$. Any object of $\mathcal{M} \times E\Sigma_*$ takes the form $(a, *)$ for some object a of \mathcal{M} , with $i(a) = (a, *)$. By definition,

$$\theta_a = \gamma(\phi(\theta)_{\eta a}, \phi(F)((1_\eta)_a)) = \gamma(\phi(\theta)_{\eta a}, 1_{Fa}) = \phi(\theta)_{\eta a}.$$

Since all objects of the **Cat**-multifunctor $\mathcal{M} \times E\Sigma_*$ are of the form ηa for some object a of \mathcal{M} , this is the only possible way of defining such **Cat**-multinatural transformation $\phi(\theta)$. Next, we check that by defining $\phi(\theta)_{(a,*)} = \theta_a$ for $a \in \text{Ob}(\mathcal{M})$, we in fact get a symmetric **Cat**-multinatural transformation $\phi(\theta): \phi(F) \rightarrow \phi(G)$. Let $n \geq 0$, $\langle a \rangle \in \text{Ob}(\mathcal{M})^n$, $b \in$

$(\text{Ob}(\mathcal{M})^n)$, $f: \langle a \rangle \rightarrow b$, and $\sigma \in \Sigma_n$, then

$$\begin{aligned} \gamma(\phi(G)(f, \sigma); \langle \phi(\theta)_{(a_j, *)} \rangle) &= \gamma(G(f\sigma^{-1})\sigma; \langle \theta_{a_j} \rangle) \\ &= \gamma\left(G(f\sigma^{-1}); \left\langle \theta_{a_{\sigma^{-1}(j)}} \right\rangle\right) \sigma \\ &= \gamma(\theta_b; F(f\sigma^{-1}))\sigma \\ &= \gamma(\theta_b; F(f\sigma^{-1})\sigma) \\ &= \gamma(\phi(\theta)_{(b, *)}, \phi(F)(f, \sigma)) \end{aligned}$$

Where we have used top and bottom equivariance, as well as the **Cat**-multinaturality of θ . Now we need to prove **Cat**-multinaturality of $\phi(\theta)$ for 2-cells. As before, the case where the 2-cell changes just the first variable is similar to what was done for 1-cells. Now, if $\langle a \rangle, b, f$ are as before and E_σ^τ is a morphism in $E\Sigma_n$, $(1_f, E_\sigma^\tau)$ is a morphism in $\mathcal{M}(\langle a \rangle; b) \times E\Sigma_n$, and

$$\begin{aligned} \gamma\left(\phi(G)(1_f, E_\sigma^\tau); \left\langle 1_{\phi(\theta)_{(a_j, *)}} \right\rangle\right) &= \gamma\left((G_{\tau\sigma^{-1}; f\tau^{-1}})\sigma; \langle 1_{\theta_{a_j}} \rangle\right) \\ &= \gamma\left(G_{\tau\sigma^{-1}; f\tau^{-1}}; \left\langle 1_{\theta_{a_{\sigma^{-1}(j)}}} \right\rangle\right) \sigma \\ &= \gamma(1_{\theta_b}; F_{\tau\sigma^{-1}; f\tau^{-1}}) \sigma \\ &= \gamma\left(1_{\phi(\theta)_{(b, *)}}; \phi(F)(1_f, E_\sigma^\tau)\right). \end{aligned}$$

In the third equality we have used pseudo symmetric **Cat**-multinaturality for θ . In conclusion, by componentwise functoriality of $\gamma, \phi(F)$ and $\phi(G)$ we conclude that **Cat**-multinaturality holds for $\phi(\theta)$ at the 2-cell level finishing the proof of the lemma. ■

Furthermore, Theorem 3.3 and Lemma 3.4 together give the following isomorphism.

3.5. COROLLARY. *If \mathcal{M}, \mathcal{N} are **Cat** multicategories, then there is an isomorphism of small categories*

$$\mathbf{Cat}\text{-Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N}) \cong \mathbf{Cat}\text{-Multicat}(\mathcal{M} \times E\Sigma_*, \mathcal{N}).$$

PROOF. Recalling the definitions from the two previous results, we define

$$\phi: \mathbf{Cat}\text{-Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N}) \longrightarrow \mathbf{Cat}\text{-Multicat}(\mathcal{M} \times E\Sigma_*, \mathcal{N}) \tag{28}$$

for pseudo symmetric **Cat**-multifunctors as in Theorem 3.3 and for pseudo symmetric **Cat**-multinatural transformations as in Lemma 3.4.

It is immediate from the definitions that ϕ is a functor. Indeed, if $\alpha: F \rightarrow G$ and $\beta: G \rightarrow H$ are pseudo symmetric **Cat**-multinatural transformations with $F, G, H: \mathcal{M} \rightarrow \mathcal{N}$

$$\phi(\beta * \alpha)_{(c, *)} = (\beta * \alpha)_c = \gamma(\beta_c, \alpha_c) = \gamma(\phi(\beta)_{(c, *)}, \phi(\alpha)_{(c, *)}) = (\phi(\beta) * \phi(\alpha))_{(c, *)}$$

We can define the inverse of ϕ, η^* , as the composite

$$\begin{array}{ccc}
 \mathbf{Cat}\text{-}\mathbf{Multicat}(\mathcal{M} \times E\Sigma_*, \mathcal{N}) & \xrightarrow{j} & \mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}(\mathcal{M} \times E\Sigma_*, \mathcal{N}) \\
 \searrow \eta^* & & \downarrow \eta_{\mathcal{M}}^* \\
 & & \mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N}).
 \end{array} \tag{29}$$

Finally, the existence part of Theorem 3.3 and Lemma 3.4, implies that $\eta^* \circ \phi$ is the identity of $\mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N})$, while the uniqueness part of both results implies that $\phi \circ \eta^*$ is the identity of $\mathbf{Cat}\text{-}\mathbf{Multicat}(\mathcal{M} \times E\Sigma_*, \mathcal{N})$. ■

The two previous results hint at the existence of a 2-adjunction between the 2-inclusion $j: \mathbf{Cat}\text{-}\mathbf{Multicat} \rightarrow \mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}$ and the 2-functor which we define next.

3.6. DEFINITION. We define the 2-functor $\psi: \mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}} \rightarrow \mathbf{Cat}\text{-}\mathbf{Multicat}$ as follows. For a **Cat**-multicategory \mathcal{M} , $\psi\mathcal{M} = \mathcal{M} \times E\Sigma_*$. For \mathcal{M}, \mathcal{N} **Cat**-multicategories, we define the component functor ψ as the composite

$$\begin{array}{ccc}
 \mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N}) & \xrightarrow{\eta_{\mathcal{N}}^*} & \mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N} \times E\Sigma_*) \\
 \searrow \psi & & \downarrow \phi \\
 & & \mathbf{Cat}\text{-}\mathbf{Multicat}(\mathcal{M} \times E\Sigma_*, \mathcal{N} \times E\Sigma_*).
 \end{array}$$

Thus, by Theorem 3.3 if $F: \mathcal{M} \rightarrow \mathcal{N}$ is a pseudo symmetric **Cat**-multifunctor, then $\psi F: \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N} \times E\Sigma_*$ is the unique symmetric **Cat**-multifunctor which makes the diagram

$$\begin{array}{ccc}
 \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\
 F \downarrow & & \downarrow j\psi F \\
 \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_*
 \end{array} \tag{30}$$

commute in $\mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}$. Similarly, by Lemma 3.4, for $\theta: F \rightarrow G$ a pseudo symmetric **Cat**-multinatural transformation between $F, G: \mathcal{M} \rightarrow \mathcal{N}$ pseudo symmetric **Cat**-multifunctors, $\psi\theta: \psi F \rightarrow \psi G$ is the unique symmetric **Cat**-multinatural transformation such that the equality of pasting diagrams

$$\begin{array}{ccc}
 \begin{array}{ccc}
 \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\
 \left(\begin{array}{ccc} \theta & \longrightarrow & \end{array} \right) & & \downarrow j\psi F \\
 \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_*
 \end{array} & = & \begin{array}{ccc}
 \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\
 F \downarrow & & j\psi F \left(\begin{array}{ccc} j\psi\theta & \longrightarrow & \end{array} \right) j\psi G \\
 \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_*
 \end{array}
 \end{array} \tag{31}$$

holds in $\mathbf{Cat}\text{-}\mathbf{Multicat}^{\text{ps}}$.

3.7. THEOREM. *There is a 2-adjunction*

$$\text{Cat-Multicat}^{\text{ps}} \begin{array}{c} \xrightarrow{\psi} \\ \perp \\ \xleftarrow{j} \end{array} \text{Cat-Multicat}$$

where j is the inclusion 2-functor.

PROOF. Following Corollary 3.5, we define the unit of the adjunction as the strict 2-natural transformation $\eta: 1_{\text{Cat-Multicat}^{\text{ps}}} \rightarrow j\psi$ having component $\eta_{\mathcal{M}}$ at a **Cat**-multicategory \mathcal{M} . We also define the counit of the adjunction $\pi: \psi j \rightarrow 1_{\text{Cat-Multicat}}$ as having component at a **Cat**-multicategory \mathcal{M} the projection $\pi_{\mathcal{M}}: \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{M}$.

The fact that η defines a strict 2-natural transformation follows directly from (30) and (31). To prove that the data of π defines a strict 2-natural transformation we need to prove that given $F: \mathcal{M} \rightarrow \mathcal{N}$ symmetric **Cat**-multifunctor, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{M} \times E\Sigma_* & \xrightarrow{\pi_{\mathcal{M}}} & \mathcal{M} \\ \psi j F \downarrow & & \downarrow F \\ \mathcal{N} \times E\Sigma_* & \xrightarrow{\pi_{\mathcal{N}}} & \mathcal{N}. \end{array}$$

Indeed, we prove that $\psi j F = F \times 1_{E\Sigma_*}$. By (30), it suffices to show that the following diagram commutes in **Cat-Multicat**^{ps}:

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\ j F \downarrow & & \downarrow j(F \times 1) \\ \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_*. \end{array} \tag{32}$$

It is clear that this diagram commutes at the level of objects, 1-cells, and 2-cells of the multicategory. The pseudo symmetry isomorphisms of both composites also agree. Indeed, for $f: \langle a \rangle \rightarrow b$ a 1-cell of \mathcal{M} and $\sigma \in \Sigma_n$, by (14), we get that

$$\begin{aligned} (j(F \times 1)\eta_{\mathcal{M}})_{\sigma;f} &= j(F \times 1)_{\sigma;\eta_{\mathcal{M}}(f)} \circ j(F \times 1)(\eta_{\mathcal{M}\sigma;f}) \\ &= (1_{(Ff)\sigma}, 1_{\sigma}) \circ (1_{(Ff)\sigma}, E_{\text{id}}^{\sigma}) \\ &= (1_{(Ff)\sigma}, E_{\text{id}}^{\sigma}) \circ (1_{(Ff)\sigma}, 1_{\sigma}) \\ &= \eta_{\mathcal{N}\sigma;Ff} \circ \eta_{\mathcal{N}}(jF_{\sigma;f}) \\ &= (\eta_{\mathcal{N}} \circ jF)_{\sigma;f}. \end{aligned}$$

To finish proving the 2-naturality of $\pi_{\mathcal{M}}$, we need to prove that given \mathcal{M}, \mathcal{N} **Cat**-multicategories, $F, G: \mathcal{M} \rightarrow \mathcal{N}$ **Cat**-multifunctors and a **Cat**-multinatural transforma-

tion $\theta: F \longrightarrow G$, the following equality of pasting diagrams holds in **Cat-Multicat**:

$$\begin{array}{ccc}
 \mathcal{M} \times E\Sigma_* & \xrightarrow{\pi_{\mathcal{M}}} & \mathcal{M} \\
 \downarrow j(F \times 1) \left(\begin{array}{c} \xrightarrow{\psi j\theta} \\ \xrightarrow{\psi j\theta} \end{array} \right) j(G \times 1) & & \downarrow G \\
 \mathcal{N} \times E\Sigma_* & \xrightarrow{\pi_{\mathcal{N}}} & \mathcal{N}
 \end{array} = \begin{array}{ccc}
 \mathcal{M} \times E\Sigma_* & \xrightarrow{\pi_{\mathcal{M}}} & \mathcal{M} \\
 \downarrow j(F \times 1) & & \downarrow F \left(\begin{array}{c} \xrightarrow{\theta} \\ \xrightarrow{\theta} \end{array} \right) G \\
 \mathcal{N} \times E\Sigma_* & \xrightarrow{\pi_{\mathcal{N}}} & \mathcal{N}
 \end{array}$$

In turn, the last equality of pasting diagrams holds since $\psi j\theta = j(\theta \times 1)$. To see this, by (31), we must show the following equality of pasting diagrams in **Cat-Multicat^{PS}** :

$$\begin{array}{ccc}
 \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\
 \downarrow jF \left(\begin{array}{c} \xrightarrow{j\theta} \\ \xrightarrow{j\theta} \end{array} \right) jG & & \downarrow j(G \times 1) \\
 \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_*
 \end{array} = \begin{array}{ccc}
 \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\
 \downarrow jF & & \downarrow j(F \times 1) \left(\begin{array}{c} \xrightarrow{j(\theta \times 1)} \\ \xrightarrow{j(\theta \times 1)} \end{array} \right) j(G \times 1) \\
 \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_*
 \end{array} \tag{33}$$

To check that this equality holds let $a \in \text{Ob}(\mathcal{M})$. We get, by (8), that

$$\begin{aligned}
 (1_{\eta_{\mathcal{N}}} * j\theta)_a &= \gamma(1_{\eta_{\mathcal{N}}(jGa)}; \eta_{\mathcal{N}}(\theta_a)) \\
 &= \gamma((1_{Ga}, 1_{\text{id}}); (\theta_a, 1_{\text{id}})) \\
 &= \gamma((\theta_a, 1_{\text{id}}); (1_{Fa}, 1_{\text{id}})) \\
 &= \gamma(j(\theta \times 1)_{\eta_{\mathcal{N}}(a)}; j(F \times 1)(1_{\eta_{\mathcal{M}}(a)})) \\
 &= (j(\theta \times 1) * \eta_{\mathcal{M}})_a.
 \end{aligned}$$

Thus, η and π are strict 2-natural transformations and we just need to prove that they satisfy the triangle identities. To prove that the identity $(1_j * \pi)(\eta * 1_j) = 1_j$ holds we need to prove that for \mathcal{M} a **Cat-multicategory** the diagram

$$\begin{array}{ccc}
 & \mathcal{M} \times E\Sigma_* & \\
 \eta_{\mathcal{M}} \nearrow & & \searrow j\pi_{\mathcal{M}} \\
 \mathcal{M} & \xrightarrow{1_{\mathcal{M}}} & \mathcal{M}
 \end{array}$$

commutes in **Cat-Multicat^{PS}**. This is clear at the level of objects, n -ary 1-cells and n -ary 2-cells. The pseudo symmetry isomorphisms of both pseudo symmetric **Cat-multifunctors** also agree since, for $f: \langle a \rangle \longrightarrow b$ an n -ary 1-cell of \mathcal{M} and $\sigma \in \Sigma_n$, we obtain, by (14),

$$((j\pi_{\mathcal{M}}) \circ \eta_{\mathcal{M}})_{\sigma;f} = (j\pi_{\mathcal{M}})_{\sigma; \eta_{\mathcal{M}}(f)} \circ j\pi_{\mathcal{M}}(\eta_{\mathcal{M}\sigma;f}) = 1_{f\sigma} = 1_{\mathcal{M}\sigma;f}.$$

The other triangle identity is $(\pi * 1_{\psi})(1_{\psi} * \eta) = 1_{\psi}$. To check it, we must prove that, given a **Cat-multicategory** \mathcal{M} , the composite

$$\mathcal{M} \times E\Sigma_* \xrightarrow{\psi \eta_{\mathcal{M}}} \mathcal{M} \times E\Sigma_* \times E\Sigma_* \xrightarrow{\pi_{\mathcal{M} \times E\Sigma_*}} \mathcal{M} \times E\Sigma_*$$

agrees with $1_{\mathcal{M} \times E\Sigma_*}$. This holds since, if $\Delta: E\Sigma_* \rightarrow E\Sigma_* \times E\Sigma_*$ denotes the diagonal map, then $\psi(\eta_{\mathcal{M}}) = 1_{\mathcal{M}} \times \Delta$. To see this, notice that by (30) all we need is to prove that the following diagram is commutative:

$$\begin{array}{ccc}
 \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\
 \eta_{\mathcal{M}} \downarrow & & \downarrow j(1 \times \Delta) \\
 \mathcal{M} \times E\Sigma_* & \xrightarrow{\eta_{\mathcal{M} \times E\Sigma_*}} & \mathcal{M} \times E\Sigma_* \times E\Sigma_*.
 \end{array} \tag{34}$$

Now, the previous diagram is evidently commutative at the level of objects, 1-cells, and 2-cells. The diagram also commutes at the level of pseudo symmetry isomorphisms since, for $f: \langle a \rangle \rightarrow b$ an n -ary 1-cell in \mathcal{M} and $\sigma \in \Sigma_n$,

$$\begin{aligned}
 (\eta_{\mathcal{M} \times E\Sigma_*} \circ \eta_{\mathcal{M}})_{\sigma;f} &= \eta_{\mathcal{M} \times E\Sigma_*; \sigma; \eta_{\mathcal{M}}(f)} \circ \eta_{\mathcal{M} \times E\Sigma_*}(\eta_{\mathcal{M}\sigma;f}) \\
 &= (1_{f\sigma}, 1_{\sigma}, E_{\text{id}}^{\sigma}) \circ (1_{f\sigma}, E_{\text{id}}^{\sigma}, 1_{\text{id}}) \\
 &= (1_{f\sigma}, 1_{\sigma}, 1_{\sigma}) \circ (1_{f\sigma}, E_{\text{id}}^{\sigma}, E_{\text{id}}^{\sigma}) \\
 &= j(1 \times \Delta)_{\sigma; \eta_{\mathcal{M}}(f)} \circ j(1 \times \Delta)(\eta_{\mathcal{M}\sigma;f}) \\
 &= (j(1 \times \Delta) \circ \eta_{\mathcal{M}})_{\sigma;f}.
 \end{aligned}$$

We conclude that the triangle identities are satisfied and thus we get the desired 2-adjunction. ■

We can use this 2-adjunction to describe the 2-category **Cat-Multicat**^{ps} in terms of symmetric **Cat**-multifunctors and symmetric **Cat**-multinatural transformations alone, thus upgrading the functors ϕ from Corollary 3.5 to an isomorphism of 2-categories.

3.8. DEFINITION. *The 2-category **D** has **Cat**-multicategories as objects. For \mathcal{M}, \mathcal{N} **Cat**-multicategories, the category of morphisms between \mathcal{M} and \mathcal{N} is*

$$\mathbf{D}(\mathcal{M}, \mathcal{N}) = \mathbf{Cat}\text{-}\mathbf{Multicat}(\mathcal{M} \times E\Sigma_*, \mathcal{N}).$$

*In particular, vertical composition of 2-cells is defined as in **Cat-Multicat**. For $F: \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$ and $G: \mathcal{N} \times E\Sigma_* \rightarrow \mathcal{Q}$ symmetric **Cat**-multifunctors, the composition $G \circ F$ is defined as the composite*

$$\mathcal{M} \times E\Sigma_* \xrightarrow{1 \times \Delta} \mathcal{M} \times E\Sigma_* \times E\Sigma_* \xrightarrow{F \times 1} \mathcal{N} \times E\Sigma_* \xrightarrow{G} \mathcal{Q}$$

*in **Cat-Multicat**. Similarly, for $F, J: \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$, $G, K: \mathcal{N} \times E\Sigma_* \rightarrow \mathcal{Q}$ symmetric **Cat**-multifunctors and $\theta: F \rightarrow J$, $\zeta: G \rightarrow K$ **Cat**-multinatural transformations, $\zeta * \theta$ is defined as the pasting*

$$\begin{array}{ccccc}
 \mathcal{M} \times E\Sigma_* & \xrightarrow{1 \times \Delta} & \mathcal{M} \times E\Sigma_* \times E\Sigma_* & \xrightarrow{F \times 1} & \mathcal{N} \times E\Sigma_* & \xrightarrow{G} & \mathcal{Q} \\
 & & \Downarrow \theta \times 1 & & \Downarrow \zeta & & \\
 & & \xrightarrow{J \times 1} & & \xrightarrow{K} & &
 \end{array}$$

*in **Cat-Multicat**.*

The previous definition makes **D** into a 2-category and the functors ϕ , and η^* from Corollary 3.5 into the components of isomorphisms of 2-categories.

3.9. THEOREM. *The data of the previous definition defines a 2-category \mathbf{D} isomorphic to $\mathbf{Cat}\text{-Multicat}^{\text{ps}}$.*

PROOF. The (horizontal) composition functors are defined so that ϕ and η^* become the componentwise functors of a 2-category isomorphism between \mathbf{D} and $\mathbf{Cat}\text{-Multicat}^{\text{ps}}$. More precisely, for \mathcal{M}, \mathcal{N} and \mathcal{Q} \mathbf{Cat} -multicategories, we will prove that the \mathbf{D} composition functor defined, $\circ' : \mathbf{D}(\mathcal{N}, \mathcal{Q}) \times \mathbf{D}(\mathcal{M}, \mathcal{N}) \rightarrow \mathbf{D}(\mathcal{M}, \mathcal{Q})$, makes the following diagram commute, where \circ denotes the horizontal composition functor of $\mathbf{Cat}\text{-Multicat}^{\text{ps}}$:

$$\begin{array}{ccc} \mathbf{D}(\mathcal{N}, \mathcal{Q}) \times \mathbf{D}(\mathcal{M}, \mathcal{N}) & \xrightarrow{\circ'} & \mathbf{D}(\mathcal{M}, \mathcal{Q}) \\ \eta^* \times \eta^* \downarrow & & \uparrow \phi \end{array} \tag{35}$$

$$\mathbf{Cat}\text{-Multicat}^{\text{ps}}(\mathcal{N}, \mathcal{Q}) \times \mathbf{Cat}\text{-Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{N}) \xrightarrow{\circ} \mathbf{Cat}\text{-Multicat}^{\text{ps}}(\mathcal{M}, \mathcal{Q}).$$

Let $G : \mathcal{N} \times \mathcal{Q}$ and $F : \mathcal{M} \times E\Sigma_* \rightarrow \mathcal{N}$ be symmetric \mathbf{Cat} -multifunctors. The commutativity of (35) for (G, F) reduces to the commutativity of the following diagram by Theorem 3.3:

$$\begin{array}{ccccc} \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* & & \\ \eta_{\mathcal{M}} \downarrow & & \downarrow j(1 \times \Delta) & & \\ \mathcal{M} \times E\Sigma_* & \xrightarrow{\eta_{\mathcal{M} \times E\Sigma_*}} & \mathcal{M} \times E\Sigma_* \times E\Sigma_* & & \\ jF \downarrow & & \downarrow j(F \times 1) & & \\ \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_* & \xrightarrow{jG} & \mathcal{Q}. \end{array}$$

This diagram in turn is commutative by (32) and (34). Now, if F, G are as before, $J : \mathcal{M} \times E\Sigma_*$ and $K : \mathcal{N} \times E\Sigma_* \rightarrow \mathcal{Q}$ are symmetric \mathbf{Cat} -multifunctors, and $\theta : F \rightarrow J$, $\zeta : G \rightarrow K$ are \mathbf{Cat} -multinatural transformations, by Lemma 3.4, the commutativity of (35) for (ζ, θ) reduces to the equality of pasting diagrams:

$$\begin{array}{ccc} \begin{array}{ccc} \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\ \eta_{\mathcal{M}} \downarrow & & \downarrow j(1 \times \Delta) \\ \mathcal{M} \times E\Sigma_* & \xrightarrow{\eta_{\mathcal{M} \times E\Sigma_*}} & \mathcal{M} \times E\Sigma_* \times E\Sigma_* \\ jF \downarrow & & \downarrow j(F \times 1) \\ \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_* \\ & & \downarrow jG \\ & & \mathcal{Q} \end{array} & = & \begin{array}{ccc} \mathcal{M} & \xrightarrow{\eta_{\mathcal{M}}} & \mathcal{M} \times E\Sigma_* \\ \eta_{\mathcal{M}} \downarrow & & \downarrow j(1 \times \Delta) \\ \mathcal{M} \times E\Sigma_* & \xrightarrow{\eta_{\mathcal{M} \times E\Sigma_*}} & \mathcal{M} \times E\Sigma_* \times E\Sigma_* \\ jF \downarrow & & \downarrow j(J \times 1) \\ \mathcal{N} & \xrightarrow{\eta_{\mathcal{N}}} & \mathcal{N} \times E\Sigma_* \\ & & \downarrow jG \\ & & \mathcal{Q} \end{array} \end{array}$$

This equality holds by (33) and makes implicit use of (32) and (34). We can thus define $\phi : \mathbf{Cat}\text{-Multicat}^{\text{ps}} \rightarrow \mathbf{D}$ in objects as the identity map, and do the same for

$\eta^* : \mathbf{D} \rightarrow \mathbf{Cat}\text{-Multicat}^{\text{ps}}$, with the component functors given for \mathcal{M} and \mathcal{N} multicategories by (28) and (29) respectively. By (35) and the fact that ϕ and η^* are componentwise isomorphisms, ϕ and η preserve vertical composition of 2-cells and horizontal composition of 1-cells and 2-cells. The fact that $\mathbf{Cat}\text{-Multicat}^{\text{ps}}$ is a 2-category implies that \mathbf{D} is a 2-category. This further turns ϕ and η^* into isomorphisms of 2-categories. ■

4. Applications to inverse K -theory

We use our understanding of pseudo symmetric \mathbf{Cat} -multifunctors to show that they preserve certain E_n -algebras for $n = 1, 2, 3, \dots, \infty$. First we define E_n \mathbf{Cat} -operads.

4.1. DEFINITION. For $n = 1, \dots, \infty$, an E_n \mathbf{Cat} -operad is a \mathbf{Cat} -operad that becomes a topological E_n -operad (in the sense of [May, 1972]) after applying the classifying space functor. A topological E_n -operad is one that has the same Σ -equivariant homotopy type as the little n -cubes operad.

4.2. EXAMPLE. An example of an E_∞ \mathbf{Cat} -operad is $E\Sigma_*$. There are also examples of E_n \mathbf{Cat} -operads for each $n = 1, 2, \dots$ in [Berger, 1996] and [Fiedorowicz et al., 2003], which furthermore have a free action of the symmetric group. Importantly, symmetric algebras over topological E_n -operads are grouplike n -fold loop spaces. Symmetric algebras over the E_n \mathbf{Cat} -operads in [Fiedorowicz et al., 2003] are n -fold monoidal categories, with the group completion of the classifying space of an n -monoidal category being an example of an n -fold loop space.

4.3. DEFINITION. Let \mathcal{M} be a \mathbf{Cat} -multicategory and \mathcal{O} a \mathbf{Cat} -operad. A pseudo symmetric algebra in \mathcal{M} over \mathcal{O} is a pseudo symmetric \mathbf{Cat} -multifunctor $\mathcal{O} \rightarrow \mathcal{M}$. For $n \in \{1, 2, \dots, \infty\}$, a symmetric E_n -algebra (respectively a pseudo symmetric E_n -algebra) in \mathcal{M} is a symmetric algebra (respectively a pseudo symmetric algebra) over an E_n -operad.

1. Let \mathcal{O} be a Σ -free E_n \mathbf{Cat} -operad. Then $\mathcal{O} \times E\Sigma_*$ is an E_n \mathbf{Cat} -operad.
2. Pseudo symmetric E_n -algebras over Σ -free E_n \mathbf{Cat} -operads are symmetric E_n -algebras for $n = 1, 2, \dots, \infty$.

PROOF. Let \mathcal{O} be a Σ -free \mathbf{Cat} -operad. We will show that $\mathcal{O} \times E\Sigma_*$ is componentwise Σ -equivariantly homotopy equivalent to \mathcal{O} (after taking nerves), that is, for each $n \geq 0$, we will show that the projection $\mathcal{O}(n) \times E\Sigma_n \rightarrow \mathcal{O}(n)$ induces a Σ_n -equivariant homotopy equivalence on classifying spaces. Since $B(\mathcal{O}(n) \times E\Sigma_n)$ and $B(\mathcal{O}(n))$ are Σ_n -CW complexes we must show that for subgroups $H \leq \Sigma_n$, the projection induces homotopy equivalences $B(\mathcal{O}(n) \times \Sigma_n)^H \rightarrow B(\mathcal{O}(n))^H$. Since the action of Σ_n on both $\mathcal{O}(n) \times E\Sigma_n$ and $\mathcal{O}(n)$ is free, the fixed point map is either empty when H is non-trivial or the projection $B(\mathcal{O}(n)) \times B(E\Sigma_n) \rightarrow B(\mathcal{O}(n))$, which is a homotopy equivalence since $B(E\Sigma_n)$ is contractible. ■

4.5. **EXAMPLE.** If \mathcal{O} is **Cat**-operad and \mathcal{M} is a **Cat**-multicategory, the pseudo symmetric algebras over \mathcal{O} agree with symmetric algebras over the operad $\mathcal{O} \times E\Sigma_*$. For example, while algebras over the commutative operad $\{*\}$ in \mathcal{M} are the commutative monoids in \mathcal{M} , pseudo symmetric algebras over $\{*\}$ in \mathcal{M} are precisely algebras over the Barratt-Eccles operad and thus, E_∞ -algebras. Similarly, pseudo symmetric algebras over the E_∞ **Cat**-operad $E\Sigma_*$, which are defined in [Yau, 2024] as pseudo symmetric E_∞ -algebras in \mathcal{M} , are algebras over $E\Sigma_* \times E\Sigma_* = E(\Sigma_* \times \Sigma_*)$ which is still an E_∞ **Cat**-operad, and thus, they are still E_∞ -algebras in the sense defined above. Thus, we have the following result.

4.6. **REMARK.** We remind the reader that Σ -freedom is not a serious restriction since there are E_n -operads in **Cat**, like those in [Berger, 1996] and [Fiedorowicz et al., 2003] which are Σ -free. As a corollary, we conclude that pseudo symmetric **Cat**-multifunctors preserve certain E_n -algebras.

4.7. **COROLLARY.** *Let \mathcal{M} and \mathcal{N} be **Cat**-multicategories and $F: \mathcal{M} \rightarrow \mathcal{N}$ be a pseudo symmetric **Cat**-multifunctor, then:*

1. *F sends commutative monoids in \mathcal{M} to E_∞ -algebras in \mathcal{N} .*
2. *F preserves E_n -algebras parameterized by Σ -free **Cat**-operads.*

We conclude our paper by applying our understanding of pseudo symmetric **Cat**-multifunctors to multifunctorial inverse K -theory. In [Johnson and Yau, 2022], Johnson and Yau define Mandell's inverse K -theory multifunctor \mathcal{P} as well as the **Cat**-multicategories that are its domain (Γ -categories) and target (permutative categories). Yau proves in [Yau, 2024] that \mathcal{P} is pseudo symmetric. We refer the interested reader [Yau, 2024] of which the following theorem is one of the main results.

4.8. **THEOREM.** [Yau, 2024] Mandell's inverse K -theory functor is a pseudo symmetric **Cat**-multifunctor $\mathcal{P}: \Gamma\text{-Cat} \rightarrow \text{PermCat}^{\text{sg}}$.

As a consequence, \mathcal{P} sends commutative monoids to E_∞ -algebras and preserves E_n -algebras parameterized by Σ -free E_n -operads, as was stated in Corollary 1.2.

References

- C. Berger. "Opérides cellulaires et espaces de lacet itérés". In: *Annales de l'Institut Fourier* Volume 46 (1996), pp. 1125–1157.
- A. K. Bousfield and E. M. Friedlander. "Homotopy theory of Γ -spaces, spectra, and bisimplicial sets". In: *Geometric applications of homotopy theory* (Proc. Conf., Evanston, Ill., 1977), II. Vol. 658. Lecture Notes in Math. Springer, Berlin-New York, 1978, pp. 80–130.

- C. Balteanu, Z. Fiedorowicz, R. Schwänzl, and R. Vogt. “Iterated monoidal categories.” In: *Adv. Math.* 176 no. 2 (2003), pp. 277–349.
- Andrew J. Blumberg and Michael A. Mandell. “Derived Koszul duality and involutions in the algebraic K-theory of spaces”. In: *J. Topol.* 4 no. 2 (2011), pp. 327–342.
- Denis-Charles Cisinski. “La classe des morphismes de Dwyer n’est pas stable par retracts”. In: *Cahiers Topologie Géom. Différentielle Catég.* 40 no. 3 (1999), pp. 227–231.
- Anthony D. Elmendorf. “Multiplicativity in Mandell’s Inverse KTheory”. Preprint available at <https://arxiv.org/pdf/2110.07512.pdf>. 2021.
- A. D. Elmendorf and M. A. Mandell. “Rings, modules, and algebras in infinite loop space theory”. In: *Adv. Math.* 205 no. 1 (2006), pp. 163–228.
- N. Gurski, N. Johnson, and A. M. Osorno. “The symmetric monoidal 2-category of permutative categories”. In: *High. Struct.* 8.1 (2024), pp. 244–320.
- B. J. Guillou, J. P. May, M. Merling, and A. M. Osorno. “Multiplicative equivariant K-theory and the Barratt-Priddy-Quillen theorem”. In: *Adv. Math.* 414 (2023), Paper No. 108865, 111 p.
- John R. Isbell. “On coherent algebras and strict algebras”. In: *J. Algebra* 13 (1969), pp. 299–307.
- N. Johnson and D. Yau. “Multifunctorial Inverse K -theory”. In: *Ann. K-Theory* 7 no. 3 (2022), pp. 507–548.
- Michael A. Mandell. “An inverse K -theory functor”. In: *Doc. Math.* vol 15 (2010), pp. 765–791.
- J. Peter May. *The geometry of iterated loop spaces*. Lecture notes in Mathematics. Berlin, Heidelberg: Springer, 1972.
- J. Peter May. “ E_∞ spaces, group completions, and permutative categories”. In: *New developments in topology. The Edited and Revised Proceedings of the Symposium on Algebraic Topology, Oxford, June 1972*. Ed. by Graeme Segal. London Math. Soc. Lecture Note Ser. 11. London-New York: Cambridge University Press, 1974, pp. 61–93.
- J. Peter May. “The construction of E_∞ ring spaces from bipermutative categories”. In: *New topological contexts for Galois theory and algebraic geometry (BIRS 2008)*. Vol. 16. Geom. Topol. Monogr. Geom. Topol. Publ., Coventry, 2009, pp. 283–330.
- J. P. May and R. Thomason. “The uniqueness of infinite loop space machines”. In: *Topology* 17 no. 3 (1978), pp. 205–224.

- Graeme Segal. “Categories and cohomology theories”. In: *Topology* 13 (1974), pp. 293–312.
- R. W. Thomason. “**Cat** as a closed model category”. In: *Cahiers Topologie Géom. Différentielle* 21 no. 3 (1980), pp. 305–324.
- R. W. Thomason. “Symmetric monoidal categories model all connective spectra”. In: *Theory Appl. Categ.* 1 no. 3 (1995), pp. 78–118.
- Donald Yau. “Multifunctorial Equivariant Algebraic K -Theory”. Available at <https://arxiv.org/abs/2404.02794>. 2024.
- Donald Yau. *The Grothendieck Construction of Bipermutative-Indexed Categories and Pseudo Symmetric Inverse K-Theory*. CRC Press, Boca Raton, FL: Chapman & Hall/CRC, 2024.
- Inna Zakharevich. “The category of Waldhausen categories is a closed multicategory”. In: *New directions in homotopy theory*. Vol. 707. Contemp. Math. Providence, RI: Amer. Math. Soc., 2018, pp. 175–194.

Department of Mathematics, University of Western Ontario
2004 Perth Dr, London, ON, Canada, N6G 2V4
Email: dmanco@uwo.ca

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS L^AT_EX₂ε is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Geoff Cruttwell, Mount Allison University: gcruttwell@mta.ca

T_EXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT T_EX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

John Bourke, Masaryk University: bourkej@math.muni.cz

Maria Manuel Clementino, Universidade de Coimbra: mmc@mat.uc.pt

Valeria de Paiva, Topos Institute: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Rune Haugseng, Norwegian University of Science and Technology: rune.haug seng@ntnu.no

Dirk Hofmann, Universidade de Aveiro: dirk@ua.pt

Joachim Kock, Universitat Autònoma de Barcelona: Joachim.Kock@uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Sandra Mantovani, Università degli Studi di Milano: sandra.mantovani@unimi.it

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Giuseppe Metere, Università degli Studi di Palermo: giuseppe.metere@unipa.it

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

Jiri Rosický, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@unige.it

Michael Shulman, University of San Diego: shulman@sandiego.edu

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be

Christina Vasilakopoulou, National Technical University of Athens: cvasilak@math.ntua.gr